


Monirah A. Al-Alshaikh, Hazem A. Ghabbour*, Abdul-Malek S. Al-Tamimi, Mohammed S.M. Abdelbaky, Santiago Garcia-Granda and Ali A. El-Emam

Crystal structure of 5-(adamantan-1-yl)-3-[(4-chloroanilino)methyl]-2,3-dihydro-1,3,4-oxadiazole-2-thione, C₁₉H₂₂ClN₃OS



DOI 10.1515/ncls-2015-0151

Received November 26, 2015; accepted January 7, 2016; available online January 29, 2016

Abstract

C₁₉H₂₂ClN₃OS, orthorhombic, P2₁2₁2₁ (no. 19), $a = 7.0418(2)$ Å, $b = 10.8802(3)$ Å, $c = 23.5506(6)$ Å, $V = 1804.36(8)$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.0413$, $wR_{\text{ref}}(F^2) = 0.1110$, $T = 100$ K.

CCDC no.: 1405608

***Corresponding author:** Hazem A. Ghabbour, Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, e-mail: ghabbour@yahoo.com; and Department of Medicinal Chemistry, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

Monirah A. Al-Alshaikh: Department of Chemistry, College of Sciences, King Saud University, Riyadh 11451, Saudi Arabia

Abdul-Malek S. Al-Tamimi: Department of Pharmaceutical Chemistry, College of Pharmacy, Prince Sattam bin Abdulaziz University, Alkharij 11942, Saudi Arabia

Mohammed S.M. Abdelbaky and Santiago Garcia-Granda:

Departamento de Química Física y Analítica, Facultad de Química, Universidad de Oviedo – CINN, Av. Julián Clavería, 8, 33006 Oviedo, (Asturias), Spain

Ali A. El-Emam: Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia

The crystal structure is shown in the figure. Tables 1–3 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Colourless, prism, size 0.2231×0.5389×0.7274 mm
Wavelength:	CuK α radiation (1.54184 Å)
μ :	30.49 cm ⁻¹
Diffractometer, scan mode:	Xcalibur, Ruby, Gemini, ω scans
$2\theta_{\text{max}}$:	140.8°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	15818, 3444
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3373
$N(\text{param})_{\text{refined}}$:	231
Programs:	CrysAlis [17], SHELX [18]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	Site	x	y	z	U_{iso}
H(9A)	4a	0.5080	-0.0521	0.1901	0.031
H(9B)	4a	0.6646	-0.0449	0.1427	0.031
H(15)	4a	-0.3588	-0.1097	0.0216	0.037
H(10)	4a	0.8080	-0.0995	0.2301	0.033
H(8A)	4a	0.4589	0.2715	0.2156	0.033
H(8B)	4a	0.3809	0.1427	0.2350	0.033
H(18)	4a	0.1008	-0.3292	0.1189	0.041
H(4A)	4a	0.7917	0.1621	0.1110	0.036
H(4B)	4a	0.7132	0.2835	0.1386	0.036
H(13A)	4a	0.1514	0.0956	-0.0290	0.036
H(13B)	4a	0.2343	-0.0282	-0.0062	0.036
H(7)	4a	0.6037	0.2154	0.3032	0.035
H(16)	4a	-0.4431	-0.2822	0.0734	0.044
H(19)	4a	0.1838	-0.1522	0.0710	0.038
H(5)	4a	1.0117	0.2359	0.1789	0.043
H(6A)	4a	0.7915	0.3347	0.2400	0.043
H(6B)	4a	0.9234	0.2473	0.2756	0.043
H(12A)	4a	1.0509	0.0515	0.2310	0.042
H(12B)	4a	0.9987	0.0194	0.1678	0.042
H(11A)	4a	0.8000	0.0395	0.3070	0.035
H(11B)	4a	0.5919	-0.0003	0.2915	0.035
H(3)	4a	-0.133(4)	0.032(3)	-0.004(2)	0.05(1)

Table 3: Atomic displacement parameters (Å²).

Atom	Site	x	y	z	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
S(1)	4a	0.08526(8)	0.33243(6)	0.03351(3)	0.0228(3)	0.0306(3)	0.0473(3)	0.0020(2)	-0.0023(2)	0.0118(3)
Cl(1)	4a	-0.2398(1)	-0.45144(6)	0.14068(3)	0.0636(5)	0.0355(3)	0.0392(3)	-0.0125(3)	0.0101(3)	0.0039(2)
O(1)	4a	0.3371(2)	0.2459(2)	0.10883(7)	0.0238(8)	0.0236(8)	0.0351(9)	0.0009(7)	-0.0003(7)	0.0015(7)
N(2)	4a	0.2306(3)	0.1054(2)	0.05303(8)	0.0245(9)	0.0250(9)	0.0279(9)	-0.0030(8)	-0.0031(8)	0.0042(7)
N(1)	4a	0.3609(3)	0.0477(2)	0.08851(8)	0.0238(9)	0.0243(9)	0.0292(9)	0.0000(8)	0.0003(8)	0.0044(8)
C(2)	4a	0.4201(3)	0.1342(2)	0.12104(9)	0.019(1)	0.023(1)	0.030(1)	-0.0006(9)	0.0042(9)	0.0033(8)
N(3)	4a	-0.0360(3)	-0.0064(2)	0.0103(1)	0.026(1)	0.028(1)	0.043(1)	0.0018(9)	-0.0098(9)	0.0059(9)
C(14)	4a	-0.0788(4)	-0.1109(2)	0.04094(9)	0.029(1)	0.028(1)	0.028(1)	0.000(1)	-0.000(1)	-0.0050(9)
C(1)	4a	0.2174(3)	0.2252(2)	0.0643(1)	0.019(1)	0.029(1)	0.033(1)	-0.0022(9)	0.0034(9)	0.0068(9)
C(9)	4a	0.6193(3)	-0.0073(2)	0.1775(1)	0.025(1)	0.021(1)	0.033(1)	0.0008(9)	-0.0020(9)	-0.0027(9)
C(15)	4a	-0.2665(4)	-0.1524(2)	0.0418(1)	0.029(1)	0.028(1)	0.035(1)	0.003(1)	-0.001(1)	-0.009(1)
C(10)	4a	0.7746(3)	-0.0134(2)	0.2231(1)	0.029(1)	0.024(1)	0.033(1)	0.0010(9)	-0.002(1)	-0.0005(9)
C(3)	4a	0.5669(3)	0.1282(2)	0.16640(9)	0.019(1)	0.021(1)	0.031(1)	-0.0024(9)	0.0024(9)	0.0003(8)
C(8)	4a	0.4931(3)	0.1863(2)	0.2221(1)	0.024(1)	0.024(1)	0.035(1)	0.0023(9)	0.0016(9)	-0.0016(9)
C(18)	4a	0.0083(4)	-0.2835(2)	0.1002(1)	0.042(1)	0.030(1)	0.030(1)	0.000(1)	-0.002(1)	0.000(1)
C(4)	4a	0.7451(3)	0.1982(2)	0.1460(1)	0.023(1)	0.033(1)	0.035(1)	-0.008(1)	0.001(1)	0.0067(9)
C(13)	4a	0.1510(4)	0.0399(2)	0.0031(1)	0.031(1)	0.031(1)	0.030(1)	-0.002(1)	-0.0042(9)	0.004(1)
C(7)	4a	0.6497(4)	0.1787(2)	0.2677(1)	0.031(1)	0.028(1)	0.030(1)	-0.002(1)	-0.0008(9)	-0.005(1)
C(17)	4a	-0.1787(4)	-0.3207(2)	0.1017(1)	0.051(2)	0.026(1)	0.029(1)	-0.006(1)	0.009(1)	0.000(1)
C(16)	4a	-0.3174(4)	-0.2562(3)	0.0725(1)	0.032(1)	0.036(1)	0.040(1)	-0.007(1)	0.008(1)	-0.011(1)
C(19)	4a	0.0581(4)	-0.1785(2)	0.0709(1)	0.029(1)	0.033(1)	0.032(1)	-0.000(1)	-0.002(1)	0.001(1)
C(5)	4a	0.8989(4)	0.1911(3)	0.1918(1)	0.024(1)	0.042(2)	0.041(1)	-0.014(1)	-0.002(1)	0.009(1)
C(6)	4a	0.8249(4)	0.2494(3)	0.2468(1)	0.036(1)	0.030(1)	0.041(1)	-0.014(1)	-0.011(1)	0.002(1)
C(12)	4a	0.9516(3)	0.0563(3)	0.2025(1)	0.019(1)	0.050(2)	0.037(1)	0.001(1)	0.001(1)	-0.001(1)
C(11)	4a	0.7023(3)	0.0444(2)	0.2781(1)	0.027(1)	0.030(1)	0.030(1)	-0.003(1)	0.0000(9)	-0.001(1)

Source of material

4-Chloroaniline (1.28 g, 0.01 mol) and 37% formaldehyde solution (1.5 mL) were added to a solution of 5-(1-adamantyl)-1,3,4-oxadiazoline-2-thione (2.36 g, 0.01 mol) in ethanol (15 mL), and the mixture was stirred at room temperature for 2 hours and allowed to stand overnight. The precipitated crude product was filtered, washed with cold ethanol, dried, and crystallized from ethanol to yield 3.46 g (92%) of the title compound. M.p. 450–452 K [9]. Colourless prismatic single crystals were obtained by slow evaporation of chloroform-ethanol solution (1:1) at room temperature. ¹H NMR (CDCl₃, 500.13 MHz): δ 1.68 (q, 6H, adamantane-CH₂), 1.93 (s, 6H, adamantine-CH₂), 2.07 (s, 3H, adamantane-CH), 5.06 (t, 1H, NH), 5.36 (d, 2H, CH₂), 6.80 (d, 2H, Ar—H, *J* = 7.0 Hz), 7.14 (d, 2H, Ar—H, *J* = 7.0 Hz). ¹³C NMR (CDCl₃, 125.76 MHz): δ 27.30, 34.35, 35.91, 38.94 (adamantane-C), 58.22 (CH₂), 115.30, 124.46, 129.10, 142.75 (Ar—C), 168.20 (C = N), 176.77 (C = S). EI-MS, *m/z* (rel. int.): 377 (M⁺ + 2, 1), 359 (M⁺, 4), 236 (82), 176 (17), 142 (20), 140 (55), 135 (100), 113 (5), 111 (26).

Experimental details

The structure was refined as an inversion twin (BASF 0.43).

Discussion

The adamantyl group was identified as an essential pharmacophore in several bioactive drugs [1, 2]. Several adamantane derivatives are currently used as efficient therapies against

influenza [3–5], herpes simplex [6] and HIV [7–9] viruses. In addition, numerous adamantane derivatives were reported to exhibit marked anticancer [10, 11], antibacterial [12–15] and antifungal [16] activities. In continuation to our ongoing interest in the chemical and biological properties of adamantane derivatives, we report herein the crystal structure of the title compound which displayed potent broad spectrum antibacterial activity [9]. The title compound is a functionalized 1,3,4-oxadiazole-2-thione derivative. The benzene ring makes a dihedral angle of 57.43(5)^o with the 1,3,4-oxadiazole-2-thione. In the crystal, the N3—H3 · · · S1 hydrogen-bonding interactions link the molecules into chains extending along the c direction. The distance of N and H3 is 2.57(4) Å and the angle is 171.0 (3)^o.

Acknowledgements: The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for funding this work through the Research Group Project No. PRG-1436-23. We also acknowledge the financial support from Spanish Ministerio de Economía y Competitividad (MINECO-13-MAT2013-40950-R, FPI grant BES-2011-046948 to MSM-A).

References

- Lamoureux, G.; Artavia, G.: Use of the adamantane structure in medicinal chemistry. *Curr. Med. Chem.* **17** (2010) 2967–2978.

2. Liu, J.; Obando, D.; Liao, V.; Lifa, T.; Codd, R.: The many faces of the adamantyl group in drug design. *Eur. J. Med. Chem.* **46** (2011) 1949–1963.
3. Davies, W. L.; Grunnert, R. R.; Haff, R. F.; McGahen, J. W.; Neumeyer, E. M.; Paulshock, M.; Watts, J. C.; Wood, T. R.; Hermann, E. C.; Hoffmann, C. E.: Antiviral activity of 1-adamantamine (amantadine). *Science* **144** (1964) 862–863.
4. Rabinovich, S.; Baldini, J. T.; Bannister, R.: Treatment of influenza. The therapeutic efficacy of rimantadine HCl in a naturally occurring influenza A2 outbreak. *Am. J. Med. Sci.* **257** (1969) 328–335.
5. Hayden, F. G.; Gwaltney, J. M. I.; Van, C. R. L.; Adams, K. F.; Giordani, B.: Comparative toxicity of amantadine hydrochloride and rimantadine hydrochloride in healthy adults. *Antimicrob. Agents Chemother.* **19** (1981) 226–233.
6. Rosenthal, K. S.; Sokol, M. S.; Ingram, R. L.; Subramanian, R.; Fort, R. C.: Tromantadine: inhibitor of early and late events in herpes simplex virus replication. *Antimicrob. Agents Chemother.* **22** (1982) 1031–1036.
7. Burstein, M. E.; Serbin, A. V.; Khakhulina, T. V.; Alymova, I. V.; Stotskaya, L. L.; Bogdan, O. P.; Manukchina, E. E.; Idanov, V. V.; Sharova, N. K.: Inhibition of HIV-1 replication by newly developed adamantane-containing polyanionic agents. *Antiviral Res.* **41** (1999) 135–144.
8. Balzarini, J.; Orzeszko, B.; Mauri, J. K.; Orzeszko, A.: Synthesis and anti-HIV studies of 2-adamantyl-substituted thiazolidin-4-ones. *Eur. J. Med. Chem.* **42** (2007) 993–1003.
9. El-Emam, A. A.; Al-Deeb, O. A.; Al-Omar, M. A.; Lehmann, J.: Synthesis, antimicrobial, and anti-HIV-1 activity of certain 5-(1-adamantyl)-2-substituted thio-1,3,4-oxadiazoles and 5-(1-adamantyl)-3-substituted aminomethyl-1,3,4-oxadiazoline-2-thiones. *Bioorg. Med. Chem.* **12** (2004) 5107–5113.
10. Sun, S. Y.; Yue, P.; Chen, X.; Hong, W. K.; Lotan, R.: The synthetic retinoid CD437 selectively induces apoptosis in human lung cancer cells while sparing normal human lung epithelial cells. *Cancer Res.* **62** (2002) 2430–2436.
11. Lorenzo, P.; Alvarez, R.; Ortiz, M. A.; Alvarez, S.; Piedrafita, F. J.; de Lira, Á. R. Inhibition of IκB kinase-β and anticancer activities of novel chalcone adamantyl carotenoids. *J. Med. Chem.* **51** (2008) 5431–5440.
12. Jia, L.; Tomaszewski, J. E.; Hanrahan, C.; Coward, L.; Noker, P.; Gorman, G.; Nikonenko, B.; Protopopova, M.: Pharmacodynamics and pharmacokinetics of SQ109, a new diamine-based antitubercular drug. *Brit. J. Pharmacol.* **144** (2005) 80–87.
13. Al-Abdullah, E. S.; Asiri, H. H.; Lahsasni, S.; Habib, E. E.; Ibrahim, T. M.; El-Emam, A. A.: Synthesis, antimicrobial, and anti-inflammatory activity, of novel *S*-substituted and *N*-substituted 5-(1-adamantyl)-1,2,4-triazole-3-thiols. *Drug Des. Dev. Ther.* **8** (2014) 505–518.
14. Kadi, A. A.; El-Brolosy, N. R.; Al-Deeb, O. A.; Habib, E. E.; Ibrahim, T. M.; El-Emam, A. A. Synthesis, antimicrobial, and anti-inflammatory activities of novel 2-(1-adamantyl)-5-substituted-1,3,4-oxadiazoles and 2-(1-adamantylamino)-5-substituted-1,3,4-thiadiazoles. *Eur. J. Med. Chem.* **42** (2007) 235–242.
15. Kadi, A. A.; Al-Abdullah, E. S.; Shehata, I. A.; Habib, E. E.; Ibrahim, T. M.; El-Emam, A. A. Synthesis, antimicrobial and anti-inflammatory activities of novel 5-(1-adamantyl)-1,3,4-thiadiazole Derivatives. *Eur. J. Med. Chem.* **45** (2010) 5006–5011.
16. Omar, K.; Geronikaki, A.; Zoumpoulakis, P.; Camoutsis, C.; Soković, M.; Čirić, A.; Glamočlija, J. Novel 4-thiazolidinone derivatives as potential antifungal and antibacterial drugs. *Bioorg. Med. Chem.* **18** (2010) 426–432.
17. Agilent. CrysAlis PRO. Agilent Technologies UK Ltd, Yarnton, England, 2014.
18. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.