

Doubled aspects of generalised dualities and integrable deformations

Saskia Demulder,^{a,b} Falk Hassler^{c,d,e} and Daniel C. Thompson^{a,b}

^a*Theoretische Natuurkunde, Vrije Universiteit Brussel & The International Solvay Institutes, B-1050 Brussels, Belgium*

^b*Department of Physics, Swansea University, Swansea, SA2 8PP, U.K.*

^c*Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC 27599-3255, U.S.A.*

^d*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, U.S.A.*

^e*Department of Physics, University of Oviedo, Avda. Federico García Lorca 18, 33007 Oviedo, Spain*

E-mail: Saskia.Demulder@vub.be, falk@fhassler.de, D.C.Thompson@Swansea.ac.uk

ABSTRACT: The worldsheet theories that describe Poisson-Lie T-dualisable σ -models on group manifolds as well as integrable η , λ and β -deformations provide examples of \mathcal{E} -models. Here we show how such \mathcal{E} -models can be given an elegant target space description within Double Field Theory by specifying explicitly generalised frame fields forming an algebra under the generalised Lie derivative. With this framework we can extract simple criteria for the R/R fields and the dilaton that extend the \mathcal{E} -model conditions to type II backgrounds. In particular this gives conditions for a type II background to be Poisson-Lie T-dualisable. Our approach gives rise to algebraic field equations for Poisson-Lie symmetric spacetimes and provides an effective tool for their study.

KEYWORDS: Flux compactifications, Space-Time Symmetries, String Duality

ARXIV EPRINT: [1810.11446](https://arxiv.org/abs/1810.11446)

Contents

1	Introduction	1
2	\mathcal{E}-models: Poisson-Lie duality and integrable theories	3
2.1	Poisson-Lie models	4
2.2	Integrable deformations	6
2.2.1	η -deformation	6
2.2.2	λ -deformations	7
3	Target space description of \mathcal{E}-models	8
3.1	A brief review of DFT_{wzw}	8
3.2	The generalised frame fields	11
3.3	Equivalence to (modified) type II supergravity	15
3.3.1	Unimodular case	16
3.3.2	Non-unimodular case	16
3.3.3	R/R sector	18
4	The \mathcal{E}-model conditions	19
4.1	NS sector	20
4.2	Dilaton	20
4.3	R/R sector	21
4.4	Fourier-Mukai transformation	21
5	Application to integrable deformations	24
5.1	Deformations based on the (m)CYBE	24
5.2	η -deformation redux	29
5.3	λ -deformation redux	31
6	Conclusion	35
A	Conventions and notation	36
B	Algebraic structures	38
C	Solution to the section condition	41
D	Fluxes in the generalised parallelizable frame	43
E	Dilaton in PL and λ models	44
F	Details of η-supergravity solution	45
G	Drinfel'd doubles and group parameterisations	47
G.1	$\mathfrak{su}(2) \oplus \mathfrak{e}_3$	47
G.2	$\mathfrak{su}(1,1) \oplus \mathfrak{e}_3$	48

1 Introduction

Abelian T-duality, the invariance of string theory when the radius of an S^1 in target space is inverted, has long served as a catalyst for theoretical developments. Given its prominent role, a long standing challenge has been to establish T-duality in more general contexts for instance when the target space admits a non-Abelian group of isometries [1]. Two decades ago, in a remarkable sequence of works [2, 3] by Klimčík and Ševera it was shown that in special circumstances one may even relax the imposition of an isometry in target space and still retain a notion of T-duality called Poisson-Lie (PL) duality.

Some caution should be exercised here; whilst the maps between non-linear sigma-models induced by non-Abelian [4] or more generally PL T-dualities [2, 3, 5, 6] are canonical transformations of the classical phase space, it is hard in general to establish them as fully fledged quantum equivalences. Indeed in a generic context one should not expect to have control of either α' or g_s effects. Optimistically one might suggest that these “dualities” constitute a map from a CFT to a new CFT' for which modular invariance may necessitate the inclusion of extra twisted sectors. The partition sums of these theories need not match. This viewpoint dates back to [7] and was recently shown to be the case in a simple $SU(2)$ non-Abelian T-dualisation [8].

Nonetheless these generalised “dualities” (and henceforth we drop the quotation marks) retain utility as solution generating techniques within supergravity and continue then to hold interest for their potential application to holography. Non-Abelian T-duality for instance can be used to construct novel examples of holographic spacetimes (see e.g. [9–11] for early works in this direction and [12] for the field theoretic interpretation). Poisson-Lie T-duality at first sight appears to concern rather complicated looking spacetimes. However this complexity can in some cases be illusory. In fact a class of integrable models, known variously as η -deformations or Yang-Baxter sigma-models, introduced by Klimčík [13] some years ago constitute exemplars of PL T-dualisable theories. A significant amount of activity has followed from the introduction of the integrable η -deformation of the $AdS_5 \times S^5$ spacetime [14, 15]. A further development has been integrable λ -deformations [16, 17] of (potentially gauged) WZW-models which are related to η -type deformations [18–20] via a Poisson-Lie duality transformation combined with an analytic continuation of certain Euler angles and couplings.

From the worldsheet perspective such generalised dualities can be rendered manifest in a *doubled formalism* much like that of Abelian T-duality introduced in [21, 22]. In these approaches one considers a sigma-model whose target space has double the number of dimensions. Half of the coordinates describing this doubled space can be eliminated to recover a standard sigma-model.¹ When this reduction can be done in multiple ways we recover T-dual related descriptions.

This philosophy was extended to generalised T-dualities in the original works [2, 3] as well as in [26]. Other recent interesting works in this direction include [27, 28]. The

¹This reduction requires the imposition of a chirality constraint which is a delicate matter quantum mechanically and in [21, 22] it is achieved at the expense of manifest Lorentz invariance, other alternatives based on gauging e.g. [23–25] may prove more amenable to a quantum treatment.

doubled space is equipped with the familiar generalised metric and $O(D, D)$ invariant inner-product and is further required to be a group manifold, \mathbb{D} , and so comes equipped with a canonical three-form. A useful presentation of the doubled worldsheet is provided by the first order formalism now coined \mathcal{E} -models [29], first introduced in [3] and developed in [30]. PL dualisable sigma models, as well as η - and λ - and β - deformations are all examples of theories that can be extracted from \mathcal{E} -models.

Abelian T-duality has an elegant target space duality symmetric formulation known as Double Field Theory (DFT) [31]. Since a worldsheet doubled formalism is available for generalised T-dualities one would hope for a similar understanding at the level of the target space. The first clues here come from studying the one-loop β -functions of the worldsheet theory [32, 33]. In [32] it was pointed out that the β -function for the generalised metric corresponds to the scalar equation of motion of a gauged supergravity with the structure constants of the doubled target space providing the embedding tensor. In DFT the way such gauged supergravities arise is by performing a Scherk-Schwarz reduction [34–40]. Thus what one requires is a precise formulation of DFT on the group manifold \mathbb{D} . DFT_{WZW} [41–43] provides exactly such an approach and the study of its relation to Poisson-Lie T-duality was initiated [44].

This manuscript will continue the development of generalised T-dualities and integrable deformations within DFT. Specifically we will show how the type II extension of DFT_{WZW} provides an immediate set of criteria that extends the structure of \mathcal{E} -models to the R/R and dilaton sector. In the case where this \mathcal{E} -model describes a PL T-dualisable NS sector, this gives rise to criteria that must be obeyed for a full type II supergravity background to be PL T-dualisable. We shall describe backgrounds for which these criteria hold as being PL symmetric. DFT_{WZW} makes this symmetry manifest and thereby significantly simplifies their analysis. For example, instead of having to cope with difficult, coupled PDEs, the field equations become algebraic.

A pivotal element in our discussion will be a generalised frame field on the spacetime which allows us to connect the fields on the doubled space with the conventional type II supergravity fields. In this work we will follow a technique suggested in [45] to construct a set of $O(D, D)$ valued generalised frame fields that furnish the algebra of \mathbb{D} via the generalised Lie derivative. In the cases we are most interested in, and that includes η -, λ - and β -deformations as well as all PL dualisable models, this construction is carried out explicitly making use of the group theoretic quantities on \mathbb{D} and its coset $\mathcal{M} = \mathbb{D}/\tilde{H}$ by a maximal isotropic subgroup \tilde{H} . Our discussion will be predominantly local in nature, however where this construction can be extended globally this provides an understanding of \mathcal{E} -models as examples of generalised parallelizable spaces [46, 47].

The η - and some β -deformations are governed by modified type II field equations [48–50]. Modified type II requires a Killing vector, I , and a one form, Z , in addition to the bosonic field content known from standard type II supergravity. Connections to DFT and ExFT of modified supergravity are discussed in [51, 52], here we show that they also arise from DFT_{WZW} if the subgroup \tilde{H} is non-unimodular. In [53–55] an open string interpretation of such I modified supergravities and integrable deformations was given. We will illustrate these ideas with a number of specific examples. They emphasize how exploit-

ing PL symmetry can make challenging calculations in integral deformations much easier and vindicate the combination of DFT techniques and integrable deformations.

The paper is organised as follows: in section 2 we review \mathcal{E} -models and how the λ - and η -deformations fit into this framework. In section 3, we develop further the implementation of Poisson-Lie T-duality in Double Field Theory [44] to show how the R/R-sector of (modified-) SUGRA can be elegantly extracted. After a short reminder of DFT_{WZW} , we prove how for a group admitting a maximally isotropic subgroup \tilde{H} together with some additional conditions there exists a generalised frame field solving the section condition of DFT. Then in section 4 we discuss how the DFT manifest implementation of Poisson-Lie T-duality can be extended to the R/R-sector and dilaton. In the last section 5 we apply the formalism to integrable deformations and provide some explicit examples of how the R/R-sector can be extracted for e.g. $\text{AdS}_3 \times S^3$ backgrounds. The goal of section 5 there is not to present novel backgrounds but to demonstrate the efficacy of the approach proposed in the paper. We conclude with a brief discussion of some of the outstanding challenges as we see them. The presentation is complemented with a number of technical appendices.

Note added. Whilst this manuscript was in its very final stages of preparation we received a preprint [56] from `math.DG` that overlaps with some of the conclusions of this paper, albeit cast in the language of Courant Algebroids rather than DFT_{WZW} .

2 \mathcal{E} -models: Poisson-Lie duality and integrable theories

To make this article self-contained, let us begin by reviewing the basic features of \mathcal{E} -models before describing the specialisation to Poisson-Lie T-duality and integrable deformations.

Our starting point² is a real Lie algebra \mathfrak{d} of even dimension, $\dim \mathfrak{d} = 2D$, equipped with non-degenerate, ad-invariant, symmetric inner-product $\langle\langle \cdot, \cdot \rangle\rangle$ that we assume to be of split-signature. Letting \mathbb{T}_A be a basis of generators for \mathfrak{d} , we shall write

$$[\mathbb{T}_A, \mathbb{T}_B] = \mathbb{F}_{AB}{}^C \mathbb{T}_C, \quad \langle\langle \mathbb{T}_A, \mathbb{T}_B \rangle\rangle = \eta_{AB}. \quad (2.1)$$

We denote the components of the matrix inverse of η_{AB} as η^{AB} and we will raise and lower indices with this.

The \mathcal{E} -model is a dynamical system that can be conveniently parametrised by a set of algebra-valued maps $j = j^A \mathbb{T}_A : S_\sigma^1 \rightarrow \mathfrak{d}$ obeying the classical current algebra

$$\{j_A(\sigma), j_B(\sigma')\}_{P.B.} = \mathbb{F}_{AB}{}^C j_C(\sigma) \delta(\sigma - \sigma') + \eta_{AB} \delta'(\sigma - \sigma'), \quad (2.2)$$

with dynamics determined by the Hamiltonian

$$\text{Ham} = \frac{1}{2} \oint d\sigma \langle\langle j(\sigma), \mathcal{E}(j(\sigma)) \rangle\rangle. \quad (2.3)$$

Here $\mathcal{E}(\mathbb{T}_A) = \mathcal{E}_A{}^B \mathbb{T}_B$, the eponymous operator, is an idempotent involution of \mathfrak{d} that is self-adjoint with respect to $\langle\langle \cdot, \cdot \rangle\rangle$. We can parametrise \mathcal{E} in terms of a *generalised metric*, \mathcal{H} , as

$$\mathcal{E}_A{}^B = \mathcal{H}_{AC} \eta^{CB}, \quad \mathcal{H}_{AB} = \mathcal{H}_{BA}, \quad \mathcal{H}_{AC} \eta^{CD} \mathcal{H}_{DB} = \eta_{AB}. \quad (2.4)$$

²A guide to notation, conventions and some algebra terminology can be found in appendix A.

We will be interested in the case, and assume it henceforth, that \mathfrak{d} admits a maximally isotropic subalgebra $\tilde{\mathfrak{h}} \subset \mathfrak{d}$. Then the \mathcal{E} -model can be reduced to a conventional non-linear sigma-model whose target space is the coset $\mathcal{M} = \mathbb{D}/\tilde{H}$ where \mathbb{D}, \tilde{H} are the groups corresponding to respectively $\mathfrak{d}, \tilde{\mathfrak{h}}$. To fix notation we let $T_A = (\tilde{T}^a, T_a)$ where \tilde{T}^a are generators of $\tilde{\mathfrak{h}}$ and T_a are the remaining generators whose span we denote \mathfrak{k} . In this basis the inner-product can be taken to be

$$\eta_{AB} = \begin{pmatrix} 0 & \delta^a_b \\ \delta_a^b & 0 \end{pmatrix}. \tag{2.5}$$

It is important to stress that in the decomposition $\mathfrak{d} = \tilde{\mathfrak{h}} \oplus \mathfrak{k}$ we place no requirement on \mathfrak{k} ; that is to say in general \mathbb{D}/\tilde{H} is neither a group manifold itself nor a symmetric space however the examples we shall be most interested in here will indeed be of this type.

The non-linear sigma-model that follows from the \mathcal{E} -model is described by an action [3, 29]³

$$S_{\mathbb{D}/\tilde{H}} = \tilde{k} S_{\text{WZW}}[m] - \frac{\tilde{k}}{\pi} \int d\sigma d\tau \langle\langle \mathcal{P}(m^{-1}\partial_+m), m^{-1}\partial_-m \rangle\rangle, \tag{2.6}$$

$$S_{\text{WZW}}[m] = \frac{1}{2\pi} \int d\sigma d\tau \langle\langle m^{-1}\partial_+m, m^{-1}\partial_-m \rangle\rangle + \frac{1}{24\pi} \int_{M_3} \langle\langle m^{-1}dm, [m^{-1}dm, m^{-1}dm] \rangle\rangle. \tag{2.7}$$

Here we have parametrised a group element on \mathbb{D} as $g(X^I) = \tilde{h}(\tilde{x}_{\tilde{i}})m(x^i)$ where $X^I = (\tilde{x}_{\tilde{i}}, x^i)$ are local coordinates on \mathbb{D} such that $\tilde{x}_{\tilde{i}}, \tilde{i} = 1 \dots D$, are local coordinates on $\tilde{H} \subset \mathbb{D}$ and $x^i, i = 1 \dots d$, are local coordinates that parametrise the coset. The first term in eq. (2.6) denotes the WZW action on \mathbb{D} , defined with the inner-product $\langle\langle \cdot, \cdot \rangle\rangle$, evaluated on the coset representative m . The second term, whose coefficient is -2 times that of the kinetic term of the WZW model, is defined with a projector obeying [29]

$$\text{Im } \mathcal{P} = \tilde{\mathfrak{h}}, \quad \text{Ker } \mathcal{P} = (1 + \text{ad}_m \cdot \mathcal{E} \cdot \text{ad}_{m^{-1}})\mathfrak{d}. \tag{2.8}$$

2.1 Poisson-Lie models

Let us now discuss the special case where \mathfrak{d} is a Drinfel'd double i.e. $\mathfrak{d} = \tilde{\mathfrak{h}} \oplus \mathfrak{h}$ with *both* $\tilde{\mathfrak{h}}, \mathfrak{h}$ maximally isotropic subalgebras. This is the setting of Poisson-Lie T-duality. In this case we can identify the coset with the Lie group manifold $\mathbb{D}/\tilde{H} \cong H$ and so in the action eq. (2.6) the representative $m(x)$ can be considered an element of the group H . Since $m^{-1}dm$ is valued in \mathfrak{h} , which is an isotropic with respect to $\langle\langle \cdot, \cdot \rangle\rangle$, the WZW part of the action eq. (2.6) is identically zero and what remains can be cast in the form of a sigma-model:

$$S_{\mathbb{D}/\tilde{H}} = \frac{1}{\pi s} \int d^2\sigma e_+^a (E_0^{-1} + \Pi)_{ab}^{-1} e_-^b = \frac{1}{\pi s} \int d^2\sigma (G(x) - B(x))_{ij} \partial_+ x^i \partial_- x^j, \tag{2.9}$$

³Here we restore an overall normalisation $\frac{\tilde{k}}{2\pi}$, which depending on the specific properties of \mathbb{D}/\tilde{H} , may require a quantisation in order to define the WZ term unambiguously in a path integral.

in which $m^{-1}\partial_{\pm}m = e_{\pm}^a T_a = e^a{}_i \partial_{\pm} x^i T_a$ are the light cone components of the left-invariant one-forms pulled back to the worldsheet and the normalisation is $s = \tilde{k}^{-1}$. Later we shall also require the right-invariant one-forms $\partial_{\pm}mm^{-1} = v_{\pm}^a T_a = v^a{}_i \partial_{\pm} x^i T_a$ together with the vector fields $e_a = e^i{}_a \partial_i$ and $v_a = v^i{}_a \partial_i$ that generate respectively right and left actions. The matrix Π is derived from the adjoint action:

$$\text{ad}_m \mathbb{T}_A = m \mathbb{T}_A m^{-1} = M_A{}^B \mathbb{T}_B, \quad \Pi^{ab} = M^{ac} M^b{}_c. \quad (2.10)$$

The D^2 constant parameters in $E_0 = G_0 - B_0$ are related to those of the generalised metric introduced in eq. (2.4) in the standard way

$$\mathcal{H}_{AB} = \begin{pmatrix} G_0^{-1} & -G_0^{-1} B_0 \\ B_0 G_0^{-1} & G_0 - B_0 G_0^{-1} B_0 \end{pmatrix}. \quad (2.11)$$

In general the target space metric corresponding to the sigma model eq. (2.9) is unappetising and lacking isometry however it has a rather special algebraic structure. Although the currents J_a corresponding to left action on H are not conserved in the usual sense they do obey a non-commutative conservation law

$$\partial_+ J_{a-} + \partial_- J_{a+} = \tilde{F}^{bc}{}_a J_{b+} J_{c-}, \quad (2.12)$$

in which we emphasise that the structure constants appearing on the right hand side are those of the $\tilde{\mathfrak{h}}$. In terms of the target space data, $E_{ij} = G_{ij} - B_{ij}$, this places a requirement that

$$L_{V_a} E_{ij} = -\tilde{F}^{bc}{}_a e^k{}_b e^l{}_c E_{ik} E_{lj}. \quad (2.13)$$

This condition on the target space is referred to as a Poisson-Lie symmetry.⁴

At this stage we make an important observation; using the curved space G_{ij} and B_{ij} that define the sigma model eq. (2.9) we may define a coordinate dependent $O(D, D)$ generalised metric

$$\hat{\mathcal{H}}_{\hat{i}\hat{j}}(x) = \begin{pmatrix} G^{-1} & -G^{-1}B \\ BG^{-1} & G - BG^{-1}B \end{pmatrix}_{\hat{i}\hat{j}}. \quad (2.14)$$

A tedious but straightforward calculation reveals that

$$\hat{\mathcal{H}}_{\hat{i}\hat{j}}(x) = \hat{E}^A{}_{\hat{i}}(x) \mathcal{H}_{AB} \hat{E}^B{}_{\hat{j}}(x), \quad (2.15)$$

where

$$\hat{E}^A{}_{\hat{i}}(x) = \begin{pmatrix} 1 & 0 \\ \Pi & 1 \end{pmatrix}^A{}_B \begin{pmatrix} e^{-T} & 0 \\ 0 & e \end{pmatrix}^B{}_i = M^A{}_B \begin{pmatrix} v^{-T} & 0 \\ 0 & v \end{pmatrix}^B{}_i. \quad (2.16)$$

The hats on the indices and frame fields are introduced to emphasize dependence only on the coordinates x^i and not on the ‘‘dual’’ $\tilde{x}_{\hat{i}}$, i.e. $\partial_{\hat{i}} = (0, \partial_i)$ — in the terminology of DFT

⁴Taking a further Lie derivative of this relation invokes an integrability condition, namely that viewed as a map $\mathfrak{h} \rightarrow \mathfrak{h} \wedge \mathfrak{h}$ the structure constants $\tilde{F}^{bc}{}_a$ are required to define a one-cocycle obeying the co-Jacobi identity. As explained in the appendix B, this property can be understood as the infinitesimal version of H being a Poisson-Lie group, giving justification for the name.

we have picked a solution to the section condition. Notice also that the frame fields are (coordinate dependent) elements of $O(D, D)$.

Of course we could swap the rôle of the two subgroups. If instead we parametrize $\mathfrak{g}(X) = \tilde{m}(\tilde{x})h(x)$ we can reduce to a theory on the coset $\tilde{\mathcal{M}} = \mathbb{D}/H \cong \tilde{H}$. In that case we find the Poisson-Lie T-dual theory to eq. (2.9) given by an action

$$S_{\mathbb{D}/H} = \frac{1}{\pi s} \int d^2\sigma \tilde{e}_{+a} \left(E_0 + \tilde{\Pi} \right)^{-1 ab} \tilde{e}_{-b} = \frac{1}{\pi s} \int d^2\sigma (\tilde{G} - \tilde{B})^{\tilde{i}\tilde{j}} \partial_+ \tilde{x}_{\tilde{i}} \partial_- \tilde{x}_{\tilde{j}}, \quad (2.17)$$

where $\tilde{m}^{-1} \partial_{\pm} \tilde{m} = \tilde{e}_{\pm a} \tilde{T}^a = \tilde{e}_a^{\tilde{i}} \partial_{\pm} \tilde{x}_{\tilde{i}} \tilde{T}^a$ and $\tilde{\Pi}$ defined via the adjoint action. An important feature is that the two PL sigma-models are related by a canonical transformation [2, 3, 5, 6] at the classical level which can be derived from a generating functional

$$F = \oint d\sigma \theta(\tilde{x}, x), \quad (2.18)$$

in which θ is the pull back of a one form to S_{σ}^1 whose form is known only implicitly. However an elegant expression can be given for its derivative [6]

$$\omega = d\theta = 2(\mathcal{O}^{-1})_a^b e^a \wedge \tilde{e}_b + (\mathcal{O}^{-1} \tilde{\Pi})_{ab} e^a \wedge e^b - (\mathcal{O}^{-T} \Pi)^{ab} \tilde{e}_a \wedge \tilde{e}_b, \quad \mathcal{O} = \text{id} - \tilde{\Pi} \Pi. \quad (2.19)$$

2.2 Integrable deformations

An application of \mathcal{E} -models is to provide a universal description of two superficial distinct classes of integrable deformations known as η - and λ -deformed theories [29]. Let us review some salient features of these deformations which we shall return to in some detail later.

2.2.1 η -deformation

In its simplest form the η -model is a deformation of the principal chiral model on a group manifold G defined in terms of an operator \mathcal{R} , an endomorphism of \mathfrak{g} obeying the modified classical Yang-Baxter equation

$$[\mathcal{R}X, \mathcal{R}Y] - \mathcal{R}([\mathcal{R}X, Y] + [X, \mathcal{R}Y]) = -c^2 [X, Y], \quad \forall X, Y \in \mathfrak{g}, \quad (2.20)$$

where $c^2 \in \{-1, 0, 1\}$. We require that \mathcal{R} be skew-symmetric with respect to the Cartan-Killing form $\langle t_a, t_b \rangle = \kappa_{ab} = -\frac{1}{2h\nu} f_{ac}^d f_{bd}^c$ with $[t_a, t_b] = f_{ab}^c t_c$ with $\{t_a\}$ the generators of \mathfrak{g} .

The η -deformation corresponds to taking the choice $c^2 = -1$ with \mathcal{R} acting to swap positive and negative roots and as zero on the Cartan and is defined by the action

$$S_{\eta} = \frac{1}{\pi t} \int d^2\sigma \langle v_+, (1 - \eta \mathcal{R})^{-1} v_- \rangle. \quad (2.21)$$

This theory is of particular interest since it preserves the integrability [57] of the principal chiral model (at least classically).⁵ What may not be immediately obvious is that this is

⁵In actuality, integrability of theory in eq. (2.21) is ensured for any value of $c^2 \in \{-1, 0, 1\}$ [58, 59] and the case of $c = 0$ is of relevance in describing e.g. TsT deformations [60, 61].

an example of a model admitting Poisson-Lie T-duality and thus an \mathcal{E} -model. Indeed the action (2.21) can be brought into PL form of eq. (2.9) with the identification

$$(E_0^{-1})^{ab} = \frac{\kappa^{ab}}{\eta} - R^{ab}, \quad \Pi^{ab} = R^{ab} - D[g]^{ac} R_c^d D[g^{-1}]_d^b, \quad s = \tilde{k}^{-1} = t\eta, \quad (2.22)$$

in which we have defined $\mathcal{R}(t_a) = R_a^b t_b$ and $\text{ad}_g t_a = g t_a g^{-1} = D[g]_a^b t_b$ for $g \in G$, and raised indices with κ^{ab} . To understand the \mathcal{E} -model corresponding to this sigma-model, one needs to identify the corresponding double \mathfrak{d} and idempotent operator \mathcal{E} . Note that eq. (2.20) ensures that the bracket

$$[X, Y]_{\mathcal{R}} = [\mathcal{R}X, Y] + [X, \mathcal{R}Y], \quad (2.23)$$

obeys the Jacobi identity and thus defines a second Lie-algebra we call $\mathfrak{g}_{\mathcal{R}}$. It is a standard result that $\mathfrak{d} = \mathfrak{g} + \mathfrak{g}_{\mathcal{R}}$ can be identified with the complexification $\mathfrak{d} = \mathfrak{g}^{\mathbb{C}}$ which, when viewed as a real Lie algebra with elements $Z = X + iY$, $X, Y \in \mathfrak{g}$, can be equipped with an inner-product given by the imaginary part of the Cartan-Killing form. Under the Iwasawa decomposition $\mathfrak{d} = \mathfrak{g}^{\mathbb{C}} = \mathfrak{g} + (\mathfrak{a} + \mathfrak{n})$ both \mathfrak{g} and $\tilde{\mathfrak{h}} = \mathfrak{a} + \mathfrak{n}$ are maximal isotropic subalgebras. Finally the operator \mathcal{E} is given by [29]

$$\mathcal{E} : Z \mapsto \frac{i}{2} (\eta - \eta^{-1}) Z - \frac{i}{2} (\eta + \eta^{-1}) Z^{\dagger}.$$

2.2.2 λ -deformations

Appearing at first sight to be a rather different class of integrable models, λ -deformations can be thought of as a re-summed marginally relevant current-current perturbation of a WZW-model on a group manifold G . The λ -deformed WZW model is specified by the action [16]

$$S_{\lambda} = k S_{WZW}[g] + \frac{k}{\pi} \int d^2 \sigma \langle \partial_+ g g^{-1}, (\lambda^{-1} - \text{ad}_{g^{-1}})^{-1} g^{-1} \partial_- g \rangle. \quad (2.24)$$

Here we use the WZW action for a group element $g \in G$ as in eq. (2.6) but with the inner-product simply given by the Cartan-Killing form, $\kappa = \langle \cdot, \cdot \rangle$. In addition to the metric and B -field obtained from the above action, the construction of the λ -deformed theory [16] requires a Gaussian elimination of fields which when performed in a path integral gives rise to a dilaton

$$\phi_{\lambda} = \phi_0 - \frac{1}{2} \log \det(1 - \lambda \text{ad}_{g^{-1}}), \quad (2.25)$$

in which ϕ_0 is constant. The λ -deformation can be recast into an \mathcal{E} -model for which $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{g}$, whose elements are a pair $\{X, Y\}$, equipped with an inner-product

$$\langle\langle \{X_1, Y_1\}, \{X_2, Y_2\} \rangle\rangle = \langle X_1, X_2 \rangle - \langle Y_1, Y_2 \rangle. \quad (2.26)$$

With this inner-product it is clear that the diagonally embedded G is a subgroup and a maximal isotropic. However the anti-diagonal, whilst being the complementary isotropic, is not a subgroup. The specification of the \mathcal{E} -model is completed by defining

$$\mathcal{E} : \{X, Y\} \mapsto \frac{1 + \lambda^2}{1 - \lambda^2} \{X, -Y\} - \frac{2\lambda}{1 - \lambda^2} \{Y, -X\}, \quad (2.27)$$

from which a flat space generalised metric can be obtained via eq. (2.4). The metric and B-field of the λ -model can be obtained by dressing this generalised metric with flat (algebra) indices with an appropriate frame field as in eq. (2.14). The construction of this frame field is slightly more involved than in the case of the PL model, principally because the anti-diagonal embedding of G is not a subgroup and we are not dealing with a Drinfel'd double. This feature is crucial to ensure that the WZ term in eq. (2.6) plays a role. A second delicate matter is to relate the coset representative $m(x)$, and quantities derived from it, to those obtained in terms of the group element $g(x)$ defining the λ -model. We shall return to both these points in the sequel.

3 Target space description of \mathcal{E} -models

We begin this section by reviewing double field theory on a group manifold, DFT_{wzw} , which will be our framework to implement \mathcal{E} -models. We will then show how the section condition can be solved by introducing a set of frame fields that further describe the generalised geometry of $\mathcal{M} = \mathbb{D}/\tilde{H}$. We will explain how modified supergravity arises out of this procedure.

3.1 A brief review of DFT_{wzw}

We now present a more consolidated target space perspective of the discussion in the previous section. For this we shall employ the framework of DFT_{wzw} [42]; a specification of the $O(D, D)$ symmetric double field theory that assumes an underlying group manifold, \mathbb{D} , of dimension $2D$. The corresponding algebra \mathfrak{d} is as in eq. (2.1), and in particular is equipped with an ad-invariant inner-product, η , of split signature that will be used to raise and lower indices.

We introduce a group element, $\mathbf{g}(X)$, depending on X^I , $I = 1 \dots 2D$, local coordinates on \mathbb{D} and the left-invariant Maurer-Cartan forms $\mathbf{g}^{-1}d\mathbf{g} = \mathbb{E}^A{}_I \mathbb{T}_A dX^I$ from which is constructed $D_A = \mathbb{E}_A{}^I \partial_I$, (with $\mathbb{E}_A{}^I$ the inverse transpose of $\mathbb{E}^A{}_I$) a set of vector fields generating a right action obeying $[D_A, D_B] = \mathbb{F}_{AB}{}^C D_C$.

The NS/NS sector. The common NS sector of DFT_{wzw} consists of a generalised metric \mathcal{H}^{AB} , which *a priori* may depend on all of the X^I , and a generalised dilaton d . The dynamics are encoded by a target space action [42],

$$\begin{aligned}
 S_{\text{NS}} = \int d^{2D} X e^{-2d} & \left(\frac{1}{8} \mathcal{H}^{CD} \nabla_C \mathcal{H}_{AB} \nabla_D \mathcal{H}^{AB} - \frac{1}{2} \mathcal{H}^{AB} \nabla_B \mathcal{H}^{CD} \nabla_D \mathcal{H}_{AC} \right. \\
 & \left. - 2 \nabla_A d \nabla_B \mathcal{H}^{AB} + 4 \mathcal{H}^{AB} \nabla_A d \nabla_B d + \frac{1}{6} F_{ACD} F_B{}^{CD} \mathcal{H}^{AB} \right). \quad (3.1)
 \end{aligned}$$

Here we have introduced a covariant derivative ∇ that acts on a vector density V^A with weight w as,

$$\nabla_A V^B = D_A V^B + \frac{1}{3} F_{AC}{}^B V^C - w F_A V^B. \quad (3.2)$$

The generalised metric has weight $w = 0$ whilst the generalised dilaton e^{-2d} has $w = 1$ and $\nabla_A d = -\frac{1}{2} e^{2d} \nabla_A e^{-2d}$. The density correction makes use of $F_A = D_A \log \det \mathbb{E}$.

object	gen.-diffeomorphisms	2D-diffeomorphisms	global $O(D, D)$
\mathcal{H}_{AB}	tensor	scalar	tensor
$\nabla_A d$	not covariant	scalar	1-form
e^{-2d}	scalar density ($w=1$)	scalar density ($w=1$)	invariant
χ	spinor	scalar density ($w=\frac{1}{2}$)	spinor
η_{AB}	invariant-tensor	invariant-scalar	invariant
$\mathbb{F}_{AB}{}^C$	invariant	invariant	tensor
$E_A{}^I$	invariant	vector	1-form
S_{NS}	invariant	invariant	invariant
$S_{\text{R/R}}$	invariant	invariant	invariant
D_A	not covariant	covariant	covariant
∇_A	not covariant	covariant	covariant

Table 1. Transformation properties of objects under DFT_{WZW} symmetries.

The local symmetries of the action comprise:

1. *Generalised diffeomorphisms* mediated by the generalised Lie derivative

$$\mathcal{L}_\xi V^A = \xi^B \nabla_B V^A - V^B \nabla_B \xi^A + \eta^{AB} \eta_{CD} V^C \nabla_B \xi^D + w \nabla_B \xi^B V^A, \quad (3.3)$$

2. *Conventional 2D-diffeomorphisms* mediated by the Lie derivative

$$L_\xi V^A = \xi^B D_B V^A - w \xi^B F_B V^A + w D_B \xi^B V^A. \quad (3.4)$$

It should be emphasised here that under the conventional 2D-diffeomorphisms objects with curved space indices I, J etc. transform tensorially whereas those with algebra indices A, B , transform as scalars. In particular with respect to this transformation \mathcal{H}_{AB} is a scalar and η_{AB} is an invariant (i.e. constant) scalar. See table 1 for further details.

Closure of the local symmetry algebra necessitates that fields and gauge parameters, and products thereof, can depend on coordinates in only a restricted way. This restriction is called the section condition and reads

$$(D_A f_1 - w_1 F_A f_1)(D^A f_2 - w_2 F^A f_2) = 0, \quad (3.5)$$

in which f_1 and f_2 indicate any field or combinations of fields with the corresponding weights w_1 and w_2 , respectively. Notionally solving this condition should amount to giving a splitting of coordinates $X^I = (\tilde{x}_{\tilde{i}}, x^i)$ in to physical $\{x^i\}$, on which fields can depend, and non-physical $\{\tilde{x}_{\tilde{i}}\}$ on which fields cannot depend. Once a solution to the section condition is adopted of course the full conventional 2D-diffeomorphism symmetry is broken, and all that survives can in fact be absorbed into the generalised diffeomorphisms.

Having the action (3.1), we can derive the corresponding equations of motion by varying it with respect to the generalised metric and the generalised dilaton. Doing so, we find [42]

$$\delta S_{\text{NS}} = \int d^{2D} X e^{-2d} \mathcal{K}_{AB} \delta \mathcal{H}^{AB} \quad \text{and} \quad \delta S_{\text{NS}} = -2 \int d^{2D} X e^{-2d} \mathcal{R} \delta d, \quad (3.6)$$

with

$$\begin{aligned}
 \mathcal{K}_{AB} = & \frac{1}{8} \nabla_A \mathcal{H}_{CD} \nabla_B \mathcal{H}^{CD} - \frac{1}{4} [\nabla_C - 2(\nabla_C d)] \mathcal{H}^{CD} \nabla_D \mathcal{H}_{AB} + 2 \nabla_{(A} \nabla_{B)} d \\
 & - \nabla_{(A} \mathcal{H}^{CD} \nabla_D \mathcal{H}_{B)C} + [\nabla_D - 2(\nabla_D d)] [\mathcal{H}^{CD} \nabla_{(A} \mathcal{H}_{B)C} + \mathcal{H}^C_{(A} \nabla_C \mathcal{H}^D_{B)}] \\
 & + \frac{1}{6} F_{ACD} F_B{}^{CD}, \tag{3.7}
 \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{R} = & 4 \mathcal{H}^{AB} \nabla_A \nabla_B d - \nabla_A \nabla_B \mathcal{H}^{AB} - 4 \mathcal{H}^{AB} \nabla_A d \nabla_B d + 4 \nabla_A d \nabla_B \mathcal{H}^{AB} \\
 & + \frac{1}{8} \mathcal{H}^{CD} \nabla_C \mathcal{H}_{AB} \nabla_D \mathcal{H}^{AB} - \frac{1}{2} \mathcal{H}^{AB} \nabla_B \mathcal{H}^{CD} \nabla_D \mathcal{H}_{AC} + \frac{1}{6} F_{ACD} F_B{}^{CD} \mathcal{H}^{AB}. \tag{3.8}
 \end{aligned}$$

In order to obtain the field equations, one has to take into account that $\delta \mathcal{H}^{AB}$ is not an arbitrary rank two tensor but restricted to symmetric $O(D, D)$ generators. Therefore, one introduces the generalised Ricci tensor [42, 62]

$$\mathcal{R}_{AB} = 2 P_{(A}{}^C \mathcal{K}_{CD} \bar{P}_{B)}{}^D \quad \text{with} \quad P_{AB} = \frac{1}{2} (\eta_{AB} + \mathcal{H}_{AB}) \quad \text{and} \quad \bar{P}_{AB} = \frac{1}{2} (\eta_{AB} - \mathcal{H}_{AB}). \tag{3.9}$$

It projects out the irrelevant components of \mathcal{K} and allows to write the field equations for the NS/NS sector in the compact form

$$\mathcal{R}_{AB} = 0 \quad \text{and} \quad \mathcal{R} = 0. \tag{3.10}$$

The R/R sector. Let us now examine the R/R sector for which the target space action on \mathbb{D} reads [63]

$$S_{\text{R/R}} = \frac{1}{4} \int d^2 D X (\nabla \chi)^\dagger S_{\mathcal{H}} \nabla \chi. \tag{3.11}$$

Here χ is a Majorana-Weyl spinor of $\text{Spin}(D, D)$ and depending on its chirality encodes either type IIA or IIB theories. A natural way to parameterise this spinor is in terms of even or odd differential forms with degree up to D . Let us denote these forms as $C^{(p)}$ so that

$$\chi = \sum_{p=0}^D \frac{1}{2^{p/2} p!} C_{a_1 \dots a_p}^{(p)} \Gamma^{a_1} \dots \Gamma^{a_p} |0\rangle, \tag{3.12}$$

in which the Γ -matrices $\Gamma^A = (\Gamma_a, \Gamma^\alpha)$ obey $\{\Gamma^A, \Gamma^B\} = 2\eta^{AB}$ and $|0\rangle$ is the Clifford vacuum annihilated by the Γ_a . The action of an $O(D, D)$ element O on a spinor, denoted as S_O , is implicitly defined by the Clifford relation

$$\Gamma^A = S_O \Gamma^B S_O^{-1} O_B{}^A. \tag{3.13}$$

The covariant derivative for spinors entering the action is defined as

$$\nabla \chi = \Gamma^A \nabla_A \chi \quad \text{with} \quad \nabla_A \chi = D_A \chi - \frac{1}{12} \mathbb{F}_{ABC} \Gamma^{BC} \chi - \frac{1}{2} F_A \chi, \tag{3.14}$$

where we take into account that χ transforms as a density with weight $w = 1/2$. The Dirac operator $\not{\nabla}$ is nilpotent providing that $\mathbb{F}_{ABC}\mathbb{F}^{ABC} = 0$, this requirement in fact follows from the section condition of DFT and we shall assume it to be the case.

Generalised diffeomorphisms act on the spinor as

$$\mathcal{L}_\xi\chi = \xi^A\nabla_A\chi + \frac{1}{2}\nabla_A\xi_B\Gamma^{AB}\chi + \frac{1}{2}\nabla_A\xi^A\chi, \quad (3.15)$$

and under 2D-diffeomorphisms it transforms exactly as in eq. (3.4) as a scalar with density 1/2. The field strengths are defined as $G = \not{\nabla}\chi$. In order that eq. (3.11) describes the correct degrees of freedom a self-duality condition must be imposed [63]

$$G = -\mathcal{K}G \quad \text{with} \quad \mathcal{K} = C^{-1}S_{\mathcal{H}}, \quad (3.16)$$

in which C is the charge conjugation matrix.⁶

The variation of the action with respect to χ gives rise to the equations of motion

$$\not{\nabla}(\mathcal{K}G) = 0, \quad (3.19)$$

which is automatically satisfied providing the self-duality constraint (3.16) and Bianchi identity are imposed. Furthermore, the NS/NS sector equations of motion (3.10) receive the additional contribution from also varying the R/R part of the action [63]

$$\mathcal{R}_{AB} - \frac{e^{2d}}{16}\mathcal{H}_{(A}{}^C G^\dagger\Gamma_{B)C}\mathcal{K}G = 0. \quad (3.20)$$

3.2 The generalised frame fields

In order to present concrete solutions to the section condition let us restrict our attention to the case relevant to \mathcal{E} -models, i.e. that \mathfrak{d} admits a subalgebra $\tilde{\mathfrak{h}} \subset \mathfrak{d}$. Let $\mathbb{T}_A = (\tilde{T}^a, T_a)$ where \tilde{T}^a are generators of $\tilde{\mathfrak{h}}$ and T_a are the remaining generators whose span we denote \mathfrak{k} . The subalgebra is maximally isotropic with respect to η . The space \mathfrak{k} is automatically maximally isotropic but not necessarily a subalgebra. Locally in a patch, one can always decompose a group element $g \in \mathbb{D}$ as

$$g(X^I) = \tilde{h}(\tilde{x}_{\tilde{i}})m(x^i), \quad \tilde{h} \in \tilde{H} \quad \text{and} \quad m \in \exp(\mathfrak{k}). \quad (3.21)$$

This splitting should be extended globally, working patchwise if a global section m is unavailable [64]. Note that the coset-representative $m(x^i)$ is chosen to be just the exponential of coset generators; this represents a preferred choice of coordinates on \mathbb{D}/\tilde{H} which will be employed in what follows.

⁶Charge conjugation is defined by its action

$$C\Gamma^a C^{-1} = \Gamma_a = (\Gamma^a)^\dagger \quad \text{and} \quad C\Gamma_a C^{-1} = \Gamma^a = (\Gamma_a)^\dagger \quad (3.17)$$

on the Γ -matrices. This constraint requires that $(C^{-1}S_{\mathcal{H}})^2 = 1$ and therefore that $D(D-1)/2$ is odd. Thus, we can only impose it ($D \leq 10$) for [63]

$$D = \{10, 7, 6, 3, 2\}. \quad (3.18)$$

Now we make one further important requirement, namely we demand that

$$\eta^{ij} = 0 \quad \text{of} \quad \eta^{IJ} = \mathbb{E}_A^I \eta^{AB} \mathbb{E}_B^J. \quad (3.22)$$

It was shown in [65] when \mathbb{D}/\tilde{H} is identified with a group manifold (i.e. \mathfrak{d} is a Drinfel'd double) or is a symmetric coset then eq. (3.22) follows directly from eq. (3.21). This will be the case in all examples we are interested in this paper, however in anticipation that there may be more general solutions we keep this as a separate requirement.

Then the section condition in DFT is implemented by demanding physical fields just depend on the coordinates x^i of the target space \mathbb{D}/\tilde{H} and not on \tilde{x}_i . A slight subtlety arises for densities f of weight w where only the combination

$$f |\det \tilde{e}_a^{\tilde{i}}|^{-w} = \hat{f}(x^i) \quad \text{with} \quad \tilde{T}^a \tilde{e}_a^{\tilde{i}} d\tilde{x}_{\tilde{i}} = \tilde{h}^{-1} d\tilde{h}, \quad (3.23)$$

depends on x^i , whilst f depends on all coordinates. The physical fields are then the generalised metric and the corrected dilaton:

$$\hat{\mathcal{H}}^{\hat{I}\hat{J}}(x^i) \quad \text{and} \quad \hat{d}(x^i) = d + \frac{1}{2} \log |\det \tilde{e}_a^{\tilde{i}}|. \quad (3.24)$$

The last equation takes into account that e^{-2d} is the covariant density with weight $w = 1$. Similarly in the R/R sector the coordinate dependence of χ is restricted to

$$S_{\hat{E}}\chi = \hat{\chi} \sqrt{|\det \tilde{e}_a^{\tilde{i}}|} \quad (3.25)$$

where $\hat{\chi}$ depends on the physical coordinates x^i only and $S_{\hat{E}}$ will be the spinorial counterpart of a certain frame-field we shall now define.

We now need to express the actions eq. (3.1) and (3.11) in terms of these restricted quantities $\hat{\mathcal{H}}^{\hat{I}\hat{J}}, \hat{d}$ and $\hat{\chi}$ which can be thought of as living in the generalised tangent space of $\mathcal{M} = \mathbb{D}/\tilde{H}$. To do so we shall show that when the factorisation eq. (3.21) is assumed we can define a set of generalised frame fields $\hat{E}_A^{\hat{I}}$ that obey

- i. $\hat{E}_A^{\hat{I}}$ is an $O(D, D)$ element ,
- ii. $\hat{E}_A^{\hat{I}}$ only depends on the physical coordinates x^i ,
- iii. $\hat{E}_A^{\hat{I}}$ gives rise to the frame algebra,

$$\hat{\mathcal{L}}_{\hat{E}_A} \hat{E}_B^{\hat{I}} = \mathbb{F}_{AB}^C \hat{E}_C^{\hat{I}}, \quad (3.26)$$

where \mathbb{F}_{AB}^C are the structure constants of the double \mathbb{D} and $\hat{\mathcal{L}}$ is the generalised Lie derivative of generalised geometry

$$\hat{\mathcal{L}}_{\xi} V^{\hat{I}} = \xi^{\hat{J}} \partial_{\hat{J}} V^{\hat{I}} + (\partial^{\hat{I}} \xi_{\hat{J}} - \partial_{\hat{J}} \xi^{\hat{I}}) V^{\hat{J}}. \quad (3.27)$$

At this stage we are working locally however where these frame fields can be globally extended they would define a *generalised Leibniz parallelisation* on $\mathcal{M} = \mathbb{D}/\tilde{H}$ [46, 47].

Also note that because of condition **ii** we may use the term generalised Lie derivative and Courant bracket are interchangeable here.

The hatted notation on indices is introduced to emphasise quantities that take values in the generalised tangent space of \mathcal{M} , i.e. when the section condition has been solved so for example

$$V^{\hat{I}} = \begin{pmatrix} v_i \\ v^i \end{pmatrix}, \quad \partial_{\hat{I}} = \begin{pmatrix} 0 & \partial_i \end{pmatrix}, \quad \text{and} \quad \eta_{\hat{I}\hat{J}} = \begin{pmatrix} 0 & \delta_j^i \\ \delta_i^j & 0 \end{pmatrix}. \quad (3.28)$$

In the present context we have the following useful theorem:

Theorem 1. *For each group \mathbb{D} , with a non-degenerate, bilinear, ad-invariant split form η , and a maximally isotropic subgroup \tilde{H} , there exists generalised frame fields on $\mathcal{M} = \mathbb{D}/\tilde{H}$ where eq. (3.21) and eq. (3.22) hold that obey conditions **i–iii** above. These are realized by*

$$\hat{E}_A^{\hat{I}} = M_A^B \hat{V}_B^{\hat{I}} = M_A^B \begin{pmatrix} v^b{}_i & 0 \\ v_b{}^j \rho_{ji} & v_b{}^i \end{pmatrix}_B^{\hat{I}}, \quad (3.29)$$

with

$$M_A^B \mathbb{T}_B = m \mathbb{T}_A m^{-1}, \quad T_a v^a{}_i dx^i + \tilde{T}^a A_{ai} dx^i = \mathbb{T}_A V^A dx^i = dmm^{-1}, \quad v^a{}_i v_a{}^j = \delta_i^j, \quad (3.30)$$

and ρ_{ij} the components of a two-form

$$\rho^{(2)} = \frac{1}{2} \rho_{ij} dx^i \wedge dx^j = \omega^{(2)} - \Omega^{(2)}, \quad (3.31)$$

in which

$$\omega^{(2)} = \frac{1}{2} v^a{}_i A_{aj} dx^i \wedge dx^j, \quad (3.32)$$

and $\Omega^{(2)}$ chosen such that

$$d\Omega^{(2)} = \Omega^{(3)} = \frac{1}{12} \langle\langle dmm^{-1}, [dmm^{-1}, dmm^{-1}] \rangle\rangle = \frac{1}{12} \mathbb{F}_{ABC} V^A \wedge V^B \wedge V^C. \quad (3.33)$$

Proof. Condition **i** is trivially satisfied. The adjoint action of any group element in \mathbb{D} , and in particular M_A^B , is an $O(D, D)$ element as η is adjoint invariant. The second part of (3.29), $\hat{V}_B^{\hat{I}}$, is also $O(D, D)$ valued, indeed it is the product of a b -field transformation and a $\text{GL}(D)$ action. By construction the frame fields only depend on the coordinates $\{x^i\}$ and the condition **ii** is automatic. Finally we have to check the frame algebra condition **iii**. First we make use of the easy identity

$$\partial_{\hat{I}} M_A^B = V^C{}_{\hat{I}} M_A^D \mathbb{F}_{CD}{}^B, \quad (3.34)$$

to show that

$$\hat{E}_{C\hat{I}} \hat{\mathcal{L}}_{\hat{E}_A} \hat{E}_B^{\hat{I}} = M_A^D M_B^E M_C^F (T_{DEF} + S_{DEF}), \quad (3.35)$$

in which

$$T_{ABC} = 3 \hat{V}_{[A}^{\hat{I}} \partial_{\hat{I}} (\hat{V}_B^{\hat{J}}) \hat{V}_{C] \hat{J}}, \quad S_{ABC} = 3 \Lambda_{[A}^G \mathbb{F}_{BC]G}, \quad \Lambda_A^B = \hat{V}_A^i V^B{}_i = \begin{pmatrix} 0 & 0 \\ v_a{}^i A_{bi} & \delta_a^b \end{pmatrix}. \quad (3.36)$$

Since the structure constants are invariant under the adjoint action the proof is completed provided

$$T_{ABC} + S_{ABC} = \mathbb{F}_{ABC}. \quad (3.37)$$

This can be verified component by component:

$$\begin{aligned} S^{abc} &= 0, & T^{abc} &= 0, \\ S^{ab}{}_c &= \tilde{F}^{ab}{}_c, & T^{ab}{}_c &= 0, \\ S^a{}_{bc} &= 2F^a{}_{bc} + 2\tilde{F}^{da}{}_{[b}\Lambda_{c]d}, & T^a{}_{bc} &= -F^a{}_{bc} - 2\tilde{F}^{da}{}_{[b}\Lambda_{c]d}, \\ S_{abc} &= 3F_{abc} + 3F^d{}_{[ab}\Lambda_{c]d}, & T_{abc} &= 3v_a{}^i v_b{}^j v_c{}^k \partial_{[i}\rho_{jk]}. \end{aligned} \quad (3.38)$$

The proof is concluded by substituting the derivative of ρ from eq. (3.33) calculated using

$$d\omega^{(2)} = -\frac{1}{4} (\mathbb{F}_{ABc} V^A \wedge V^B \wedge v^c - \mathbb{F}_{AB}{}^c V^A \wedge V^B \wedge A_c), \quad (3.39)$$

which follows from the Maurer-Cartan identity for V^A . □

Comment 1. Using these frame fields we construct derivative operators $\partial_{\hat{I}} = \hat{E}^A{}_{\hat{I}} D_A$. As detailed in appendix C, $\eta^{\hat{I}\hat{J}} \partial_{\hat{I}} \partial_{\hat{J}} \equiv 0$ and therefore the section condition of DFT is solved.

Comment 2. The twisting of a Courant bracket by an $\mathbf{H} \in H_3(M, \mathbb{R})$ (see e.g. [66]) provides an interpretation for the use of $\Omega^{(2)}$, which may not exist globally (even if the decomposition eq. (3.21) does). Defining new generalised frame fields $\hat{E}'_A{}^{\hat{I}}$ as in eq. (3.29) but with now $\rho_{ij} = \omega_{ij}^{(2)}$ we have that

$$\hat{\mathcal{L}}_{\hat{E}'_A} \hat{E}'_B{}^{\hat{I}} = F_{AB}{}^C \hat{E}'_C{}^{\hat{I}} + \begin{pmatrix} \hat{E}'_A{}^j \hat{E}'_B{}^k \Omega_{jki}^{(3)} \\ 0 \end{pmatrix}, \quad (3.40)$$

such that $\mathbf{H} = \Omega^{(3)}$ appears as such a twisting. If \mathfrak{k} is also a subalgebra (as in the case of a Drinfel'd double) this vanishes.

Comment 3. The assumption of eq. (3.22) allows us to introduce, in addition to η_{AB} and \mathcal{H}_{AB} , the structure

$$\tilde{\omega}_{AB} = \begin{pmatrix} 0 & -1 \\ 1 & 2A_{ai} v^i{}_b \end{pmatrix}. \quad (3.41)$$

It will be shown in a forthcoming paper by one of the authors [67] that $\tilde{\omega}$, dressed with an appropriate adjoint action to transport it around the group manifold \mathbb{D} , and η equips $T\mathbb{D}$ with an (almost) para-hermitian structure. An interesting question, beyond the present scope, is to establish the circumstances in which \mathcal{H}_{AB} will allow a full Born geometry of [68]. Recently these structures were examined for the case of Drinfel'd double [69] and DFT [70].

Let us close this section with some further properties of the frame field that will be employed in the sequel. We consider the quantity

$$\hat{\Omega}_{\hat{I}\hat{J}\hat{K}} = -\partial_{\hat{I}} \hat{E}_{\hat{J}A} \hat{E}_{\hat{K}}{}^A. \quad (3.42)$$

An immediate consequence of the frame algebra is that

$$\widehat{\Omega}_{[IJK]} = \frac{1}{3} \widehat{F}_{IJK} \equiv \frac{1}{3} \widehat{E}^A{}_I \widehat{E}^B{}_J \widehat{E}^C{}_K \mathbb{F}_{ABC}. \quad (3.43)$$

We shall also need the contraction

$$\widehat{\Omega}_{\widehat{K}} = \eta^{IJ} \widehat{\Omega}_{IJK}, \quad (3.44)$$

which can be simplified by making use of eq. (3.34) to show

$$\widehat{V}_A{}^I \widehat{\Omega}_{\widehat{I}} = \Lambda_D{}^E \mathbb{F}_{AE}{}^D - \partial_{\widehat{I}} \widehat{V}_A{}^I. \quad (3.45)$$

Then it follows using the Maurer-Cartan identity for $V^A \mathbb{T}_A = dmm^{-1}$ that

$$\widehat{\Omega}_{\widehat{I}} = \left(-v_a{}^i \tilde{f}^a, \partial_i \log \det v^a{}_j - \Omega_{ij}^{(2)} v_a{}^j \tilde{f}^a \right)_{\widehat{I}}, \quad (3.46)$$

in which

$$\tilde{f}^a = \tilde{F}^{ab}{}_b, \quad f_a = F_{ab}{}^b. \quad (3.47)$$

In the special case of a Drinfel'd double, the final term involving $\Omega^{(2)}$ vanishes. Here we see that $\widehat{\Omega}_{\widehat{I}}$ will play an important role in the case that $\tilde{\mathfrak{h}}$ is *non-unimodular*, i.e when $\tilde{F}^{ab}{}_b \neq 0$. The quantities $\widehat{F}_{\widehat{IJK}}$ can be thought of as generalised fluxes and, whilst not essential for what follows, this view is explored in the appendix D.

3.3 Equivalence to (modified) type II supergravity

In what follows, we apply the idea of [44] and rewrite the action (3.1) using the generalised frame field. We parametrise the generalised metric as in eq. (2.14) and write

$$\widehat{\mathcal{H}}^{\widehat{I}\widehat{J}} = \widehat{E}_A{}^{\widehat{I}} \mathcal{H}^{AB} \widehat{E}_A{}^{\widehat{J}}. \quad (3.48)$$

Similarly, we write the generalised dilaton, recalling eq. (3.24) as

$$\widehat{d} = \phi - \frac{1}{4} \log |\det g_{ij}| = d + \frac{1}{2} \log |\det \tilde{e}^a{}_{\tilde{k}}|. \quad (3.49)$$

Taking into account that,

$$\partial_{\widehat{I}} \widehat{\varphi}(x^i) = \widehat{E}^A{}_{\widehat{I}} D_A \widehat{\varphi}(x^i), \quad (3.50)$$

we can make use of the property that $\widehat{E}_A{}^{\widehat{I}}$ satisfies the frame algebra (3.26) to pull covariant derivatives to generalised tangent bundle. An illustrative example is ∇_A acting on a weightless vector V^B for which

$$\nabla_A V^B \rightarrow \partial_{\widehat{I}} \widehat{V}^{\widehat{J}} + (\widehat{\Omega}_{[\widehat{I}\widehat{K}\widehat{L}]} - \widehat{\Omega}_{\widehat{I}\widehat{K}\widehat{L}}) \eta^{\widehat{L}\widehat{J}} \widehat{V}^{\widehat{K}} \quad (3.51)$$

with $\widehat{\Omega}_{\widehat{IJK}}$ defined in eq. (3.42). The generalization to higher rank tensors follows immediately.

Next, we take a look at the covariant derivative for the generalised dilaton for which we must pay attention to the weight factor,

$$\nabla_A d = D_A d + \frac{1}{2} D_A \log |\det \mathbb{E}^B{}_I| = D_A \widehat{d} + \frac{1}{2} D_A \log |\det v^a{}_i|. \quad (3.52)$$

Making use of eq. (3.46) we have

$$\nabla_A d \rightarrow \partial_{\hat{I}} \widehat{d} + \mathbf{X}_{\hat{I}} + \frac{1}{2} \widehat{\Omega}_{\hat{I}}, \quad \mathbf{X}_{\hat{I}} = \frac{1}{2} \begin{pmatrix} \tilde{f}^a v_a{}^i \\ \tilde{f}^a v_a{}^j \Omega_{ij}^{(2)} \end{pmatrix}. \quad (3.53)$$

When $\tilde{\mathfrak{h}}$ is non-unimodular we have by definition $\mathbf{X}_{\hat{I}} \neq 0$ [51]. In the following, we will first consider the NS sector, treating unimodular and non-unimodular cases in turn, and then we discuss the R/R sector pertaining to both cases.

3.3.1 Unimodular case

Pulling all quantities in the action (3.1) to the generalised tangent space, and doing some algebra, we obtain for $\mathbf{X}_{\hat{I}} = 0$ the action of DFT with the section condition solved

$$\begin{aligned} S_{\text{NS}} = V_{\tilde{H}} \int d^D x e^{-2\widehat{d}} & \left(\frac{1}{8} \widehat{\mathcal{H}}^{\hat{K}\hat{L}} \partial_{\hat{K}} \widehat{\mathcal{H}}_{\hat{I}\hat{J}} \partial_{\hat{L}} \widehat{\mathcal{H}}^{\hat{I}\hat{J}} \right. \\ & \left. - 2 \partial_{\hat{I}} \widehat{d} \partial_{\hat{J}} \widehat{\mathcal{H}}^{\hat{I}\hat{J}} - \frac{1}{2} \widehat{\mathcal{H}}^{\hat{I}\hat{J}} \partial_{\hat{J}} \widehat{\mathcal{H}}^{\hat{K}\hat{L}} \partial_{\hat{L}} \widehat{\mathcal{H}}_{\hat{I}\hat{K}} + 4 \widehat{\mathcal{H}}^{\hat{I}\hat{J}} \partial_{\hat{I}} \widehat{d} \partial_{\hat{J}} \widehat{d} \right). \end{aligned} \quad (3.54)$$

All occurrences of $\widehat{\Omega}_{\hat{I}\hat{K}\hat{L}}$ either directly cancel or occur in contractions that vanish due to working in a particular solution of the section condition. Let us emphasise that in eq. (3.54) the section condition has been implemented, the fields only depend on the coordinates x , the integral is only over these coordinates, and the integration over \tilde{x} has been performed with a volume $V_{\tilde{H}}$ arising from the dilaton factor in the measure. It is by now well established [63, 71] that the equations of motion derived from this theory can be equated to the common NS sector (super)gravity field equations for g_{ij}, B_{ij}, ϕ (see appendix A for the supergravity field equations used).

3.3.2 Non-unimodular case

If \tilde{H} is not unimodular, we instead obtain generalised type II [48]. This is a modification at the level of the equations of motion, described in detail in appendix A, that depends crucially on a Killing vector I obeying

$$L_I g = 0, \quad L_I H = 0, \quad (3.55)$$

where L_I is the conventional Lie derivative along I , and a one form Z further constrained to obey

$$dZ + \iota_I H = 0, \quad \iota_I Z = 0. \quad (3.56)$$

The conditions eq. (3.56) allows the construction of a differential which acts on the formal sum of forms

$$\mathbf{d} = d + H \wedge - Z \wedge - \iota_I, \quad \mathbf{d}^2 = -L_I, \quad (3.57)$$

such that when the differential form is invariant under I the differential is nilpotent. This differential operator will be important when discussing the R/R sector.

The first of eq. (3.56) may be integrated to yield

$$Z = d\phi + \iota_I B - V, \tag{3.58}$$

in which $H = dB$ locally and $L_I B = dV$. In the absence of the modifications due to the Killing vector I the scalar field ϕ coincides with the conventional dilaton.⁷ In the language of mDFT [51, 72], the corresponding modifications to the DFT equations of motion are implemented by a shift in the derivative of the DFT dilaton

$$\partial_{\hat{I}} \hat{d} \rightarrow \partial_{\hat{I}} \hat{d} + \mathbf{X}_{\hat{I}}. \tag{3.60}$$

The DFT shift vector in (3.60) is related to the modified supergravity vectors by,

$$\mathbf{X}_{\hat{I}} = \begin{pmatrix} I^i \\ -V_i \end{pmatrix}. \tag{3.61}$$

The DFT vector $\mathbf{X}^{\hat{I}}$ is not an arbitrary, instead reflecting the requirements eq. (3.55)–(3.59) it is constrained to be a generalised Killing vector⁸ obeying,

$$\hat{\eta}^{\hat{I}\hat{J}} \mathbf{X}_{\hat{I}} \mathbf{X}_{\hat{J}} = 0, \quad \hat{\mathcal{L}}_{\mathbf{X}} \hat{\mathcal{H}}^{\hat{I}\hat{J}} = 0 \quad \text{and} \quad \hat{\mathcal{L}}_{\mathbf{X}} \hat{d} = 0. \tag{3.62}$$

Since we know already in the unimodular case that the DFT equations of motion are recovered, it follows that in the non-unimodular case the mDFT equations are recovered with the identification of the DFT vector $\mathbf{X}^{\hat{I}}$ with that in eq. (3.53) i.e. with

$$I = \frac{1}{2} \tilde{f}^a v_a^i \partial_i, \quad V = \iota_I \Omega^{(2)}. \tag{3.63}$$

Whilst V here depends on a choice of $\Omega^{(2)}$ it was shown in [51] that in fact there is a gauge freedom that allows one to take V to be zero.

It is immediate that the first of eq. (3.62) holds but we now investigate under what circumstances the remaining constraints of eq. (3.62) are valid. Here we make use of the generalised frame fields and transport the results back to the flat indices with $\xi^A = \mathbf{X}^{\hat{I}} \hat{E}_{\hat{I}}^A$ and $\mathcal{H}^{AB} = \hat{E}_{\hat{I}}^A \hat{\mathcal{H}}^{\hat{I}\hat{J}} \hat{E}_{\hat{J}}^B$. A short calculation shows that,

$$\hat{E}_{\hat{I}}^A \hat{E}_{\hat{J}}^B \hat{\mathcal{L}}_{\mathbf{X}} \hat{\mathcal{H}}^{\hat{I}\hat{J}} = \xi^C D_C \mathcal{H}^{AB}, \tag{3.64}$$

⁷To make contact with the notation of [51] we define $U = \iota_I B - V$ such that $Z = d\phi + U$. The split of $d\phi$ and U is somewhat arbitrary since one could shift $\phi \rightarrow \phi + \alpha$ and $U \rightarrow U - d\alpha$ and so can be fixed by demanding

$$L_I \phi = 0, \quad \iota_I U = 0. \tag{3.59}$$

⁸To see this recall that a DFT gauge transformation generated by a vector $\mathbf{V}_{\hat{I}} = (v^i, \tilde{v}_i)$ acts as $\delta \mathcal{H}_{\hat{I}\hat{J}} = \hat{\mathcal{L}}_{\mathbf{V}} \mathcal{H}_{\hat{I}\hat{J}}$ and in the solution to the section condition $\partial_{\hat{I}} = (0, \partial_i)$ simply generates diffeomorphism $\delta g = L_v g$ and gauge transformations $\delta B = L_v B + d\tilde{v}$.

and hence if \mathcal{H}^{AB} is constant (as is in case of \mathcal{E} models) the second of eq. (3.62) holds. For the third of eq. (3.62) we have

$$\begin{aligned} \widehat{\mathcal{L}}_{\mathbf{X}}\widehat{d} &= \mathbf{X}^{\hat{I}}\partial_{\hat{I}}\widehat{d} - \frac{1}{2}\partial_{\hat{I}}\mathbf{X}^{\hat{I}} \\ &= \mathbf{X}^{\hat{I}}\left(\partial_{\hat{I}}\widehat{d} + \frac{1}{2}\partial_{\hat{I}}\log\det v_i^a\right) - \frac{1}{4}\tilde{f}^a f_a + \frac{1}{4}\tilde{f}^e F_e{}^{pq}\Lambda_{pq}. \end{aligned}$$

Now taking the trace of the Jacobi identity for the subalgebra $\tilde{\mathfrak{h}}$ yields $\tilde{f}^a \tilde{F}^{bc}{}_a \equiv 0$ such that the final term in the above vanishes. A consequence of the Jacobi identities for \mathfrak{d} is that $\mathbb{F}_{ABC}\mathbb{F}^{ABC} = 4\tilde{f}^a f_a$. Since we require $\mathbb{F}_{ABC}\mathbb{F}^{ABC} = 0$ to avoid violations of the section condition for the cases we are interested in $\tilde{f}^a f_a = 0$.

We can then make use of eq. (3.24) to recast the result in terms of conventional volume preserving 2D-diffeomorphism acting on d :

$$\widehat{\mathcal{L}}_{\mathbf{X}}\widehat{d} = L_{\xi}d \equiv -\frac{1}{2}e^{2d}L_{\xi}e^{-2d}. \tag{3.65}$$

Hence when the DFT_{WZW} dilaton, d , is invariant under 2D-diffeomorphisms then indeed we obey the criteria in eq. (3.62) and it is evident that we reproduce the field equations of modified SUGRA.

It is interesting to ask what happens at the level of the action since it is thought that generalised SUGRA does not admit an action [52]. So what goes wrong when we try to derive an action analogous to (3.54) by translating to the generalised tangent space? To solve this puzzle, remember that we need integration by parts to make sense of the action (3.54) and i.e. derive the field equations. This operation requires that the identity

$$\int dX^{2D}\partial_I(|\det \mathbb{E}|\mathbb{E}_A{}^I\phi) = \int dX^{2D}|\det \mathbb{E}|D_A\phi \tag{3.66}$$

holds. A quick calculation shows that this relation requires $\mathbb{F}_{AB}{}^B = 0$, which is always the case because the full double is always unimodular. However in (3.54), we also integrate out the non-physical directions $\{\tilde{x}_{\tilde{i}}\}$ to obtain an action just on the physical target space \mathbb{D}/\tilde{H} . For the unimodular case this is perfectly fine because integration by parts works on \mathbb{D}/\tilde{H} as well as on \tilde{H} independently. But in the non-unimodular case, $\tilde{F}^{ab}{}_b = \tilde{f}^a \neq 0$ obstructs integration by parts on \tilde{H} . Therefore, we are not allowed to integrate out \tilde{H} and write an action just on \mathbb{D}/\tilde{H} . Instead we require a genuinely doubled action. That also explains the problems in conventional DFT/EFT to find an action. There, the integration is only performed over the physical space after solving the section condition, while in DFT_{WZW} it is always over the full space.

3.3.3 R/R sector

As for the NS/NS sector, we now want to show that this description is equivalent to the R/R sector of type IIA/B supergravity, or modified type II SUGRA if \tilde{H} is not unimodular. So, we pull all quantities to the generalised tangent space. We start with the covariant derivative

$$|\det \tilde{e}^a{}_i|^{-1/2}S_{\widehat{E}}\nabla\chi = \left(\not{\partial} - (\not{\partial}S_{\widehat{E}})S_{\widehat{E}}^{-1} - \frac{1}{12}\widehat{F}_{\hat{I}\hat{J}\hat{K}}\widehat{\Gamma}^{\hat{I}\hat{J}\hat{K}} - \frac{1}{2}\not{\partial}\log|\det v^a{}_i|\right)\widehat{\chi} \tag{3.67}$$

which arises after substituting $\widehat{\chi} = |\det \tilde{e}^a_i|^{-1/2} S_{\widehat{E}} \chi$ and identifying

$$\widehat{\Gamma}^I = S_{\widehat{E}} \Gamma^A S_{\widehat{E}}^{-1} \widehat{E}_A{}^I, \quad \not\partial = \widehat{\Gamma}^I \partial_I. \quad (3.68)$$

We can simplify further as

$$\begin{aligned} -(\not\partial S_{\widehat{E}}) S_{\widehat{E}}^{-1} &= \frac{1}{4} \widehat{\Omega}_{IJK} \widehat{\Gamma}^I \widehat{\Gamma}^{JK} = \frac{1}{12} \widehat{F}_{IJK} \widehat{\Gamma}^{IJK} + \frac{1}{2} \widehat{\Omega}^J{}_{JI} \widehat{\Gamma}^I \\ &= \frac{1}{12} \widehat{F}_{IJK} \widehat{\Gamma}^{IJK} + \frac{1}{2} \not\partial \log |\det v^a_i| - \mathbf{X}_I \widehat{\Gamma}^I. \end{aligned} \quad (3.69)$$

Thus we have

$$\widehat{G} = |\det \tilde{e}^a_i|^{-1/2} S_{\widehat{E}} G = |\det \tilde{e}^a_i|^{-1/2} S_{\widehat{E}} \not\partial \chi = (\not\partial - \mathbf{X}_I \widehat{\Gamma}^I) \widehat{\chi}. \quad (3.70)$$

We are now able to consider the self-duality constraint eq. (3.16) pulled to the generalised tangent space which gives

$$\widehat{G} = -C^{-1} S_{\widehat{\mathcal{H}}} \widehat{G}. \quad (3.71)$$

To cast the results in the simplest form we follow [63] and make use of the decomposition of the spinor representative of the generalised metric⁹

$$S_{\widehat{\mathcal{H}}} = S_B^{-1} S_{g^{-1}} S_B \quad \text{with} \quad S_B = \exp(-B_{ij} \widehat{\Gamma}^{ij}). \quad (3.72)$$

Defining

$$\widehat{\mathcal{F}} = e^\phi S_B (\not\partial - \mathbf{X}_I \widehat{\Gamma}^I) \widehat{\chi}, \quad (3.73)$$

then, from [63], the self-duality condition reads,

$$\widehat{\mathcal{F}} = -S_g C^{-1} \widehat{\mathcal{F}}. \quad (3.74)$$

We also define a different set of potential $\widehat{\alpha} = e^\phi S_B \widehat{\chi}$ such that

$$\widehat{\mathcal{F}} = e^\phi S_B (\not\partial - \mathbf{X}_I \widehat{\Gamma}^I) e^{-\phi} S_B^{-1} \widehat{\alpha} = \mathbf{d}\widehat{\alpha} \quad (3.75)$$

Here note the appearance of the exterior derivative introduced in eq. (3.57). This is exactly as the R/R sector enters in mDFT in [48, 51]. Combining the Bianchi identity $\mathbf{d}\widehat{\mathcal{F}} = 0$ and eq. (3.57) shows that the Lie derivative $L_I \widehat{\mathcal{F}} = 0$ without imposing any further constraints on the R/R fields.

4 The \mathcal{E} -model conditions

Here we define how the condition for Poisson-Lie symmetry or more generally the structure behind an \mathcal{E} -model can be simply stated in the context of DFT_{WZW} . Namely we propose:

The conditions of an \mathcal{E} -model are that the fields \mathcal{H}^{AB} , d and G of DFT_{WZW} are invariant under volume preserving 2D-diffeomorphisms.

In this section we shall follow through this proposal to constrain the structure of the dilaton and R/R sector.

⁹This follows from writing $\widehat{\mathcal{H}} = \begin{pmatrix} 1 & 0 \\ B & 1 \end{pmatrix} \begin{pmatrix} g^{-1} & 0 \\ 0 & g \end{pmatrix} \begin{pmatrix} 1 & -B \\ 0 & 1 \end{pmatrix}$.

4.1 NS sector

In this sector the condition simply implies that the \mathcal{H}^{AB} is a constant, exactly matching the set up in section 2. Applying this restriction, and similar on the dilaton that we turn to momentarily, the equations of motion simplify significantly. The Ricci scalar reduces to the scalar potential of gauged supergravity

$$\mathcal{R} = \frac{1}{12} \mathbb{F}_{ACE} \mathbb{F}_{BDF} (3\mathcal{H}^{AB} \eta^{CD} \eta^{EF} - \mathcal{H}^{AB} \mathcal{H}^{CD} \mathcal{H}^{EF}), \quad (4.1)$$

without section condition violating contributions $1/6 \mathbb{F}_{ABC} \mathbb{F}^{ABC}$. For the generalised curvature \mathcal{R}_{AB} , we find

$$\mathcal{R}_{AB} = \frac{1}{8} (\mathcal{H}_{AC} \mathcal{H}_{BF} - \eta_{AC} \eta_{BF}) (\mathcal{H}^{KD} \mathcal{H}^{HE} - \eta^{KD} \eta^{HE}) \mathbb{F}_{KH}^C \mathbb{F}_{DE}^F. \quad (4.2)$$

This results matches perfectly with the RG flow calculation for a double sigma model presented in [32, 33] in which

$$\frac{\partial \mathcal{H}_{AB}}{\partial \log \mu} = \mathcal{R}_{AB}. \quad (4.3)$$

4.2 Dilaton

For the dilaton we have to take into account that the covariant quantity e^{-2d} has weight $w = 1$. Hence, we demand

$$L_\xi e^{-2d} \equiv \xi^A D_A e^{-2d} - \xi^A F_A e^{-2d} + D_A \xi^A e^{-2d} = 0, \quad (4.4)$$

where we recall $F_A = D_A \log |\det \mathbb{E}^B_I|$. The last term vanishes, because the $2D$ -diffeomorphisms which we are considering are area preserving. This leaves us with

$$\xi^I \partial_I (2d + \log |\det v^a_i| + \log |\det \tilde{v}_a^{\tilde{i}}|) = 0. \quad (4.5)$$

Plugging in the expression for the generalised dilaton

$$d = \phi - \frac{1}{4} \log |\det g_{ij}| - \frac{1}{2} \log |\det \tilde{v}_a^{\tilde{i}}|, \quad (4.6)$$

we obtain the condition

$$\phi - \frac{1}{4} \log |\det g_{ij}| + \frac{1}{2} \log |\det v^a_i| = \phi_0, \quad (4.7)$$

with ϕ_0 a constant.

For the case of unimodular PL models it can be seen in a few lines that this condition is fulfilled by the dilaton introduced using heavy duty mathematical treatment in [73]. The details of this equivalence are provided as appendix material in section E. Similarly for λ -models this prescription provides the known dilaton, also detailed in E.

4.3 R/R sector

We demand that the field strength $G = \nabla\chi$ is invariant under arbitrary $2d$ diffeomorphisms i.e.

$$L_\xi G \equiv \xi^A D_A G - \frac{1}{2}(\xi^A F_A - D_A \xi^A)G = 0. \quad (4.8)$$

In general for a scalar density, G , of 2D-diffeomorphisms of weight w (and here $w = \frac{1}{2}$) we could define

$$G = |\det \mathbb{E}|^w G_0 \quad (4.9)$$

and the invariance condition is satisfied for G_0 being constant. Here we have a further consequence since we can make use of the definition of the covariant derivative to show this requires that

$$\nabla_A G = -\frac{1}{12} \mathbb{F}_{ABC} \Gamma^{BC} G, \quad (4.10)$$

and as a consequence, assuming the Bianchi identity $0 = \nabla\chi G$, upon contraction with a gamma matrix we have a necessary condition

$$\mathbb{F}_{ABC} \Gamma^{ABC} G = 0. \quad (4.11)$$

Notice that the operator \hat{F} is nilpotent by virtue of the standard properties of Γ -matrices and the Jacobi identity of $\mathbb{F}_{AB}{}^C$. Taking into account the dilaton and the R/R spinor weights we have that the equation of motion involves purely constant algebraic quantities

$$\mathcal{R}_{AB} - \frac{e^{2\phi_0}}{16} \mathcal{H}_{(A}{}^C G_0^\dagger \Gamma_{B)C} \mathcal{K} G_0 = 0. \quad (4.12)$$

When transported to generalised tangent space via $\hat{G} = |\det \tilde{v}_a{}^i|^{-1/2} S_{\hat{E}} G$ we simply have

$$\hat{F} \hat{G} = \frac{1}{12} \hat{F}_{\hat{I}\hat{J}\hat{K}} \hat{\Gamma}^{\hat{I}\hat{J}\hat{K}} \hat{G} = 0. \quad (4.13)$$

Notice further that $S_{\hat{E}}$ contains three factors, the first is the spinor counterpart S_M of the adjoint action $M_A{}^B$, the second is a B-field S_ρ shift induced by the two form ρ and the third is the spinor counterpart $S_{\hat{V}}$ of the $GL(D)$ transformation induced by the vector fields $v_a{}^i$. Now this last transformation $S_{\hat{V}}$ carries with it a multiplicative factor of $|\det v|^{-\frac{1}{2}}$. This factor combines with the $|\det \tilde{v}|^{-1/2}$ to cancel the same factors coming from the weighting and pragmatically speaking in the end to pass to the target space it will be sufficient to calculate $S_\rho S_M G_0$. Where the context is clear we shall not overcrowd and already burdensome notation with the subscript G_0 and understand the push to the generalised tangent space in the above sense.

4.4 Fourier-Mukai transformation

An alternative approach to study the transformation of R/R fluxes is a Fourier-Mukai transformation. This idea was already applied to Abelian [74] and non-Abelian T-duality [75]. Here we show that our previous results allow us to write the R/R flux transformations also in terms of a Fourier-Mukai transformation for the full Poisson-Lie T-duality. Especially,

we give an explicit construction for the gauge invariant flux ω of the topological defect mediating the transformation.

Let us first set up our notation. We have two (pseudo)-Riemannian target spaces \mathcal{M} and $\widetilde{\mathcal{M}}$ which are connected by Poisson-Lie T-duality. We are not restricted to the cases where there are two maximally isotropic subgroups in a single decomposition of \mathfrak{d} , one could imagine taking an algebra \mathfrak{d} and by performing global $O(D, D)$ rotation making two different Manin quasi-triple decompositions. Both target spaces are D -dimensional and we denote their coordinates as x^i and $\tilde{x}^{\tilde{i}}$, respectively. Furthermore, their metrics, g_{ij} and $\tilde{g}_{\tilde{i}\tilde{j}}$, are used to define a Hodge star on both of them.¹⁰ We are interested in their R/R flux $F_{(p)}$ and $\widetilde{F}_{(p)}$. They are governed by the self-duality conditions

$$F_{(p)} = (-1)^{\frac{(D-p)(D-p-1)}{2}} \star F_{(D-p)} \tag{4.15}$$

and the same for $\widetilde{F}_{(p)}$. These fluxes can be related by the Fourier-Mukai transformation

$$\widetilde{\mathcal{F}} = \frac{s}{V(\mathcal{M})} \int_{\mathcal{M}} \mathcal{F} \wedge e^\omega, \tag{4.16}$$

where s is the signature of the metric on \mathcal{M} and $V(\mathcal{M})$ denotes its volume. It arises after integration over the volume form $v(\mathcal{M}) = \sqrt{|\det g|} dx^1 \wedge \dots \wedge dx^D$. The remaining, essential ingredient in the equation is the two-form ω . In order to fix this form, we need to remember how Poisson-Lie T-duality works for the R/R fluxes in DFT. \widehat{F} and $\widetilde{\widehat{F}}$ are represented by Majorana-Weyl spinor of $O(D, D)$. Using the generalised frames field $\widehat{E}_A^{\widehat{I}}$ and $\widetilde{\widehat{E}}_A^{\widetilde{I}}$ of the two backgrounds we can write down the $O(D, D)$ transformation [44]

$$\widetilde{\widehat{O}}^{\widetilde{I}}_{\widehat{j}} = \widetilde{\widehat{E}}_A^{\widetilde{I}} \widetilde{\widehat{E}}^A_{\widehat{j}} \tag{4.17}$$

relating these two spinors. It acts as

$$\widetilde{\widehat{F}} = \sqrt{|\det \tilde{e}^{a_{\tilde{i}} e_a^j}|} S_{\widetilde{B}} S_{\widetilde{O}} S_{-B} \widehat{F}. \tag{4.18}$$

A canonical way to parameterise the $O(D, D)$ element \widehat{O} is

$$\widehat{O}^{\widehat{I}}_{\widehat{j}} = \begin{pmatrix} r_{\widehat{i}}^{\widehat{j}} + b_{\widehat{i}\widehat{k}}^{\widehat{l}} r_{\widehat{l}}^{\widehat{k}} \beta^{\widehat{l}\widehat{j}} & b_{\widehat{i}\widehat{k}}^{\widehat{l}} r_{\widehat{l}}^{\widehat{k}} \\ r_{\widehat{k}}^{\widehat{i}} \beta^{\widehat{k}\widehat{j}} & r_{\widehat{j}}^{\widehat{i}} \end{pmatrix}. \tag{4.19}$$

It allows us to directly identify ω with

$$\begin{aligned} \omega &= -\frac{1}{2} \text{Tr} \log(r_a^b) - B - \frac{1}{2} \beta_{ij} dx^i \wedge dx^j + \frac{1}{2} b_{\tilde{i}\tilde{j}} d\tilde{x}^{\tilde{i}} \wedge d\tilde{x}^{\tilde{j}} + \widetilde{B} - r_{\tilde{i}\tilde{j}} d\tilde{x}^{\tilde{i}} \wedge dx^j \\ &= \omega_{(0,0)} + \omega_{(2,0)} + \omega_{(0,2)} + \omega_{(1,1)}. \end{aligned} \tag{4.20}$$

¹⁰We use the explicit expression

$$(\star A)_{i_1 \dots i_p} = \frac{1}{\sqrt{|\det g|(D-p)!}} g_{i_1 j_1} \dots g_{i_p j_p} \epsilon^{k_p+1 \dots k_D j_1 \dots j_p} A_{k_p+1 \dots k_D} \tag{4.14}$$

for the Hodge star with $\epsilon^{12 \dots D} = 1$. In this section we chose to restore “upstairs” positions for the indices $\tilde{x}^{\tilde{i}}$ and “downstairs” for dual algebra generators \widetilde{T}_a — this is so as not to interfere with the standard notation for differential forms.

In the first line, we lowered the two indices of β^{ij} with the metric g_{ij} and the same for the second index of $r_{\tilde{i}}^j$. Additionally, we denote a p -form on \mathcal{M} and a q -form on $\widetilde{\mathcal{M}}$ as $\omega_{(p,q)}$. As we will see later, the contribution $\omega_{(0,0)}$, which depends on $r_a^b = \tilde{e}_a^{\tilde{i}} r_{\tilde{i}}^j e^b_j$, vanishes if we have a Drinfeld double and is only relevant for Manin quasi-triple. Finally, B and \tilde{B} are the B -fields on the target space and its dual.

To show that the expression presented in eq. (4.20) is indeed the correct form of ω , we calculate

$$\tilde{\mathcal{F}} = \frac{s}{V(\mathcal{M})} \int_{\mathcal{M}} (F_{(D)} \wedge (1 + \omega_{(0,0)} + \omega_{(0,2)}) + F_{(D-1)} \wedge \omega_{(1,1)} + F_{(D-2)} \wedge \omega_{(2,0)} + \dots) \tag{4.21}$$

up to the linear order in ω and compare it with the DFT result. We have to take into account the two properties

$$\star 1 = s v(\mathcal{M}) \quad \text{and} \quad \star F_{(p)} \wedge \varphi_{(p)} = \frac{s(-1)^{(D-p)p}}{p!} F_{i_1 \dots i_p} \varphi^{i_1 \dots i_p} v(\mathcal{M}), \tag{4.22}$$

of the Hodge star. They allow to simplify each term appearing in the expansion (4.21) to

$$F_{(D-p)} \wedge \omega_{(p,q)} = s(-1)^{p(D-1)} \frac{1}{p!q!} F_{i_1 \dots i_p} \omega^{i_1 \dots i_p}_{\tilde{j}_1 \dots \tilde{j}_q} v(\mathcal{M}) \wedge d\tilde{x}^{\tilde{j}_1} \wedge \dots \wedge d\tilde{x}^{\tilde{j}_q}. \tag{4.23}$$

Applying this relation, we find

$$\begin{aligned} \tilde{\mathcal{F}} = & \left(F \left(1 - \frac{1}{2} \text{Tr} \log(r_a^b) \right) + \frac{1}{2} F(b_{\tilde{i}\tilde{j}} + B_{\tilde{i}\tilde{j}}) d\tilde{x}^{\tilde{i}} \wedge d\tilde{x}^{\tilde{j}} \right. \\ & \left. + (-1)^D F_i r_{\tilde{j}}^i d\tilde{x}^{\tilde{j}} - \frac{1}{2} F_{ij} (B^{ij} + \beta^{ij}) + \dots \right). \end{aligned} \tag{4.24}$$

Note that this relation crucially relies on the assumption that the R/R fluxes admit Poisson-Lie symmetry. Otherwise we would not be able to perform the integration and cancel the volume factor in front of the integral. One can check that performing the spinor transformation (4.18) gives exactly the same result. Thus the ansatz (4.20) is correct. Finally, one has to take into account that in IIA/IIB there are either just even or odd contributions for F_p . Thus, eq. (4.24) simplifies accordingly. For D even, the Fourier-Mukai transformations maps the R/R sector of IIB to IIB and of IIA to IIA, while for D odd both are exchanged. Taking into account not just the leading order, but also all higher order terms (the are hidden in \dots in (4.24)), one can extract all R/R fields of the dual background. However, from the computational effort this is in general more challenging as performing the spinor transformation giving in (4.18) directly.

If we specialise to Poisson-Lie T-duality on a Drinfel'd double, using the generalised frame field (3.29), we find

$$b_{\tilde{i}\tilde{j}} = 0, \quad \beta^{ij} = e_c^i (\Pi^{cd} - \tilde{\Pi}^{cd}) e_a^j \quad \text{and} \quad r_{\tilde{i}}^j = \tilde{e}^a_{\tilde{i}} e_a^j. \tag{4.25}$$

The last equation implies that $r_a^b = \delta_a^b$ and thus $\omega_{(0,0)}=0$. Writing furthermore the metric on \mathcal{M} as $g_{ij} = e^a_i g_{ab} e^b_j$, we obtain

$$\omega = \tilde{B} - B - \frac{1}{2} g_{ac} (\Pi^{cd} - \tilde{\Pi}^{cd}) g_{db} e^a \wedge e^b - g_{ab} \tilde{e}^a \wedge e^b. \tag{4.26}$$

For the case of non-Abelian T-duality, $\Pi^{ab} = 0$, $\tilde{\Pi}^{ab} = -f^{ab}{}_c x^c$ and g_{ab} is constant. Then the equation for ω above simplifies to

$$\omega = \tilde{B} - B - \frac{1}{2}g_{ac}g_{bd}f^{cd}{}_e x^e e^a \wedge e^b - g_{ab}dx^a \wedge e^b \tag{4.27}$$

and matches the result in [75].

5 Application to integrable deformations

In the following we will give examples of how the formalism described in this paper can be applied to \mathcal{E} -models. In particular we will show how to recover the R/R-sector and dilaton completing the SUGRA embedding for the η - and the λ -models.

In the first subsection we study these theories at the level of the DFT_{WZW} defined on \mathbb{D} and then show that we recover the conventional target space descriptions on $\mathcal{M} = \mathbb{D}/\tilde{H}$. Whilst the solutions presented here are not new to the literature they serve to demonstrate all the features we have described thus far.

5.1 Deformations based on the (m)CYBE

Each solution, \mathcal{R} , of the mCYBE on \mathfrak{g} gives rise to a canonical group manifold $\mathbb{D} = \mathfrak{g} \oplus \mathfrak{g}_{\mathcal{R}}$ as described in the appendix B. The structure constants of \mathbb{D} are related to those of \mathfrak{g} (denoted by $f_{ab}{}^c$) according to

$$F_{abc} = 0, \quad F_{ab}{}^c = f_{ab}{}^c, \quad \tilde{F}^{ab}{}_c = R^{ad}f_{cd}{}^b - R^{bd}f_{cd}{}^a = \tilde{f}^{ab}{}_c, \quad \tilde{F}^{abc} = 0. \tag{5.1}$$

For the YB-deformations described by the action (2.21), the generalised metric reads

$$\mathcal{H}^{AB} = \begin{pmatrix} \eta\kappa_{ab} & -\eta\kappa_{ac}R^{cb} \\ \eta R^{ac}\kappa_{cb} & \frac{\kappa^{ab}}{\eta} - \eta R^{ac}\kappa_{cd}R^{db} \end{pmatrix}, \tag{5.2}$$

in which κ is the Cartan-Killing form on \mathfrak{g} . Here η can be considered a deformation parameter. One can simplify the form of \mathcal{H}^{AB} considerably by performing the $O(D,D)$ transformation

$$\mathcal{O}_{\check{A}}{}^B = \begin{pmatrix} \sqrt{\eta}\delta^a{}_b & -\sqrt{\eta}R^{ab} \\ 0 & \frac{1}{\sqrt{\eta}}\delta_a{}^b \end{pmatrix}. \tag{5.3}$$

This leaves η_{AB} invariant and gives rise to

$$\check{\mathcal{H}}^{\check{A}\check{B}} = \mathcal{O}_{\check{C}}{}^{\check{A}}\mathcal{H}^{CD}\mathcal{O}_{\check{D}}{}^{\check{B}} = \begin{pmatrix} \kappa_{ab} & 0 \\ 0 & \kappa^{ab} \end{pmatrix}. \tag{5.4}$$

The transformed components of the structure coefficients become, after using the mCYBE,

$$\check{F}_{abc} = 0, \quad \check{F}_{ab}{}^c = \frac{1}{\sqrt{\eta}}f_{ab}{}^c, \quad \check{F}^{ab}{}_c = 0, \quad \check{F}^{abc} = \eta^{3/2}c^2\kappa^{ad}\kappa^{be}f_{de}{}^c. \tag{5.5}$$

Notice here we need not specify the value of c , and the following considerations will hold for all cases. The generalised curvature capturing the field equations for the metric and the B-field in this rotated frame reads

$$\check{\mathcal{R}}^{\check{A}\check{B}} = \frac{h^\vee(1-c^2\eta^2)^2}{4\eta} \begin{pmatrix} \kappa_{ab} & 0 \\ 0 & -\kappa^{ab} \end{pmatrix}. \quad (5.6)$$

After transforming back to the original frame, we obtain

$$\mathcal{R}^{AB} = \mathcal{O}_{\check{C}}^A \check{\mathcal{R}}^{\check{C}\check{D}} \mathcal{O}_{\check{D}}^B = \frac{h^\vee(1-c^2\eta^2)^2}{4} \begin{pmatrix} \kappa_{ab} & -\kappa_{ac}R^{cb} \\ R^{ac}\kappa_{cb} & -\frac{\kappa^{ab}}{\eta^2} - R^{ac}\kappa_{cd}R^{db} \end{pmatrix}. \quad (5.7)$$

This is consistent with the renormalisation of the sigma-model eq. (2.21) given by

$$\frac{\partial \mathcal{H}^{AB}}{\partial \log \mu} = \mathcal{R}^{AB} \quad \text{with} \quad \frac{\partial \eta}{\partial \log \mu} = \frac{h^\vee(1-c^2\eta^2)^2}{4}. \quad (5.8)$$

In the rotated frame the generalised Ricci scalar is quite easily calculated to be

$$\mathcal{R} = \frac{1}{6} (\eta^3 c^4 - 6\eta c^2 - 3\eta^{-1}) h^\vee \dim \mathfrak{g}. \quad (5.9)$$

There is no solution for $\mathcal{R} = 0$, the dilaton equation of DFT_{WZW} which holds for arbitrary η and c . However we may extend our considerations to include a direct sum of simple algebras

$$\mathfrak{g} = \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_N. \quad (5.10)$$

In this case, we can choose a different scaling for the inner product imposed on each simple factor \mathfrak{g}_i :

$$\kappa_{ab}^{(i)} = -\frac{1}{2h^\vee \alpha_i} f_{ac}^{(i)d} f_{bd}^{(i)c}. \quad (5.11)$$

In this way we will be able to engineer a cancelation of contributions to the curvature coming from each group factor, as is typical between AdS and internal factors of supergravity solutions. In principle we could have done this already for the simple case, but there such a scaling amounts to an trivial overall rescaling of the solution. For N simple factors, we have $N - 1$ additional degrees of freedom for which the dilaton field equation $\mathcal{R} = 0$ implies the constraint,

$$\sum_{i=1}^N \alpha_i h_i^\vee \dim \mathfrak{g}_i = 0. \quad (5.12)$$

This direct sum of algebras is however insufficient to solve $\mathcal{R}_{AB} = 0$. Hence, we conclude that in general there no setup which can solve the field equations without any contributions from the R/R sector. In order for the R/R sector to compensate the NS/NS contribution, we require (again in the rotated frame where equations are simplified),

$$\check{\mathcal{H}}_{\check{C}}^{\check{A}} \check{\mathcal{R}}^{\check{C}\check{B}} = \frac{h^\vee(1-c^2\eta^2)^2}{4\eta} \begin{pmatrix} 0 & -\delta_a^b \\ \delta_b^a & 0 \end{pmatrix} = -\frac{1}{8} \check{G}^T C \Gamma^{\check{A}\check{B}} \check{G}. \quad (5.13)$$

The field \check{G} is an eigenvector of $\check{\mathcal{K}}$ with eigenvalue -1 as required by (3.16) and of definite chirality. We discuss this condition in the following.

As the NS/NS sector, the R/R sector should also exhibit Poisson-Lie symmetry and in particular eq. (4.11) has to hold

$$\check{F}_{\check{A}\check{B}\check{C}}\Gamma^{\check{A}\check{B}\check{C}}\check{G} = 0, \tag{5.14}$$

at least if there are no sources like D-branes. Expanding this constraint into components, we obtain

$$\left(3f_{ab}{}^c\Gamma^{ab}\Gamma_c + c^2\eta^2\check{f}^{abc}\Gamma_{abc}\right)\check{G} = 0. \tag{5.15}$$

At this stage we should like to be explicit about solutions for \check{G} . To do so we found it convenient to recast our manipulations in terms of an $O(D)$ Drinfel'd $\check{\mathbf{G}}^{\underline{\alpha}\underline{\beta}}$ which can be related to \check{G} by a vectorisation map $\check{G} = \text{vec } \check{\mathbf{G}}$. The presentation of this somewhat technical procedure is detailed below and can be skipped on a first reading jumping instead to the explicit solution in the case of an example $\mathfrak{g} = \mathfrak{sl}(2) \oplus \mathfrak{su}(2)$.

Bispinorisation. The strategy will be to find a representation of \check{G} such that the self-duality and chirality constraints are automatically implemented and the only thing that remains to be taken care of is (5.15). We introduce the γ -matrices for the D -dimensional target space obeying the Clifford algebra,

$$\{\gamma_a, \gamma_b\} = 2\kappa_{ab}. \tag{5.16}$$

Assuming that D is even (which in our cases it shall be) they furthermore can be brought into the form,

$$(\gamma_a)^{\underline{\alpha}}{}_{\underline{\beta}} = \begin{pmatrix} 0 & (\gamma_a)_{\alpha\beta} \\ (\gamma_a)^{\alpha\beta} & 0 \end{pmatrix}, \quad \text{with} \quad (\gamma_{D+1})^{\underline{\alpha}}{}_{\underline{\beta}} = \begin{pmatrix} \delta_{\alpha}^{\beta} & 0 \\ 0 & -\delta_{\beta}^{\alpha} \end{pmatrix} \quad \text{and} \quad C_{\underline{\alpha}\underline{\beta}} = \begin{pmatrix} 0 & \delta_{\beta}^{\alpha} \\ -\delta_{\alpha}^{\beta} & 0 \end{pmatrix} \tag{5.17}$$

denoting the chirality and charge conjugation matrices. We express the 2^D components of \check{G} as a bispinor $\check{\mathbf{G}}^{\underline{\alpha}\underline{\beta}}$, where $\underline{\alpha}, \underline{\beta}, \dots$ are Dirac spinor indices which run from 0 to D . To get back and forth between these two representations, we use the vectorization

$$\check{G} = \text{vec}(\check{\mathbf{G}}) = \left(\check{\mathbf{G}}^{00}, \dots, \check{\mathbf{G}}^{D0}, \check{\mathbf{G}}^{D1}, \dots, \check{\mathbf{G}}^{DD}\right)^T. \tag{5.18}$$

The $O(D, D)$ Γ -matrices can now be written as

$$\Gamma_a = \frac{1}{\sqrt{2}}(\gamma_a \otimes 1 - i\gamma_{D+1} \otimes \gamma_a) \quad \text{and} \quad \Gamma^a = \frac{1}{\sqrt{2}}(\gamma^a \otimes 1 + i\gamma_{D+1} \otimes \gamma^a). \tag{5.19}$$

At first glance this new representation looks somewhat unwieldy. However, it has the advantage that the operator $\check{\mathcal{K}}$ and the chirality Γ_{2D+1} have a very convenient form:

$$\check{\mathcal{K}} = -(1 \otimes \gamma_{D+1}), \quad \Gamma_{2D+1} = (\gamma_{D+1} \otimes \gamma_{D+1}). \tag{5.20}$$

Remember that \check{G} has to be an eigenvector of $\check{\mathcal{K}}$ with eigenvalue -1 . Furthermore, it has to have a fixed chirality which also makes it an eigenvector of Γ_{2D+1} . The eigenvalue under

this operator decides whether we are capturing a type IIA or a IIB theory. In the bispinor representation solving these two constraints requires just to pick a particular subblock of $\check{\mathbf{G}}^{\alpha\beta}$, namely

block	$\check{\mathbf{G}}^{\alpha\beta}$	$\check{\mathbf{G}}^{\alpha}_{\beta}$	$\check{\mathbf{G}}_{\alpha}^{\beta}$	$\check{\mathbf{G}}_{\alpha\beta}$	
eigenvalue \mathcal{K}	+1	-1	+1	-1	(5.21)
eigenvalue Γ_{2D+1}	+1	-1	-1	+1.	

The condition encoding the Poisson-Lie symmetry (5.15) reads

$$[(3 + c^2\eta^2)(u \otimes 1 + i\gamma_{D+1} \otimes u) + 3(1 - c^2\eta^2)(\gamma^a \otimes u_a + iu_a\gamma_{D+1} \otimes \gamma^a)] \check{G} = 0, \quad (5.22)$$

with

$$u = f_{abc}\gamma^{abc} \quad \text{and} \quad u_a = f_{abc}\gamma^{bc}. \quad (5.23)$$

Note that because \check{G} has to be an eigenvector of both $\check{\mathcal{K}}$ and Γ_{2D+1} , two combinations of the terms in this constraint have to vanish individually. This leaves us with the two equations

$$\begin{aligned} [(3 + c^2\eta^2)(u \otimes 1) + 3(1 - c^2\eta^2)(\gamma^a \otimes u_a)] \check{G} &= 0, \\ [(3 + c^2\eta^2)(1 \otimes u) + 3(1 - c^2\eta^2)(u_a \otimes \gamma^a)] \check{G} &= 0. \end{aligned} \quad (5.24)$$

In the following, we do not want to discuss all solutions of these equations, but only the ones that have a chance to give rise to backgrounds which solve the field equations. To this end, we restrict our attention to \check{G} 's that are invariant under the action of \mathfrak{g} . More explicitly we impose

$$(u_a \otimes 1 + 1 \otimes u_a) \check{G} = 0. \quad (5.25)$$

Using this identity, (5.15) simplifies to

$$c^2\eta^2(u \otimes 1) \check{G} = 0 \quad \text{and} \quad c^2\eta^2(1 \otimes u) \check{G} = 0. \quad (5.26)$$

In particular, for the β -deformations for which $c^2 = 0$, the condition (5.25) is sufficient and in all other cases, we have to additionally impose the two constraints above (5.26).

In order to see what singles out these solutions, we have to take a closer look at the left hand side of R/R corrected field equation (5.13). To satisfy this equation the contributions from Γ^{ab} and Γ_{ab} to the left hand side vanish completely. Therefore, we just have to calculate the remaining:

$$\check{G}^T C \Gamma^a_b \mathcal{K} \check{G} = \check{G}^T (\gamma_{D+1} \gamma^1 \gamma^a \otimes \gamma_{D+1} \gamma^1 \gamma_b) \check{G} = \pm \text{Tr} \left((\check{G} \gamma^1 \gamma^a)^T \gamma^1 \gamma_b \check{G} \right). \quad (5.27)$$

This equation assumes a target space with Minkowski signature¹¹ with the time direction matching γ^1 and the $+/-$ depends on whether \check{G} is chiral/anti-chiral. Here, we have used the charge conjugation matrix on $O(D, D)$ spinors given by

$$C = i\gamma_{D+1} \gamma^1 \otimes \gamma_{D+1} \gamma^1. \quad (5.28)$$

¹¹For an Euclidean spacetime, we would just have to drop the γ^1 's.

For a simple Lie group the Killing metric is up to a scaling factor the only invariant bilinear form. But this implies that because \check{G} is invariant, the left hand side of (5.27) has to be proportional to δ_b^a . So the only thing we have to fix is the normalization of \check{G} . According to (5.13) it becomes,

$$\text{Tr} \left((\check{\mathbf{G}}\gamma^1\gamma^a)^T \gamma^1\gamma_a \check{\mathbf{G}} \right) = \mp \frac{h^\vee (1 - c^2\eta^2)^2 \dim \mathfrak{g}}{\eta}. \quad (5.29)$$

If we have more than one simple factor, like in (5.10), there are additional constraints on \check{G} :

$$\text{Tr} \left((\check{\mathbf{G}}\gamma^1\gamma^a)^T (P_i)^b{}_a \gamma^1\gamma_b \check{\mathbf{G}} \right) = \mp \frac{\alpha_i h_i^\vee (1 - c^2\eta^2)^2 \dim \mathfrak{g}_i}{\eta}, \quad (5.30)$$

where $(P_i)^b{}_a$ denotes a projector on the i th simple factor.

Example. Let us illustrate this procedure for deformations of $\text{AdS}_3 \times \text{S}^3$. In this particular case, the two relevant Lie algebras are

$$\mathfrak{g}_1 = \mathfrak{sl}(2) \quad \text{and} \quad \mathfrak{g}_2 = \mathfrak{su}(2) \quad \text{with} \quad h_1^\vee = h_2^\vee = 2, \quad \dim \mathfrak{g}_1 = \dim \mathfrak{g}_2 = 3. \quad (5.31)$$

In order to solve the field equation for the dilaton (5.12), we choose

$$\alpha_1 = 1 \quad \text{and} \quad \alpha_2 = -1. \quad (5.32)$$

This results in κ_{ab} of Minkowski signature, as required to describe $\text{AdS}_3 \times \text{S}^3$. A compatible R/R sector arises from (5.25). The corresponding R/R bispinor has the two solutions¹²

$$\check{\mathbf{G}}^{\alpha\beta} \sim \text{diag}(1, 1, 1, 1) \quad \text{and} \quad \check{\mathbf{G}}_{\alpha\beta} \sim \text{diag}(1, -1, -1, 1) \quad (5.33)$$

after restricting to the components of $\check{\mathbf{G}}$ with \mathcal{K} eigenvalue -1 . Only the second one solves the additional constraint (5.26), which is required for $c^2 \neq 0$. Furthermore, the first solution can not be normalized such that (5.29) is satisfied for both the $\mathfrak{sl}(2)$ and $\mathfrak{su}(2)$ factors. Thus, we conclude that for arbitrary c , there is only one R/R field configuration

$$\check{\mathbf{G}}_{\alpha\beta} = \frac{1}{\sqrt{2\eta}} (1 - c^2\eta^2) \text{diag}(1, -1, -1, 1) \quad (5.34)$$

that admits Poisson-Lie symmetry and in connection with the NS/NS sector solved all field equations. An alternative way to write this solution is

$$\check{G} = \frac{1 - c^2\eta^2}{12\sqrt{\eta}} f_{abc} \Gamma^{abc} |0\rangle, \quad (5.35)$$

where $|0\rangle$ denotes the vector which is annihilated by all Γ_a .

¹²We use the chiral γ -matrices

$$\begin{aligned} (\gamma_1)_{\alpha\beta} &= \begin{pmatrix} 0 & \sigma_1 \\ -\sigma_1 & 0 \end{pmatrix}, & (\gamma_2)_{\alpha\beta} &= \begin{pmatrix} i\sigma_2 & 0 \\ 0 & -i\sigma_2 \end{pmatrix}, & (\gamma_3)_{\alpha\beta} &= \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}, \\ (\gamma_4)_{\alpha\beta} &= \begin{pmatrix} 0 & -i\sigma_2 \\ i\sigma_2 & 0 \end{pmatrix}, & (\gamma_5)_{\alpha\beta} &= \begin{pmatrix} 0 & i\sigma_3 \\ -i\sigma_3 & 0 \end{pmatrix}, & (\gamma_6)_{\alpha\beta} &= \begin{pmatrix} -\sigma_2 & 0 \\ 0 & \sigma_2 \end{pmatrix}. \end{aligned}$$

They are conjugated by $\gamma_a^{\alpha\beta} = \epsilon^{\alpha\beta\gamma\delta} (\gamma_a)_{\gamma\delta}$ and give rise to the Killing metric $\delta_{ab} = \text{diag}(-1, 1, 1, 1, 1, 1)$. σ_i denotes the three Pauli matrices with $\sigma_i^2 = 1$ and $\epsilon^{\alpha\beta\gamma\delta}$ is totally anti-symmetric with $\epsilon^{1234} = 1$.

5.2 η -deformation redux

In this case the target space $\mathcal{M} = \mathbb{D}/\tilde{H}$ is equivalent to a group manifold H whose algebra corresponds to the direct sum of algebras introduced in eq. (5.10). Parametrising \mathcal{M} by a group element $g \in H$ (with e and v corresponding left and right Maurer-Cartan forms respectively and $M_A{}^B \mathbb{T}_B = g \mathbb{T}_A g^{-1}$) we have from theorem 1 the generalised frame field

$$\hat{E}_A{}^{\hat{I}} = \begin{pmatrix} e^a{}_i & \Pi^{ab} e_b{}^i \\ 0 & e_a{}^i \end{pmatrix}, \quad (5.36)$$

in which we recall $\Pi^{ab} = M^{ac} M^b{}_c$. The target space metric and the B -field are readily extracted from $\hat{\mathcal{H}}^{\hat{I}\hat{J}} = \hat{E}_A{}^{\hat{I}} \mathcal{H}^{AB} \hat{E}_B{}^{\hat{J}}$ and read¹³

$$\begin{aligned} ds^2 &= g_{ab} e^a e^b = \eta \kappa_{ab} e^a \otimes e^b - \frac{\eta^3}{1 + \eta^2} \beta^{ab} \beta^{cd} \kappa_{bd} \kappa_{ae} \kappa_{cf} e^e \otimes e^f, \\ B &= \frac{\eta^2}{2(1 + \eta^2)} \left(\kappa_{ac} \beta^{cd} \kappa_{db} e^a \wedge e^b \right). \end{aligned} \quad (5.37)$$

Here we have used the β -parametrisation of the generalised metric

$$\hat{\mathcal{H}}^{\hat{I}\hat{J}} = \begin{pmatrix} e^a{}_i & 0 \\ 0 & e_a{}^i \end{pmatrix} \begin{pmatrix} \tilde{g}_{ab} & \tilde{g}_{ac} \beta^{cb} \\ -\beta^{ac} \tilde{g}_{cb} & \tilde{g}^{ab} - \beta^{ac} \tilde{g}_{cd} \beta^{db} \end{pmatrix} \begin{pmatrix} e^b{}_j & 0 \\ 0 & e_b{}^j \end{pmatrix}, \quad (5.38)$$

of the generalised metric for which

$$\tilde{g}_{ab} = \eta \delta_{ab} \quad \text{and} \quad \beta^{ab} = \Pi^{ab} - R^{ab} = -M^a{}_c M^b{}_d R^{cd}. \quad (5.39)$$

In this parametrisation the metric and the B -field in flat indices arise from inverting

$$\left(\frac{1}{\eta} \kappa^{-1} - \beta \right)_{ab}^{-1} = g_{ab} - B_{ab}, \quad (5.40)$$

whereas the curved version are obtained after contraction with $e^a{}_i$. A comparison with the action (2.9) gives rise to

$$(G_0 - B_0)^{-1 ab} = \tilde{g}^{ab} + \beta^{ab} - \Pi^{ab} = \frac{1}{\eta} \kappa^{ab} - R^{ab} \quad (5.41)$$

which is equivalent to (2.22). The dilaton is determined by the Poisson-Lie condition (4.7) and has to have the form

$$\phi = \phi_0 + \frac{1}{4} \log |\det g_{ij}| - \frac{1}{2} \log |\det e^a{}_i| = \phi_0 + \frac{1}{4} \log |\det g_{ab}|, \quad (5.42)$$

where according to (4.7), ϕ_0 is a free constant.

¹³The overall factor of η in front of the metric may look unfamiliar but the reader should recall that the normalisation \tilde{k} of the \mathcal{E} -model, in which the DFT equations are perturbative is related to the normalisation of the sigma-model by a corresponding factor $\frac{1}{\eta}$.

Since in general \tilde{H} will be non-unimodular we will have solutions of modified type II SUGRA encoded in the DFT vector

$$\mathbf{X}^{\hat{I}} = \frac{1}{2} \begin{pmatrix} R^{bc} f_{bc}{}^a v_a{}^i \\ 0 \end{pmatrix}, \quad (5.43)$$

from which the Killing vector I of modified supergravity is

$$I^i = -\frac{1}{2} R^{bc} f_{bc}{}^a v_a{}^i = -\frac{1}{2} \beta^{bc} f_{bc}{}^a e_a{}^i. \quad (5.44)$$

To complete the NS sector specification of the modified type II SUGRA one also needs the one-form Z defined in eq. (3.58) which gives rise to

$$Z = \frac{1}{4} g^{ab} dg_{ab} + \iota_I B. \quad (5.45)$$

Finally, we have to obtain the R/R fluxes. To this end, we begin with a solution \check{G} of Poisson-Lie condition (5.15) evaluated in the simplified rotated frame and then calculate

$$\sum_{p=1} \frac{1}{p! 2^{p/2}} \hat{G}_{a_1 \dots a_p}^{(p)} \Gamma^{a_1 \dots a_p} = \sqrt{\eta} S_\beta \check{G} = \sqrt{\eta} \exp\left(\frac{1}{4} \beta^{ab} \Gamma_{ab}\right) \check{G} \quad (5.46)$$

in flat indices and again contract with $e^a{}_i$ to get the curved versions. Converting this into a polyform one can construct the fluxes $\hat{\mathcal{F}} = e^\phi e^{-B} \hat{G}$ which obey the flux equations and Bianchi identities $\mathbf{d}\hat{\mathcal{F}} = 0$ with \mathbf{d} the modified exterior derivative of eq. (3.57).

An example is the $\text{AdS}_3 \times \text{S}^3$ from the last section for which a realization of the Drinfel'd double provided in appendix G. However there is no need to resort here to an explicit parametrisation since the geometry can be entirely written in terms of e^a and β^{ab} whose exterior derivatives are easily obtained as

$$de^a = -\frac{1}{2} f_{bc}{}^a e^b \wedge e^c, \quad d\beta^{ab} = 2f_{cd}{}^{[a} \beta^{cb]} e^d. \quad (5.47)$$

The metric, B -field and vector I are already given in the simple forms above and in addition we have that for this example the dilaton is constant and

$$\phi = \log\left(\frac{\eta^{3/2}}{1 + \eta^2}\right) + \phi_0, \quad H = dB = 0, \quad Z = 0. \quad (5.48)$$

Evaluating (5.46) for the solution given in eq. (5.35), gives rise to

$$\hat{G}^{(1)} = -\frac{1 + \eta^2}{\sqrt{2}} \beta^{ab} f_{abc} e^c, \quad \hat{G}^{(3)} = \frac{1 + \eta^2}{3\sqrt{2}} f_{abc} e^a \wedge e^b \wedge e^c. \quad (5.49)$$

At this stage the preceding discussion establishes that we have a solution of modified supergravity, or rather a six-dimensional truncated version thereof. As a consistency check and for completeness we provide details of the uplift to a full ten-dimensional solution in appendix F.

5.3 λ -deformation redux

We now describe the λ -model in this framework. Here the underlying Double is formed from $\mathfrak{d} = \mathfrak{g} + \mathfrak{g}$ with the maximal isotropic subgroup \tilde{H} being the diagonal embedding $G_{\text{diag}} \subset \mathbb{D}$ (see B for further details), whose generators are \tilde{T}^a in the canonical basis. In this case however the complementary isotropic does not form a subgroup as can be seen from the structure constants of \mathfrak{d} given in terms of those of \mathfrak{g} by

$$F_{ab}{}^c = 0, \quad F_{abc} = \frac{1}{\sqrt{2}} f_{abc} = \frac{1}{\sqrt{2}} f_{ab}{}^f \kappa_{fc}, \quad \tilde{F}^{abc} = 0, \quad \tilde{F}^{ab}{}_c = \frac{1}{\sqrt{2}} f^{ab}{}_c = \frac{1}{\sqrt{2}} \kappa^{ad} \kappa^{be} f_{de}{}^f \kappa_{fc}, \quad (5.50)$$

with others given by symmetry enforced by the ad-invariance of η . This algebra admits a \mathbb{Z}_2 grading so that $\mathcal{M} = \mathbb{D}/\tilde{H} = (G \times G)/G_{\text{diag}}$ is a symmetric space and we can apply the construction of theorem 1 to obtain generalised frame fields that describe the geometry. To do so requires some care however in the parametrisation of coset representatives that we now explain.

Before doing so let us mention the alternative route to describe the λ -deformation as the Poisson-Lie T-duality of the η -model and a subsequent analytic continuation. This can be made quite manifest at the level of \mathcal{E} -models [29] and therefore in DFT_{WZW}. It is worthwhile briefly recasting this argument in the language we use here by identifying a frame where the double $\mathfrak{d} = \mathfrak{g}^{\mathbb{C}}$ decomposes, up to an analytic continuation, into $\mathfrak{d} = \tilde{\mathfrak{h}} \oplus \mathfrak{k} = \mathfrak{g}_{\text{diag}} \oplus \mathfrak{g}_{\text{antidiag}}$. Starting from the frame (5.5) we perform a rotation

$$\mathcal{O}_{A'}{}^{\check{B}} = \begin{pmatrix} 0 & \kappa^{ab} \\ \kappa_{ab} & 0 \end{pmatrix}, \quad (5.51)$$

to obtain structure coefficients $\mathbb{F}'_{A'B'C'} = \mathcal{O}_{A'}{}^{\check{D}} \mathcal{O}_{B'}{}^{\check{E}} \mathcal{O}_{C'}{}^{\check{F}} \check{\mathbb{F}}_{\check{D}\check{E}\check{F}}$ that have components

$$F'_{abc} = -\eta^{3/2} f_{ab}{}^d \kappa_{dc}, \quad F'_{ab}{}^c = 0, \quad F'^{ab}{}_c = \frac{1}{\sqrt{\eta}} \kappa^{ad} \kappa^{be} \kappa_{cf} f_{de}{}^f, \quad F'^{abc} = 0. \quad (5.52)$$

One can write down “generators” for the commutation relations (5.52),

$$T_a = \sqrt{\eta} \{-it_a, it_a\} \quad \text{and} \quad \tilde{T}^a = \frac{\kappa^{ab}}{\sqrt{\eta}} \{t_b, t_b\}, \quad t_a \in \mathfrak{g}, \quad (5.53)$$

however we see that the generators here are not anti-hermitian and hence an analytic continuation

$$T_a \rightarrow iT_a \quad (5.54)$$

must be taken in order to match (up to scaling) the structure of $\mathfrak{d} = \mathfrak{g} + \mathfrak{g}$.

We now resume the construction of the generalised frame fields. We must set the representative m for coset \mathbb{D}/\tilde{H} . To be explicit we make the choice of parameterisation of the coset representative¹⁴ $m = \{\bar{g}, \bar{g}^{-1}\}$ with $\bar{g} \in G$. However to match directly to the λ -model of eq. (2.24) which is parametrised by a group element g , a further identification

¹⁴Here we deviate from [29] in which the coset representative is chosen as $m = \{g, e\}$, the reason will be that this choice is the one that matches the parametrisations used in theorem 1.

is needed namely that $\bar{g}^2 = g$. We let $\bar{e}, \bar{v}, \bar{D}$ be the left/right-invariant forms and adjoint action on \mathfrak{g} constructed from \bar{g} which can be related to those constructed from g via,

$$e^a{}_i = (1 + \bar{D}^{-T})^a{}_b \bar{e}^b{}_i, \quad \bar{D}^2 = D. \quad (5.55)$$

The adjoint action of m on \mathfrak{d} , i.e. $m\mathbb{T}_A m^{-1} = M_A{}^B \mathbb{T}_B$, is given by

$$M_A{}^B = \frac{1}{2} \begin{pmatrix} \kappa^{-1} X^+ \kappa & \kappa^{-1} X^- \\ X^- \kappa & X^+ \end{pmatrix}, \quad X^\pm = \bar{D} \pm \bar{D}^{-1}, \quad (5.56)$$

and is easily seen to be an $O(D, D)$ element preserving η_{AB} . The one-form dmm^{-1} determines a veilbein and \mathfrak{h} -valued connection according to

$$dmm^{-1} = V^A{}_i \mathbb{T}_A dx^i = T_a v^a{}_i dx^i + \tilde{T}^a A_{ai} dx^i, \quad (5.57)$$

with

$$v^a{}_i = \frac{1}{\sqrt{2}} (\bar{v} + \bar{e})^a{}_i, \quad A_{ai} = \frac{1}{\sqrt{2}} \kappa_{ab} (\bar{v} - \bar{e})^b{}_i. \quad (5.58)$$

From these we can construct a two-form

$$\omega^{(2)} = \frac{1}{2} v^a{}_i A_{aj} dx^i \wedge dx^j = \frac{1}{4} (\kappa \bar{D}^T - \bar{D} \kappa)_{ab} \bar{e}^a \wedge \bar{e}^b, \quad (5.59)$$

such that

$$d\omega^{(2)} = -\frac{1}{4} f_{abc} (\bar{e}^a \wedge \bar{e}^b \wedge \bar{v}^c + \bar{v}^a \wedge \bar{v}^b \wedge \bar{e}^c). \quad (5.60)$$

In addition there is a globally defined three from

$$\Omega^{(3)} = \frac{1}{12} \langle\langle dmm^{-1}, [dmm^{-1}, dmm^{-1}] \rangle\rangle = \frac{1}{6} f_{abc} \bar{e}^a \wedge \bar{e}^b \wedge \bar{e}^c, \quad (5.61)$$

for which *locally* we can introduce a suitable potential $d\Omega^{(2)} = \Omega^{(3)}$.

Then in theorem 1 we have in combination

$$\rho = \omega^{(2)} - \Omega^{(2)} \rightarrow d\rho = -\frac{1}{12} f_{abc} e^a \wedge e^b \wedge e^c \quad (5.62)$$

giving the three form H_{WZW} that comes from $S_{WZW}[g]$.

We now have all the ingredients to introduce a generalised frame field, itself also an $O(d, d)$ element. We, as per comment 2, will strip out the **H**-flux contribution given by ρ and consider the frame field

$$\hat{E}'_A{}^{\hat{I}} = \frac{1}{2} \begin{pmatrix} \frac{1}{\sqrt{2}} \kappa^{-1} (1 + D) \kappa & \sqrt{2} \kappa^{-1} (1 - D) \\ \frac{1}{\sqrt{2}} (1 - D) \kappa & \sqrt{2} (1 + D) \end{pmatrix}_A{}^B \begin{pmatrix} e & 0 \\ 0 & e^{-T} \end{pmatrix}_B{}^{\hat{I}}. \quad (5.63)$$

in which we have massaged the expressions such that they only depend on the left/right-invariant forms and adjoint action on \mathfrak{g} constructed from g so as to match the parametrisation of the λ -model of eq. (2.24). A useful decomposition of the frame field is given by

$$\hat{E}'_A{}^{\hat{I}} = \begin{pmatrix} r^{-T} & 0 \\ 0 & r \end{pmatrix} \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix} \begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix} \begin{pmatrix} e & 0 \\ 0 & e^{-T} \end{pmatrix} \quad (5.64)$$

with

$$r^{-T} = \frac{1}{2\sqrt{2}} (1 + D^{-T}), \quad b = \frac{1}{8}(D^{-1} - D)\kappa \quad \text{and} \quad \beta = 2\kappa^{-1} \frac{1 - D}{1 + D}. \quad (5.65)$$

From the definition of the operator \mathcal{E} in the λ -model we have the generalised metric in flat space given by

$$\mathcal{H}_{AB} = \begin{pmatrix} \epsilon^{-\frac{1}{2}}\kappa^{ab} & 0 \\ 0 & \epsilon^{\frac{1}{2}}\kappa_{ab} \end{pmatrix}, \quad \epsilon^{\frac{1}{2}} = \frac{1 - \lambda}{1 + \lambda}, \quad (5.66)$$

from which as usual we construct the curved space generalised metric. From this the metric and B-field are readily extracted as

$$\begin{aligned} ds_\lambda^2 &= \frac{1}{2} ((O_{g^{-1}} + O_g - 1)\kappa)_{ab} e^a \otimes e^b, \\ B_\lambda &= B_{\text{WZW}} + \frac{1}{4} (O_{g^{-1}} - O_g)_{ab} e^a \wedge e^b, \end{aligned} \quad (5.67)$$

in which $H_{\text{WZW}} = dB_{\text{WZW}}$ and

$$O_g = (1 - \lambda D)^{-1}. \quad (5.68)$$

As in our previous discussion, and detailed in appendix E, the PL conditions on the dilaton determine that

$$\phi_\lambda = \phi_0 - \frac{1}{2} \log \det(1 - \lambda D^{-1}), \quad (5.69)$$

in which ϕ_0 is constant. This matches the dilaton obtained due to a Gaussian elimination of fields in the construction of [16].

Since \mathfrak{g} is assumed to be unimodular we have that $\tilde{\mathfrak{h}}$ is also unimodular, and thus we are in the situation of conventional (not modified) supergravity. What remains is to determine the RR fluxes from the PL conditions.

To be totally analogous with the discussion of the η deformations we should actually further perform an $O(D, D)$ rotation

$$\mathcal{O}_{\check{A}}^B = \begin{pmatrix} \epsilon^{\frac{1}{4}}\delta^{ab} & 0 \\ 0 & \epsilon^{-\frac{1}{4}}\delta_a^b \end{pmatrix}, \quad (5.70)$$

such that the structure constants become

$$\check{F}_{ab}{}^c = 0, \quad \check{F}_{abc} = \frac{\epsilon^{-\frac{3}{4}}}{\sqrt{2}} f_{abc}, \quad \check{\check{F}}^{abc} = 0, \quad \check{\check{F}}^{ab}{}_c = \frac{\epsilon^{\frac{1}{4}}}{\sqrt{2}} f^{ab}{}_c. \quad (5.71)$$

In this frame the PL condition invokes

$$\left(\check{\check{\Gamma}}^{abc} f_{abc} + 3\epsilon f^{ab}{}_c \check{\check{\Gamma}}^{ab}{}^c \right) \check{G} = 0. \quad (5.72)$$

Notice here we have a direct similarity between the same constraint found in the η branch. Indeed, comparing this with eq. (5.15) we simply have to swap the index positions, up and down, (corresponding to the rotation eq. (5.51)), and identify $\lambda = (\eta + i)(\eta - i)^{-1}$. Thus

knowing already how to solve this equation we can construct the curved space spinor R/R field strengths via

$$\widehat{G}' = \sum_{p=1} \frac{1}{p!2^{p/2}} \widehat{G}'_{a_1 \dots a_p} \Gamma^{a_1 \dots a_p} = S_\beta S_b S_r S_{\mathcal{O}} \check{G}, \quad (5.73)$$

in flat indices and the final result arise after contraction with the left-invariant from e^a . Note that here we are dressing with the spinor representative of \widehat{E}' in which the B -shift induced by ρ has been stripped off. To get the raw \widehat{G} that obey $d\widehat{G} = 0$ one must apply the spinor representative of a B -shift for B_{WZW} .

Example. Let us now come again to the $\text{SL}(2) \times \text{SU}(2)$ example. Like for the η -deformation in the last section, it is convenient to express all supergravity fields just in terms of the invariant tensors of the Lie algebra \mathfrak{g} and the adjoint action D_{ab} . From this point of view β^{ab} is the only quantity in (5.73) which also includes the inverse of $1 + D$. However, it always appears in either one of the two combinations

$$r^a{}_c \beta^{cb} = \frac{1}{\sqrt{2}} (\kappa^{ab} - D^{ab}) \quad \text{or} \quad \beta_{ac} b^{cb} = -\frac{1}{2} \delta_a^b + \frac{1}{4} (D_a{}^b + D^b{}_a). \quad (5.74)$$

At this stage we find it convenient to pick a particular representation for both $\text{SU}(2)$ and $\text{SL}(2)$ elements:

$$g_{\text{SL}(2)} = \begin{pmatrix} \cosh \alpha_1 - \sinh \alpha_1 \cosh \alpha_2 & -\sinh \alpha_1 \sinh \alpha_2 e^{\alpha_3} \\ \sinh \alpha_1 \sinh \alpha_2 e^{-\alpha_3} & \cosh \alpha_1 + \sinh \alpha_1 \cosh \alpha_2 \end{pmatrix}, \quad (5.75)$$

$$g_{\text{SU}(2)} = \begin{pmatrix} \cos \beta_1 + i \sin \beta_1 \cos \beta_2 & \sin \beta_1 \sin \beta_2 e^{-i\beta_3} \\ -\sin \beta_1 \sin \beta_2 e^{i\beta_3} & \cos \beta_1 - i \sin \beta_1 \cos \beta_2 \end{pmatrix}.$$

Here α_2 will become the time-like direction. To reduce paper we write $c_i = \cos \beta_i$ and $\text{ch}_i = \cosh \alpha_i$ etc. The line element is given by

$$ds^2 = \frac{1+\lambda}{1-\lambda} d\alpha_1^2 + \frac{(1-\lambda^2)\text{sh}_1^2}{\Delta} (-d\alpha_2^2 + \text{sh}_2^2 d\alpha_3^2) + \frac{1+\lambda}{1-\lambda} d\beta_1^2 + \frac{(1-\lambda^2)s_1^2}{\Theta} (d\beta_2^2 + s_2^2 d\beta_3^2), \quad (5.76)$$

and the B -field and dilaton

$$B = B_{\text{WZW}} + B_0, \quad B_0 = 4\lambda\Delta^{-1} \text{ch}_1 \text{sh}_1^3 \text{sh}_2 d\alpha_2 \wedge d\alpha_3 + 4\lambda\Theta^{-1} c_1 s_1^3 s_2 d\beta_2 \wedge d\beta_3$$

$$B_{\text{WZW}} = 2\text{sh}_1^2 \text{sh}_2 \alpha_3 d\alpha_1 \wedge d\alpha_2 + 2s_1^2 s_2 \beta_3 d\beta_1 \wedge d\beta_2, \quad \Phi = \phi_0 - \frac{1}{2} \log \Delta \Theta, \quad (5.77)$$

with

$$\Delta = 1 + \lambda^2 - 2\lambda \text{ch}_1, \quad \Theta = 1 + \lambda^2 - 2\lambda c_1. \quad (5.78)$$

We can apply directly eq. (5.73) and after some work we find

$$\widehat{G}'^{(1)} = n (\text{sh}_1 c_1 d\alpha_1 - \text{ch}_1 s_1 d\beta_1)$$

$$\widehat{G}'^{(3)} = n \left(s_1 \text{sh}_1^3 \text{sh}_2 d\alpha_2 \wedge d\alpha_3 \wedge d\beta_1 - c_1 \text{ch}_1 s_1^2 \text{sh}_2 d\alpha_1 \wedge d\alpha_2 \wedge d\alpha_3 \right. \\ \left. - s_1^3 s_2 \text{sh}_1 d\alpha_1 \wedge d\beta_2 \wedge d\beta_3 - c_1 \text{ch}_1 s_1^2 s_2 d\beta_1 \wedge d\beta_2 \wedge d\beta_3 \right)$$

$$\widehat{G}'^{(5)} = n \left(\text{ch}_1 s_1^3 s_2^2 d\alpha_1 \wedge d\alpha_2 \wedge d\alpha_3 \wedge d\beta_2 \wedge d\beta_3 + c_1 s_1^2 s_2^2 \text{sh}_3^3 \text{sh}_2 d\alpha_2 \wedge d\alpha_3 \wedge d\beta_1 \wedge d\beta_2 \wedge d\beta_3 \right), \quad (5.79)$$

which have to be twisted with the WZW contribution to the B-field in order to obtain $\widehat{G} = e^{B_{\text{WZW}}} \sum_{n=1}^3 \widehat{G}'^{(2n-1)}$ which satisfies the Bianchi identity $d\widehat{G} = 0$. All that remains to be fixed is the normalisation $n^2 = \frac{\lambda^2}{8} e^{-2\phi_0}$. One can lift this to a solution in 10d by adding an auxiliary T^4 exactly as is done in the appendix F.

6 Conclusion

In this work we have continued the development of Poisson-Lie T-duality, based on a Drinfel'd double, by describing explicitly its embedding into DFT. We are further able to extend these ideas to include so-called \mathcal{E} -models (which incorporate e.g. integrable λ -deformed theories), the required Drinfel'd double is relaxed to that of a Manin quasi-triple. In either case we are able to extend the conditions for Poisson-Lie symmetry or an \mathcal{E} -model to the dilaton, recovering rather simply a result that took some effort in the mathematical literature, and to the R/R sector which to date had been treated in a somewhat ad-hoc fashion.

The condition of having an \mathcal{E} -model can be understood in the context of DFT_{WZW} defined on a $2D$ -dimensional group manifold \mathbb{D} of an algebra \mathfrak{d} as demanding invariance under the $2D$ diffeomorphism symmetry. Choosing a solution to the section condition amounts to finding a subalgebra $\tilde{\mathfrak{h}} \in \mathfrak{d}$, reducing the dynamics to that defined on the coset $\mathcal{M} = \mathbb{D}/\tilde{H}$. When $\tilde{\mathfrak{h}}$ is unimodular, the equations of motion of regular DFT (within a solution to the section condition) are recovered, and when non-unimodular those corresponding to a known modification of DFT (and SUGRA) are found. In this way, for the backgrounds we are considering, the equations of motion for DFT_{WZW} become algebraic and match those derived some years ago from the context of the doubled worldsheet in the bosonic sector. Similarly we are able to reduce the considerations of the R/R-sector to essentially an algebraic problem.

Critical is that we are able to construct generalised frame fields that are $O(D, D)$ valued and moreover close under the generalised Lie derivative to generate structure constants of the Lie-algebra \mathfrak{d} . This allows to translate the algebraic results for the DFT_{WZW} defined on \mathbb{D} to conventional target space fields on the physical spacetime \mathcal{M} . We demonstrate this technology with examples corresponding to η and λ integrable deformations. This is a quite satisfying result, viewed as conventional geometries these deformations look anything but simple, whereas in this language their underlying simplicity becomes transparent. It seems plausible that more general integrable sigma models [76–79] may give rise to such a structure.

One of the original motivations for this study was to see if by using DFT_{WZW} we could resolve the long standing questions concerning the global properties of non-Abelian duality transformations. At first sight things seem promising since we have an underlying doubled group manifold \mathbb{D} . However, a closer look shows this is not quite the full story. Firstly our constructions only make use of the algebra \mathfrak{d} , additional input is required to specify the global structure of \mathbb{D} , e.g. there may be discrete quotients to be taken. A second challenge is that we assume a factorisation of the group elements of the form $g(X) = \tilde{h}(\tilde{x})m(x)$ with particular parametrisation of $m(x)$ as the exponent of coset generators. Neither the factorisation of g nor the identity-connectedness of m are guaranteed to hold globally.

Thus further work is needed to establish the patching required to extend our construction. However, suppose that this procedure can be completed and our frames given a global definition. In that case we would have specified a generalised parallelisation for \mathcal{M} and would provide an explicit demonstration of the fact [47] that the reduction on such spaces constitutes a consistent truncation [80, 81].

This work prompts many interesting directions. The most obvious is to describe the dressing coset procedure in this language (something that we intend to report on shortly [82]), and eventually the extension to semi-symmetric spaces with the application to the full $\text{AdS}_5 \times S^5$ superstring in mind. In these more general cases we also intend to detail the question of supersymmetry, by making manifest the idea that whilst naively broken in conventional SUGRA it is recovered in DFT by allowing Killing spinors to have dependence on the ‘dual’ coordinates [83, 84]. In the present work we also showed that the PL T-duality rule on R/R fields can be recast in the format of a Fourier-Mukai transformation, something which was known to be the case for Abelian T-duality [74]. It is well known that D-brane charges admit a K-theory classification [85, 86] and that this Fourier-Mukai transformation can be understood as implementing T-duality at the level of K-theory. So one might (optimistically perhaps given the state of knowledge of global properties) hope to understand the Poisson-Lie transformation at this level. Looking further ahead the prospect of using the algebraic description of these backgrounds to study higher order corrections [87–89] is enticing as is the interpretation of such generalised dualities in the context of the Exceptional Field Theory approach to M-theory.

Acknowledgments

We gratefully acknowledge B. Hoare, F. Rudolph and C. Strickland-Constable for helpful discussions. FH and SD would like to thank the organisers of Corfu Summer Institutes 2018 where some results of this work were presented, and the COST network for supporting the visit of FH to this workshop. The work of DCT is supported by a Royal Society University Research Fellowship URF 150185, and in part by the STFC grant ST/P00055X/1. FH is partially supported by the Spanish Government Research Grant FPA2015-63667-. SD is supported by the FWO-Vlaanderen through aspirant fellowship and in part through the project FWO G020714N.

A Conventions and notation

There are many different groups, algebras, subgroups and subalgebras encountered in this paper — we list the main definitions in table 2. Commensurate with this is an abundance of indices outlined in table 3.

Sigma-models and supergravity. We consider 2d non-linear sigma models in Lorentzian signature given by

$$S = \frac{1}{\pi s} \int d\sigma d\tau \partial_+ X^i (G(X) - B(X))_{ij} \partial_- X^j, \quad (\text{A.1})$$

	Double \mathfrak{d}	Lagrangian subalg. $\tilde{\mathfrak{h}}$	Lagrangian compl. \mathfrak{k}	Algebra \mathfrak{g} :
Exponentiation	\mathcal{D}	\tilde{H}	$\exp(\mathfrak{k})/H$ for \mathfrak{k} subsp./subalg.)	G :
Algebra gen.	\mathbb{T}_A	\tilde{T}^a	T_a	t_a :
Inner product	$\langle\langle \mathbb{T}_A, \mathbb{T}_B \rangle\rangle = \eta_{AB}$			$\langle T_a, T_b \rangle = \kappa_{ab}$:
Structure cst	$\mathbb{F}_{AB}{}^C$	$\tilde{F}^{ab}{}_c$	$F_{ab}{}^c$	$f_{ab}{}^c$:
Group element	$\mathfrak{g}(X^I)$	$\tilde{h}(\tilde{x}^{\tilde{i}})$	$m(x^i)$	g : $\tilde{g}^2 = g$
Adjoint action	$\mathbb{M}_A{}^B$	\tilde{M}^{ab}	$M_a{}^b$	$D_a{}^b$: $\bar{D}_a{}^b$
L/R MC forms	$\mathbb{E}^A{}_I/\mathbb{V}^A{}_I$	$\tilde{E}^{A\tilde{i}}/\tilde{V}^{A\tilde{i}}$	$E^A{}_i/V^A{}_i$:
L/R MC comps		$\tilde{e}_a{}^{\tilde{i}}/\tilde{v}_a{}^{\tilde{i}}$	$e^a{}_i/v^a{}_i/A_{ai}$	$\underline{e}^a{}_i/\underline{v}^a{}_i$: $\bar{e}^a{}_i/\bar{v}^a{}_i$

Table 2. The variety of algebras, groups and group elements used.

	tensor	indices
Flat frame	T_A	$A, B, C, \dots = 1 \dots 2D$
Rotated flat frame	$\check{T}_{\check{A}}$	$A, B, C, \dots = 1 \dots 2D$
Doubled curved space	T_I	$I, J, K, \dots = 1 \dots 2D$
Generalised tangent space	$\hat{T}_{\hat{I}}$	$\hat{I}, \hat{J}, \hat{K}, \dots = 1 \dots 2D$

Table 3. The variety of indices used.

in which $\partial_{\pm} = \frac{1}{2}(\partial_{\tau} \pm \partial_{\sigma})$. This sign choice for the NS two-form field means that for a constant G and B the Hamiltonian

$$\text{Ham} = \dot{X}^i P_i - L = \frac{1}{4\pi s} \mathcal{Z}^M \mathcal{H}_{MN} \mathcal{Z}^N, \quad \mathcal{Z}^M = (2\pi s P_i, \partial_{\sigma} X^i), \quad (\text{A.2})$$

is written with the generalised metric defined as

$$\mathcal{H}_{MN} = \begin{pmatrix} G^{-1} & -G^{-1}B \\ BG^{-1} & G - BG^{-1}B \end{pmatrix}. \quad (\text{A.3})$$

The NS sector supergravity equations are given by (for type IIB)

$$\begin{aligned} 0 &= R_{mn} + 2\nabla_{mn}\Phi - \frac{1}{4}H_{mpq}H_n{}^{pq} \\ &\quad - e^{2\Phi} \left(\frac{1}{2}(F_1^2)_{mn} + \frac{1}{4}(F_3^2)_{mn} + \frac{1}{96}(F_5^2)_{mn} - \frac{1}{4}g_{mn} \left(F_1^2 + \frac{1}{6}F_3^2 \right) \right), \\ 0 &= d[e^{-2\Phi} \star H] + F_1 \wedge \star F_3 + F_3 \wedge F_5, \\ 0 &= R + 4\nabla^2\Phi - 4(\partial\Phi)^2 - \frac{1}{12}H^2, \end{aligned} \quad (\text{A.4})$$

in which $H = dB$. For the R/R fields we have Hodge duals defined (in $d = 10$ dimensions) according to $F_{(p)} = -(-1)^{p(p+1)/2} \star F_{(d-p)}$ for which the poly-form $\mathcal{F} = \sum_p F_{(p)}$ obeys

$$d_H \mathcal{F} = (d + H \wedge) \mathcal{F} = 0. \quad (\text{A.5})$$

The Hodge star operator is such that $\star^2\omega_{(p)} = s(-1)^{p(d-p)}\omega_{(p)}$, where s is the signature. For IIB we have

$$\mathcal{F} = F_{(1)} + F_{(3)} + F_{(5)} - \star F_{(3)} + \star F_{(1)}, \quad F_{(5)} = \star F_{(5)}. \quad (\text{A.6})$$

We occasionally also use $\mathcal{F} = e^{-B}\mathcal{G}$ and $\mathcal{F} = e^{\Phi}\mathcal{F}$.

In modified supergravity [48] we have instead

$$\begin{aligned} 0 &= R_{mn} - \frac{1}{4}H_{mpq}H_n{}^{pq} + \nabla_m X_n + \nabla_n X_m \\ &\quad - \left(\frac{1}{2}(\mathcal{F}_1^2)_{mn} + \frac{1}{4}(\mathcal{F}_3^2)_{mn} + \frac{1}{96}(\mathcal{F}_5^2)_{mn} - \frac{1}{4}g_{mn} \left(\mathcal{F}_1^2 + \frac{1}{6}\mathcal{F}_3^2 \right) \right), \\ 0 &= d\star H + \mathcal{F}_1 \wedge \star \mathcal{F}_3 + \mathcal{F}_3 \wedge \mathcal{F}_5 - 2\star dX - 2X \wedge \star H, \\ 0 &= R + 4\nabla_n X^n - 4X_n X^n - \frac{1}{12}H^2, \\ 0 &= \mathbf{d}\mathcal{F} \equiv (d + H \wedge - Z \wedge - \iota_I)\mathcal{F}. \end{aligned} \quad (\text{A.7})$$

Here the vector X is given by

$$X = Z + I, \quad (\text{A.8})$$

with the constraints

$$dZ + \iota_I H = 0, \quad \iota_I Z = 0, \quad L_I g = L_I H = 0. \quad (\text{A.9})$$

For the case of $I = 0$ we have that $X = d\phi$ and the conventional supergravity is recovered. In general we identify the ‘‘dilaton’’ as the exact piece of Z ;

$$Z = d\phi + \iota_I B - V, \quad L_I B = dV. \quad (\text{A.10})$$

In these equations we use the interior contraction defined as $\iota_{\mathcal{I}}\omega = \star(\mathcal{I} \wedge \star\omega)$ and recall $L_{\mathcal{I}}\omega = d\iota_{\mathcal{I}}\omega + \iota_{\mathcal{I}}d\omega$.

B Algebraic structures

Algebras and groups. We work with real Lie algebras \mathfrak{g} , and corresponding group G , of dimension $\dim G = D$ with a basis of anti-Hermitian generators $\{t_a\}$ equipped with an ad-invariant symmetric pairing given by the Cartan-Killing form, $\kappa = \langle \cdot, \cdot \rangle$, obeying

$$[t_a, t_b] = f_{ab}{}^c t_c, \quad \kappa_{ab} = \langle t_a, t_b \rangle = -\frac{1}{2h^\vee} f_{ad}{}^e f_{be}{}^d. \quad (\text{B.1})$$

Left/right-invariant forms and adjoint actions for a group element $g(x) \in G$, depending on local coordinates x^i , are defined according to

$$\begin{aligned} dg g^{-1} &= v = v^a t_a = v^a dx^i t_a, & g^{-1} dg &= e = e^a t_a = e^a dx^i t_a, \\ \text{ad}_g t &= g t_a g^{-1} = D[g]_a{}^b t_b, & v^a &= e^b D[g]_b{}^a. \end{aligned} \quad (\text{B.2})$$

This definition of the adjoint action obeys

$$D[g]^{-1} = D[g^{-1}], \quad D[g]\kappa D[g]^T = \kappa, \quad (\text{B.3})$$

we will write $D \equiv D[g]$ when clear from the context.

The Maurer-Cartan equations are

$$dv^a = +\frac{1}{2}f_{bc}{}^a v^b \wedge v^c, \quad de^a = -\frac{1}{2}f_{bc}{}^a e^b \wedge e^c, \quad dD[g]_a{}^b = v^c D[g]_a{}^d f_{cd}{}^b. \quad (\text{B.4})$$

R-matrices. We consider \mathcal{R} a skew-symmetric endomorphism of \mathfrak{g} defined as

$$\mathcal{R}(t_a) = R_a{}^b t_b, \quad R_{ab} = R_a{}^c \kappa_{cb} = -R_{ba}, \quad R^{ab} = \kappa^{ac} R_c{}^b = -R^{ba}. \quad (\text{B.5})$$

From \mathcal{R} is constructed a second bracket over the vector space \mathfrak{g} ,

$$[x, y]_{\mathcal{R}} = [\mathcal{R}(x), y] + [x, \mathcal{R}(y)], \quad [t_a, t_b]_{\mathcal{R}} = \tilde{f}_{ab}{}^c t_c, \quad \tilde{f}_{ab}{}^c = R_a{}^e f_{eb}{}^c + R_b{}^e f_{ae}{}^c. \quad (\text{B.6})$$

This will obey the Jacobi identity provided \mathcal{R} solves the modified classical Yang-Baxter equation

$$[\mathcal{R}(X), \mathcal{R}(Y)] - \mathcal{R}([X, Y]_{\mathcal{R}}) + c^2[X, Y] = 0 \quad \forall X, Y, \in \mathfrak{g}. \quad (\text{B.7})$$

We will define the algebra constructed from this bracket as $\mathfrak{g}_{\mathcal{R}}$. We have two Lie-brackets giving algebras \mathfrak{g} and $\mathfrak{g}_{\mathcal{R}}$ over the same vector space and this set up is also called a bi-algebra. Technically the construction of eq. (B.6) means this is a coboundary bi-algebra. It can be useful to define a element $\mathbf{r} \in \wedge^2 \mathfrak{g}$ as

$$\mathbf{r} = R^{ab} t_a \otimes t_b. \quad (\text{B.8})$$

Drinfel'd double. Here we consider real Lie algebra \mathfrak{d} , and corresponding group \mathbb{D} , of dimension $\dim \mathbb{D} = 2d$ with a basis of anti-Hermitian generators $\{\mathbb{T}_A\}$ equipped with an ad-invariant symmetric pairing, $\eta = \langle\langle \cdot, \cdot \rangle\rangle$, obeying

$$[\mathbb{T}_A, \mathbb{T}_B] = \mathbb{F}_{AB}{}^C \mathbb{T}_C, \quad \eta_{AB} = \langle\langle \mathbb{T}_A, \mathbb{T}_B \rangle\rangle. \quad (\text{B.9})$$

A classical double is such a real Lie algebra that admits a decomposition $\mathfrak{d} = \mathfrak{g} \oplus \tilde{\mathfrak{g}}$ as the sum of two Lie subalgebras each of dimension d that are Lagrangian (maximally isotropic with respect to η). In a basis $\mathbb{T}_A = (\tilde{T}^a, T_a)$ we have that

$$\begin{aligned} [T_a, T_b] &= F_{ab}{}^c T_c, & [\tilde{T}^a, \tilde{T}^b] &= \tilde{F}^{ab}{}^c \tilde{T}^c, & [T_a, \tilde{T}^b] &= \tilde{F}^{bc}{}_a T_c - F_{ac}{}^b \tilde{T}^c, \\ \langle\langle T_a, T_b \rangle\rangle &= \langle\langle \tilde{T}^a, \tilde{T}^b \rangle\rangle = 0, & \langle\langle T_a, \tilde{T}^b \rangle\rangle &= \delta_a{}^b, & \langle\langle \tilde{T}^a, T_b \rangle\rangle &= \delta^a{}_b. \end{aligned} \quad (\text{B.10})$$

The Jacobi identity of $\mathbb{F}_{AB}{}^C$ places a compatibility condition on the two Lie subalgebras, namely that $\delta(T_a) = \tilde{F}^{bc}{}_a T_b \otimes T_c$ viewed as a map $\mathfrak{g} \rightarrow \wedge^2 \mathfrak{g}$, should be a one-cocycle for \mathfrak{g} valued in $\wedge^2 \mathfrak{g}$ obeying

$$0 = d\delta(X, Y) \equiv \text{ad}_X \delta(Y) - \text{ad}_Y \delta(X) - \delta([X, Y]), \quad (\text{B.11})$$

in which the adjoint action extends to the tensor product as $\text{ad}_X Y = (1 \otimes \text{ad}_X + \text{ad}_X \otimes 1)Y$ for $Y \in \mathfrak{g} \otimes \mathfrak{g}$. Of particular interest will be the case when the one-cocycle is a one-coboundary

$$\delta(X) = [X, \mathbf{r}], \quad \mathbf{r} \in \mathfrak{g} \wedge \mathfrak{g}. \quad (\text{B.12})$$

Identifying $T_a = t_a$ and with $\mathbf{r} = R^{ab} t_a \otimes t_b$ we have that

$$\tilde{F}^{ab}{}_c = R^{ae} F_{ce}{}^b - R^{be} F_{ce}{}^a, \quad (\text{B.13})$$

which is nothing other than the raising of indices on $\tilde{F}_{ab}{}^c$ defined in eq. (B.6) using κ . In this case the double $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{g}_R$ and the Jacobi identity of $\mathbb{F}_{AB}{}^C$ follows from the mCYBE.

In this way we have an equivalence between such doubles and coboundary Lie-bialgebras. The cocycle δ can be integrated to give a cocycle on G valued in $\mathfrak{g} \wedge \mathfrak{g}$

$$\Pi[g] = (1 - \text{ad}_{g^{-1}} \otimes \text{ad}_{g^{-1}}) \mathbf{r}, \tag{B.14}$$

which for $g = \exp(\epsilon x)$ has $\Pi[g] \sim \epsilon[X, \mathbf{r}] = \epsilon\delta(X)$ and obeys

$$\Pi[hg] = \Pi(g) + \text{ad}_{g^{-1}} \otimes \text{ad}_{g^{-1}} \Pi[h], \quad \Pi[e] = 1. \tag{B.15}$$

In components

$$\Pi[g]^{ab} = R^{ab} - D[g^{-1}]_c^a R^{cd} D[g^{-1}]_d^b. \tag{B.16}$$

Whilst the above expression applies in the case of the coboundary specialisation, one can construct the same group cocycle for any double as follows. Let g be a group element for $G = \exp \mathfrak{g} \subset \mathbb{D}$ and using its adjoint action on \mathfrak{d} ,

$$g \mathbb{T}_A g^{-1} = D[g]_A^B \mathbb{T}_B, \quad D[g]_A^B = \begin{pmatrix} D[g]_a^b & D[g]^{ab} \\ 0 & D[g]_a^b \end{pmatrix}, \tag{B.17}$$

one has

$$\Pi[g]^{ab} = D[g]^{ac} D[g]_c^b. \tag{B.18}$$

The group composition of the adjoint action in \mathbb{D} shows that the cocycle properties eq. (B.15) holds and its derivative returns the algebra cocycle δ . The cocycle can be understood as being as element of $T_e(G) \otimes T_e(G)$, and by taking its right translation to a point g we have a bi-vector $\Pi_g \in T_g(G) \otimes T_g(G)$; this endows a Poisson structure to G making it a Lie-Poisson group manifold.

Manin pair, triple and quasi-triple. We now describe a weakening of the above structure to define a Manin quasi-triple. A pair $(\mathfrak{d}, \tilde{\mathfrak{h}})$ consisting of an algebra, \mathfrak{d} and a Lagrangian subalgebra $\tilde{\mathfrak{h}} \subset \mathfrak{d}$ is called a Manin pair. A Manin quasi-triple $(\mathfrak{d}, \tilde{\mathfrak{h}}, \mathfrak{k})$ is a Manin pair $(\mathfrak{d}, \tilde{\mathfrak{h}})$ together with a choice of complementary Lagrangian subspace \mathfrak{k} such that $\mathfrak{d} = \tilde{\mathfrak{h}} \oplus \mathfrak{k}$. Different choices of complementary subspaces are related by a twist $t \in \Lambda^2 \tilde{\mathfrak{h}}$ [66]. The salient difference to Drinfel'd double is that the complementary Lagrangian \mathfrak{k} need not be a subalgebra.

We define a basis of anti-Hermitian generators $\{\mathbb{T}_A\}$ for \mathfrak{d} equipped with an ad-invariant symmetric pairing, $\eta = \langle \cdot, \cdot \rangle$, obeying

$$[\mathbb{T}_A, \mathbb{T}_B] = \mathbb{F}_{AB}^C \mathbb{T}_C, \quad \eta_{AB} = \langle \mathbb{T}_A, \mathbb{T}_B \rangle. \tag{B.19}$$

Letting $\mathbb{T}_A = (\tilde{T}^a, T_a)$ be the decomposition $\mathfrak{d} = \tilde{\mathfrak{h}} \oplus \mathfrak{k}$ these relations read

$$\begin{aligned} [T_a, T_b] &= F_{ab}^c T_c + \phi_{abc} \tilde{T}^c, & [\tilde{T}^a, \tilde{T}^b] &= \tilde{F}^{ab}_c \tilde{T}^c, & [T_a, \tilde{T}^b] &= \tilde{F}^{bc}_a T_c - F_{ac}^b \tilde{T}^c, \\ \langle T_a, T_b \rangle &= \langle \tilde{T}^a, \tilde{T}^b \rangle = 0, & \langle T_a, \tilde{T}^b \rangle &= \delta_a^b. \end{aligned} \tag{B.20}$$

The object ϕ_{abc} is antisymmetric in all its indices and invariant under the (co-adjoint) action of $\tilde{H} = \exp \tilde{\mathfrak{h}}$.

Algebraic structure of λ - and η models. The integrable λ and η models can be placed into this algebraic framework [29, 90] in which $\mathfrak{d} = \mathfrak{g} + \mathfrak{g}$ (a Manin quasi-triple) and $\mathfrak{d} = \mathfrak{g} + \mathfrak{g}_{\mathcal{R}} = \mathfrak{g}^{\mathbb{C}}$ (a Drinfel'd double) respectively.

Consider a Lie algebra \mathfrak{g} endowed with an ad-invariant non-degenerate symmetric bilinear form κ . The construction of the λ -deformation requires a double that is the direct sum $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{g}$ equipped with the inner product

$$\langle\langle \{X_1, Y_1\}, \{X_2, Y_2\} \rangle\rangle = \langle X_1, X_2 \rangle - \langle Y_1, Y_2 \rangle.$$

The two subspaces completing the Manin quasi-triple are taken to be the diagonal subalgebra $\tilde{\mathfrak{h}} = \mathfrak{g}_{\text{diag}}$, embedded in \mathfrak{d} by the map $X \mapsto \{X, X\}/\sqrt{2}$ for $X \in \mathfrak{g}$, and the anti-diagonal subspace $\mathfrak{k} = \mathfrak{g}_{\text{anti-diag}}$ embedded as $X \mapsto \{X, -X\}/\sqrt{2}$ in \mathfrak{d} . Let $f_{ab}{}^c$ be generators of \mathfrak{g} , then in \mathfrak{d} we have that

$$F_{ab}{}^c = 0, \quad \tilde{F}^{ab}{}_c = \frac{1}{\sqrt{2}} f^{ab}{}_c = \frac{1}{\sqrt{2}} \kappa^{ad} \kappa^{be} f_{de}{}^f \kappa_{fc}, \quad \phi_{abc} = \frac{1}{\sqrt{2}} f_{abc} = \frac{1}{\sqrt{2}} f_{ab}{}^f \kappa_{fc}. \quad (\text{B.21})$$

The η -deformation on the other hand, the double is determined by the operator \mathcal{R} entering the definition of the deformation. This operator is the canonical R-matrix associated to a semi-simple Lie algebra \mathfrak{g} with Killing form κ . It acts by anti-symmetrically swapping positive and negative roots and annihilates the Cartan. As described above, since \mathcal{R} is a solution to the classical (modified) Yang-Baxter equation it defines a second Lie-bracket $[\cdot, \cdot]_{\mathcal{R}}$ on \mathfrak{g} . The double is the direct sum $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{g}_{\mathcal{R}}$, which is isomorphic to the complexification $\mathfrak{g}^{\mathbb{C}}$ of \mathfrak{g} . This double can be decomposed into a Manin pair using the Iwasawa decomposition $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g} \oplus (\mathfrak{a} + \mathfrak{n})$, where \mathfrak{g} and $\mathfrak{a} + \mathfrak{n}$ are both Lagrangian subalgebras of $\mathfrak{d} = \mathfrak{g}^{\mathbb{C}}$. The ad-invariant non-degenerate symmetric bilinear form on $\mathfrak{d} = \mathfrak{g}^{\mathbb{C}}$ is

$$\langle\langle Z_1, Z_2 \rangle\rangle = -i \kappa(Z_1, Z_2) + i \overline{\kappa(Z_1, Z_2)},$$

where $Z \in \mathfrak{g}^{\mathbb{C}}$ and $\bar{\cdot}$ denotes the complex conjugation.

C Solution to the section condition

In this appendix we show how the choice of generalised frame fields solves the section condition (3.28). Even more, this equation together with the requirement to be an $O(d, d)$ -element, will completely fix the form of the generalised frame field in (3.29) completely in terms of the element of the right-invariant form on the double $\mathfrak{d} = \tilde{\mathfrak{h}} \oplus \mathfrak{k}$,

$$\mathbb{T}_A \mathbb{W}^A{}_I dX^I = \mathbb{T}_A \mathbb{M}_B{}^A \mathbb{E}^B{}_I dX^I = \partial_I \mathbb{g} \mathbb{g}^{-1} dX^I. \quad (\text{C.1})$$

Let us now explicitly calculate $\mathbb{W}^A{}_I$ by using the parameterization of the double element $\mathbb{g} = \tilde{h}(\tilde{x}_{\tilde{i}}) m(x^i)$, $\tilde{h} \in \tilde{H}$ and $m \in \exp(\mathfrak{k})$. We obtain

$$d\mathbb{g} \mathbb{g}^{-1} = \mathbb{T}_A \mathbb{W}^A{}_I dX^I = \tilde{h} \partial_i m m^{-1} \tilde{h}^{-1} dx^i + \partial_{\tilde{i}} \tilde{h} \tilde{h}^{-1} d\tilde{x}_{\tilde{i}} = \mathbb{T}_A \mathbb{W}^A{}_i dx^i + T^a \mathbb{V}_a{}^{\tilde{i}} d\tilde{x}_{\tilde{i}} \quad (\text{C.2})$$

and $\mathbb{W}^a{}_{\tilde{i}} = 0$ because h is an element of the subgroup \tilde{H} . As usual the inverse transpose of $\mathbb{W}^A{}_I$ is denoted as $\mathbb{W}_A{}^I$, for which we have that $\mathbb{W}^{ai} = 0$.

Looking now at the generalised frame field, it is convenient to decompose it into two parts,

$$\widehat{E}_A^{\hat{I}} = M_A^B \widehat{V}_B^{\hat{I}}. \quad (\text{C.3})$$

Using this decomposition we need to check that

$$\widehat{E}^A_{\hat{I}} \mathbb{E}_A^I \partial_I = \widehat{V}^A_{\hat{I}} \mathbb{W}_A^I \partial_I = (0 \ \partial_i). \quad (\text{C.4})$$

We need the above equation to only hold on the physical fields that we are considering, which will depend only on the coordinates x^i so that we may use on the left hand side that $\partial_I = (0 \ \partial_i)$. To see the first equality of eq. (C.4) the parametrisation $\mathbf{g}(x, \tilde{x}) = \tilde{h}(\tilde{x}_{\tilde{i}})m(x^i)$ is paramount since it ensures that the differences between the adjoint action \mathbb{M} of \mathbf{g} and the adjoint action M of m don't contribute. Then eq. (C.4) reduces to

$$\left(\widehat{V}^{Ai} \mathbb{W}_A^j \partial_j \ \widehat{V}^A_{\hat{I}} \mathbb{W}_A^j \partial_j \right) = (0 \ \partial_i) \quad (\text{C.5})$$

or equivalently

$$\widehat{V}^{Ai} \mathbb{W}_A^j = 0 \quad \text{and} \quad \widehat{V}^A_{\hat{I}} \mathbb{W}_A^j = \delta_i^j. \quad (\text{C.6})$$

The second of these is satisfied providing

$$\mathbb{T}_A \widehat{V}^A_{\hat{I}} = T_a \mathbb{W}^a_{\hat{I}} + \tilde{T}^a \mathbb{W}_a^j \rho_{ji}, \quad (\text{C.7})$$

with arbitrary and to be fixed matrix ρ_{ij} . For the first component \widehat{V}^{Ai} we find

$$\mathbb{T}_A \widehat{V}^{Ai} \partial_i = \tilde{T}^a \mathbb{W}_a^i \partial_i, \quad (\text{C.8})$$

because $\mathbb{W}^{ai} = 0$. Furthermore we need to require the generalised frame field to be an $O(D, D)$ element, that is $\widehat{V}^A_{\hat{I}}$ has to have the property

$$\widehat{V}^A_{\hat{I}} \eta_{AB} \widehat{V}^B_{\hat{J}} = \eta_{\hat{I}\hat{J}}. \quad (\text{C.9})$$

This implies several constraints (those on the right being implied by those on the left):

$$\widehat{V}^A_{\hat{I}} \eta_{AB} \widehat{V}^B_{\hat{J}} = 0 \quad \langle \partial_i \mathbf{g} \mathbf{g}^{-1}, \partial_j \mathbf{g} \mathbf{g}^{-1} \rangle + 2\rho_{(ij)} = 0 \quad (\text{C.10})$$

$$\widehat{V}^{Ai} \eta_{AB} \widehat{V}^{Bj} = 0 \quad \langle \tilde{T}^a, \tilde{T}^b \rangle v_a^i v_b^j = 0 \quad (\text{C.11})$$

$$\widehat{V}^{Ai} \eta_{AB} \widehat{V}^B_{\hat{J}} = \delta_j^i \quad \langle \tilde{T}^a, T_b \rangle v_a^i v_b^j = \delta_j^i. \quad (\text{C.12})$$

The first term in the first equation on the right vanishes by assumption and the second one implies that ρ_{ij} has to be antisymmetric. All other identities follow automatically.

Summarising the discussion above, $\widehat{V}_A^{\hat{I}}$ reads

$$\widehat{V}_A^{\hat{I}} = \begin{pmatrix} v_a^i & 0 \\ v_a^j \rho_{ji} & v_a^i \end{pmatrix}, \quad (\text{C.13})$$

as claimed in (3.29). The precise form of ρ_{ij} will be fixed to ensure the frame fields obey a frame algebra under the generalised Lie derivative.

D Fluxes in the generalised parallelizable frame

It is instructive to compute the components of $F_{\hat{J}\hat{J}\hat{K}}$, the structure constants dressed by the generalised frame fields as in eq. (3.43), explicitly. In conventional notation these are denoted $\mathbf{H}, \mathbf{Q}, \mathbf{F}, \mathbf{R}$ [91]. First let us consider the case of a Drinfel'd double for which we have in general

$$\begin{aligned}\mathbf{H}_{ijk} &= 0, \\ \mathbf{F}_{ij}{}^k &= e^b{}_{[i}e^c{}_{j]}e_a{}^k F_{bc}{}^a \\ \mathbf{Q}^{ij}{}_k &= e_a{}^{[i}e_b{}^{j]}e^c{}_k(\tilde{F}^{ab}{}_c + F_{dc}{}^b\Pi^{ad}) = -e_a{}^{[i}e_b{}^{j]}\partial_k\Pi^{ab}, \\ \mathbf{R}^{ijk} &= \frac{1}{2}e_a{}^{[i}e_b{}^j e_c{}^{k]}(F_{de}{}^a\Pi^{bd}\Pi^{ce} + \tilde{F}^{ab}{}_d\Pi^{dc}), \\ &= -\frac{1}{4}e_a{}^{[i}e_b{}^j e_c{}^{k]}(\tilde{F}^{a[b}{}_d\Pi^{c]d} - 2F_{de}{}^a\Pi^{db}\Pi^{ec}) = 0.\end{aligned}$$

The identity required to show the vanishing of the R-flux is slightly involved and was provided in [6].

In the case that the Drinfel'd Double corresponds to a coboundary Lie bialgebra, i.e. both $\tilde{F}^{ab}{}_c$ and Π^{ab} are expressible in terms of an R -matrix (see appendix A) we can go a little further to express the fluxes as

$$\begin{aligned}\mathbf{H}_{ijk} &= 0, \\ \mathbf{F}_{ij}{}^k &= v^b{}_{[i}v^c{}_{j]}v_a{}^k f_{bc}{}^a, \\ \mathbf{Q}^{ij}{}_k &= 2v_a{}^{[i}v_b{}^{j]}v^c{}_k F_{dc}{}^a R^{bd} = v_a{}^{[i}v_b{}^{j]}v^c{}_k \tilde{F}^{ab}{}_c, \\ \mathbf{R}^{ijk} &= 3\left(v_a{}^{[i}v_b{}^j v_c{}^{k]} - e_a{}^{[i}e_b{}^j e_c{}^{k]}\right) F_{de}{}^a R^{bd} R^{ce} \\ &= -3c^2\left(v_a{}^{[i}v_b{}^j v_c{}^{k]} - e_a{}^{[i}e_b{}^j e_c{}^{k]}\right) \tilde{F}^{abc} = 0.\end{aligned}$$

In the two last fluxes, we used the definition of the modified Yang-Baxter equation.

Let us now turn to the more general case of a Manin quasi-triple. Here we must make a slight refinement, the H -flux also has a contribution that arises as a twisting of the Courant bracket as discussed in comment 2. In what follows we shall strip off this twisting using the frame fields

$$\hat{E}'_A{}^{\hat{I}} = M_A{}^B \begin{pmatrix} v^b{}_i & 0 \\ 0 & v_b{}^i \end{pmatrix}_B{}^{\hat{I}}, \quad (\text{D.1})$$

that obey

$$\hat{\mathcal{L}}_{\hat{E}'_A} \hat{E}'_B{}^{\hat{I}} = F_{AB}{}^C \hat{E}'_C{}^{\hat{I}} + \begin{pmatrix} \hat{E}'_A{}^j \hat{E}'_B{}^k (\Omega^{(3)} - d\omega^{(2)})_{jki} \\ 0 \end{pmatrix}. \quad (\text{D.2})$$

We then consider $F'_{\hat{I}\hat{J}\hat{K}} = F_{ABC} \hat{E}'^C{}_I \hat{E}'^B{}_J \hat{E}'^A{}_K$ and simply add back the contribution to the \mathbf{H} given by $\bar{\mathbf{H}} = \Omega^{(3)} - d\omega^{(2)}$.

We recall that from the coset representative $m(x)$ for \mathcal{D}/\tilde{H} we have

$$dm m^{-1} = v^a{}_i dx^i T_a + A_{ai} dx^i \tilde{T}^a.$$

Then evaluating the fluxes one finds

$$\begin{aligned}
 \mathbf{H}_{ijk} &= \left(\phi_{abc} M^a{}_d M^b{}_e M^c{}_f + \frac{1}{2} F_{ab}{}^c M^a{}_d M^b{}_e M_{cf} - \frac{1}{2} \tilde{F}^{ab}{}_c M_{ad} M_{be} M^c{}_f \right) v^d [{}^i v^e{}_j v^f{}_k] + \bar{\mathbf{H}}_{ijk}, \\
 \mathbf{F}_{ij}{}^k &= 2v_a{}^k v^e [{}^i v^f{}_j] \left(F_{bc}{}^d (M_{de} M^b{}_f M^{ca} - M_d{}^a M^b{}_e M^c{}_f) - \tilde{F}^{bc}{}_d (M_{be} M_c{}^a M^d{}_f - M_{be} M_{cf} M^{da}) \right. \\
 &\quad \left. + M^c{}_e M^d{}_f M^{ba} \phi_{bcd} \right), \\
 \mathbf{Q}^{ij}{}_k &= 2v_a [{}^i v_b{}^j] v^f{}_k \left(-F_{cd}{}^e (M_{ef} M^{ca} M^{db} - M_e{}^b M^c{}_f M^{da}) \right. \\
 &\quad \left. + \tilde{F}^{cd}{}_e (M_c{}^a M_d{}^b M^e{}_f + M_{cf} M_d{}^b M^{ea}) + M^d{}_f M^{ea} M^{cb} \phi_{dec} \right), \\
 \mathbf{R}^{ijk} &= \left(\phi_{abc} M^{da} M^{eb} M^{cf} + \frac{1}{2} F_{ab}{}^c M^{da} M^{eb} M_c{}^f + \frac{1}{2} \tilde{F}^{ab}{}_c M_a{}^d M_b{}^e M^{cf} \right) v_d [{}^i v_e{}^j v^k{}_f].
 \end{aligned}$$

Here

$$\bar{\mathbf{H}}_{ijk} = -3A_{a[i} A_{bj} v^c{}_k] \tilde{F}^{ab}{}_c + \phi_{abc} v^a [{}^i v^b{}_j v^c{}_k] \quad (\text{D.3})$$

is the contribution to the H-flux coming from the twisting of the Courant bracket.

Specialising to $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{g}$ relevant to the λ -model we can now go further by using the explicit form of the adjoint action $M_A{}^B$, given in eq. (5.56). Doing so we find numerous cancellations to leave

$$\begin{aligned}
 \mathbf{H}_{ijk} &= -\frac{3}{\sqrt{2}} A_{a[i} A_{bj} v^c{}_k] f^{ab}{}_c + \frac{2}{\sqrt{2}} f_{abc} v^a [{}^i v^b{}_j v^c{}_k], \\
 \mathbf{F}_{ij}{}^k &= 0, \quad \mathbf{Q}^{ij}{}_k = \frac{1}{\sqrt{2}} v_a [{}^i v_b{}^j] v^c{}_k f^{ab}{}_c, \quad \mathbf{R}^{ijk} = 0,
 \end{aligned}$$

in which κ is used to raise algebra indices out of position. It might seem contrary to have \mathbf{Q} rather than \mathbf{F} flux but it reflects the construction of the geometry as a coset $\mathbb{D}/G_{\text{diag}}$ and that $G_{\text{anti-diag}}$ is not a subgroup of \mathbb{D} .

E Dilaton in PL and λ models

Here we show that the constraint on the doubled dilaton d matches the (conventional) dilatons for both PL and λ -models.

We begin by extracting the metric for the PL model of eq. (2.9)

$$\begin{aligned}
 G &= e^T (1 + E_0^- \Pi)^{-1} G_0 (1 - \Pi E_0^+)^{-1} e \\
 &= e^T \left(1 + \tilde{g}_0^{-1} (\Pi - \tilde{B}_0) \right)^{-1} \tilde{g}_0^{-1} \left(1 - (\Pi - \tilde{B}_0) \tilde{g}_0^{-1} \right)^{-1} e,
 \end{aligned} \quad (\text{E.1})$$

in which $E_0^\pm = G_0 \pm B_0$ and $E_0^- = \left(\tilde{g}_0 - \tilde{b}_0 \right)^{-1}$ and e are the components of the left-invariant forms. It is simple to take the determinant

$$\log \det G = 2 \log \det e - \log \det \tilde{g}_0 - 2 \log \det \left(1 + \tilde{g}_0^{-1} (\Pi - \tilde{B}_0) \right) \quad (\text{E.2})$$

Since $\det e = \det v$ we conclude from (4.7)

$$\phi = \phi_0 - \frac{1}{4} \log \det \tilde{g}_0 - \frac{1}{2} \log \det \left(1 + \tilde{g}_0^{-1} (\Pi - \tilde{B}_0) \right), \quad (\text{E.3})$$

and choosing $\phi_0 = \frac{1}{4} \log \det \tilde{g}_0$, which is of course a constant, gives the result provided in the more mathematically inclined treatment of [73].

For the case of the λ -model, the expression correctly reduces to the known expression, see e.g. [92]. This can be readily seen by conveniently expressing the curved metric g_{ij} in terms of right-invariant form on G and the flat metric

$$\begin{aligned} g_{ab} &= ((1 - \lambda D^{-1})^{-1} \kappa + (1 - \lambda D)^{-1} \kappa - \kappa)_{ab} \\ &= (1 - \lambda^2) ((1 - \lambda D^{-1})^{-1} \kappa (1 - \lambda D)^{-1} \kappa)_{ab}. \end{aligned} \quad (\text{E.4})$$

Starting from the expression (4.7), we indeed obtained the correct expression for the dilaton of the λ -model:

$$\phi = \phi_0 + \frac{1}{4} \log |\det g_{ij}| - \frac{1}{2} \log |\det v^a{}_i| = \phi'_0 + \frac{1}{4} \log |\det g_{ab}| = \phi''_0 - \frac{1}{2} \log |\det(1 - \lambda D^{-1})|,$$

where we have used that the adjoint action has unit determinant. The last step is obtained by plugging (E.4) and using that $D^{-T} = \kappa^{-1} D \kappa$. All constant contribution were successively absorbed into the constant dilaton term.

F Details of η -supergravity solution

In this appendix we detail the full modified supergravity solution outlined in section 5.2. For the generators t_a of $\mathfrak{g} = \mathfrak{su}(1, 1) \oplus \mathfrak{su}(2)$ we let t_i be those of $\mathfrak{su}(1, 1)$ and $t_{\bar{i}}$ be those of $\mathfrak{su}(2)$ in a basis where the non-vanishing structure constants are given by

$$f_{12}{}^3 = f_{13}{}^2 = f_{32}{}^1 = -1, \quad f_{\bar{1}\bar{2}}{}^{\bar{3}} = f_{\bar{2}\bar{3}}{}^{\bar{1}} = f_{\bar{3}\bar{1}}{}^{\bar{2}} = -1. \quad (\text{F.1})$$

To raise and lower indices we use the ad-invariant inner-product given by

$$\kappa_{ij} = \frac{\alpha}{2} f_{ik}{}^l f_{jl}{}^k = \text{diag}(\alpha, \alpha, -\alpha), \quad \kappa_{\bar{i}\bar{j}} = -\frac{\alpha}{2} f_{\bar{i}\bar{k}}{}^{\bar{l}} f_{\bar{j}\bar{l}}{}^{\bar{k}} = \text{diag}(\alpha, \alpha, \alpha), \quad (\text{F.2})$$

in which we note that the overall normalisation of the $\mathfrak{su}(2)$ part is of opposite sign to that of the $\mathfrak{su}(1, 1)$. The solution of the $c^2 = -1$ mCYBE is given by an R-matrix with non-vanishing components

$$R_{12} = -R_{21} = -\alpha, \quad R_{\bar{1}\bar{2}} = -R_{\bar{2}\bar{1}} = \alpha. \quad (\text{F.3})$$

We supplement the six dimensional space corresponding to the deformed $\text{AdS}_3 \times S^3$ with a four-torus (with coordinates x^μ , $\mu = 1 \dots 4$) such that the NS data is

$$\begin{aligned} ds^2 &= v^a g_{ab} v^b + dx^\mu dx^\mu = v^a \left(\eta \kappa_{ab} + \frac{\eta^3}{1 + \eta^2} R_a{}^d R_{db} \right) v^b + dx^\mu dx^\mu, \\ B &= -\frac{\eta^2}{2(1 + \eta^2)} \left(R_{ab} v^a \wedge v^b \right) = \frac{\eta^2 \alpha}{(1 + \eta^2)} \left(v^1 \wedge v^2 - v^{\bar{1}} \wedge v^{\bar{2}} \right), \\ \Phi &= \log \left(\frac{\eta^{3/2}}{1 + \eta^2} \right) + \phi_0 \quad H = dB = 0. \end{aligned} \quad (\text{F.4})$$

Note that we have chosen to work with the right-invariant Maurer-Cartan forms rather than the left; this removes all coordinate dependence from the metric and fluxes. The curvatures that follow from this metric have non-vanishing components

$$\text{Ric}_{ij} = \frac{(1 + \eta^2)}{\alpha} \kappa_{ij} - \frac{(1 + \eta^2)(3 + \eta^2)}{2\eta\alpha} g_{ij}, \quad \text{Ric}_{\bar{i}\bar{j}} = -\frac{(1 + \eta^2)}{\alpha} \kappa_{\bar{i}\bar{j}} + \frac{(1 + \eta^2)(3 + \eta^2)}{2\eta\alpha} g_{\bar{i}\bar{j}}, \quad (\text{F.5})$$

and are such that the curvature scalar is zero,

$$R = -\frac{1}{2} f_{ac}{}^d f_{bd}{}^c g^{ab} - \frac{1}{4} f_{ac}{}^e f_{bd}{}^f g^{ab} g^{cd} g_{ef} = 0. \quad (\text{F.6})$$

This is fundamentally due to the choice of opposing normalisations for the $\mathfrak{su}(2)$ and $\mathfrak{su}(1, 1)$ in the inner-product.

The modified supergravity is defined by the one-form

$$I = \frac{1}{2} R^{ab} f_{ab}{}^c g_{cd} v^d = -\eta (v^{\bar{3}} + v^3), \quad (\text{F.7})$$

and related

$$Z = d\Phi + \iota_I B = 0, \quad X = I + Z = I. \quad (\text{F.8})$$

Eq. (5.35) encodes the unique solution for the R/R fluxes however this is in six-dimensions. Here we need to uplift it to 10-dimensions. In six-dimensions we define

$$\widehat{G}_{6d}^{(1)} = -\frac{1 + \eta^2}{\sqrt{2}} R^{ab} f_{abc} v^c, \quad \widehat{G}_{6d}^{(3)} = \frac{1 + \eta^2}{3\sqrt{2}} f_{abc} v^a \wedge v^b \wedge v^c,$$

and the six-dimensional poly-form

$$\widehat{\mathcal{F}}_{6d} = e^\Phi e^{-B} \left(\widehat{G}_{6d}^{(1)} + \widehat{G}_{6d}^{(3)} \right), \quad (\text{F.9})$$

which by construction has vanishing Lie derivative along \mathcal{I} i.e. $L_{\mathcal{I}} \widehat{\mathcal{F}}_{6d} = 0$. The components of this obey

$$\iota_{\mathcal{I}} \widehat{\mathcal{F}}_{6d}^{(1)} = 0, \quad d\widehat{\mathcal{F}}_{6d}^{(1)} = \iota_{\mathcal{I}} \widehat{\mathcal{F}}_{6d}^{(3)}, \quad d\widehat{\mathcal{F}}_{6d}^{(3)} = \iota_{\mathcal{I}} \widehat{\mathcal{F}}_{6d}^{(5)}, \quad \star_6 \widehat{\mathcal{F}}_{6d}^{(1)} = -\widehat{\mathcal{F}}_{6d}^{(5)}, \quad \star_6 \widehat{\mathcal{F}}_{6d}^{(3)} = \widehat{\mathcal{F}}_{6d}^{(3)}. \quad (\text{F.10})$$

From this we can build a ten-dimensional R/R poly-form

$$\mathcal{F}_{10d} = \mu \widehat{\mathcal{F}}_{6d} \wedge (1 + \omega - \text{vol}_4), \quad (d + H \wedge -Z \wedge -\iota_I) \mathcal{F}_{10d} = 0 \quad (\text{F.11})$$

in which μ is a normalisation to be fixed, and $\omega = \vec{n} \cdot \vec{\omega}$ is expanded in the basis of self-dual three forms on T^4 and $\text{vol}_4 = dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^4$. With this in mind it is quite easy now to verify the modified supergravity equations of eq. (A.4) are satisfied providing the normalisation of the R/R sector are set such that

$$\mu = e^{-\phi_0} (2\eta(1 + \vec{n}^2))^{-\frac{1}{2}}. \quad (\text{F.12})$$

Explicitly we have

$$\begin{aligned}
 \mathcal{F}_{10d}^{(1)} &= \rho \left(v^3 + v^{\bar{3}} \right), \quad \mathcal{F}_{10d}^{(3)} = \rho \left(v^3 + v^{\bar{3}} \right) \wedge \omega + \frac{\alpha \rho}{(1 + \eta^2)} \left(v^{123} - v^{\bar{1}\bar{2}\bar{3}} - \eta^2 v^{12\bar{3}} + \eta^2 v^{\bar{1}\bar{2}3} \right) \\
 \mathcal{F}_{10d}^{(5)} &= -\rho \left(v^3 + v^{\bar{3}} \right) \wedge \text{vol}_4 + \frac{\alpha \rho}{(1 + \eta^2)} \left(v^{123} - v^{\bar{1}\bar{2}\bar{3}} - \eta^2 v^{12\bar{3}} + \eta^2 v^{\bar{1}\bar{2}3} \right) \wedge \omega \\
 &\quad + \frac{\alpha^2 \eta^2 \rho}{(1 + \eta^2)^2} \left(v^{123\bar{1}\bar{2}} + v^{12\bar{1}\bar{2}\bar{3}} \right),
 \end{aligned} \tag{F.13}$$

in which we let $\rho = \eta(1 + \bar{\eta}^2)^{-\frac{1}{2}}$.

G Drinfel'd doubles and group parameterisations

G.1 $\mathfrak{su}(2) \oplus \mathfrak{e}_3$

We work with the following basis of generators

$$\begin{aligned}
 T_1 &= \frac{i}{2} \mathbb{I} \otimes \sigma_1 & T_2 &= \frac{i}{2} \mathbb{I} \otimes \sigma_2 & T_3 &= \frac{i}{2} \mathbb{I} \otimes \sigma_3 \\
 \tilde{T}^1 &= -\frac{1}{2} \sigma_3 \otimes \sigma_1 - \frac{i}{2} \mathbb{I} \otimes \sigma_2 & \tilde{T}^2 &= -\frac{1}{2} \sigma_3 \otimes \sigma_2 + \frac{i}{2} \mathbb{I} \otimes \sigma_1 & \tilde{T}^3 &= -\frac{1}{2} \sigma_3 \otimes \sigma_3.
 \end{aligned} \tag{G.1}$$

Defining projectors

$$P^\pm = \frac{1}{2} (\mathbb{I} \pm \sigma_3) \otimes \mathbb{I}, \tag{G.2}$$

allows us to realise the inner-product as

$$\langle\langle \mathbb{T}_A, \mathbb{T}_B \rangle\rangle = i \text{Tr} \left(P^+ \mathbb{T}_A P^+ \mathbb{T}_B - P^- \mathbb{T}_A P^- \mathbb{T}_B \right) = \eta_{AB}. \tag{G.3}$$

The \mathcal{R} -matrix that gives this a bialgebra structure is

$$\mathcal{R} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{G.4}$$

and we parametrise an $SU(2)$ element as

$$G = \begin{pmatrix} g & 0 \\ 0 & g \end{pmatrix}, \quad g = \begin{pmatrix} e^{-ix} \sqrt{1-r^2} & -e^{-i\phi} r \\ e^{i\phi} r & e^{ix} \sqrt{1-r^2} \end{pmatrix}, \tag{G.5}$$

such that the left-invariant forms defined by $G^{-1}dG = e^a T_a$ read

$$\frac{1}{2} (e^1 \pm ie^2) = \frac{e^{\mp i(x-\phi)}}{\sqrt{1-r^2}} (r(r^2-1)d(x+\phi) \pm idr), \quad e^3 = (r^2-1)dx + r^2d\phi. \tag{G.6}$$

The metric on S^3 is obtained as

$$\frac{-1}{4} \text{Tr} (G^{-1}dGG^{-1}dG) = (1-r^2)dx^2 + \frac{dr^2}{1-r^2} + r^2d\phi^2, \tag{G.7}$$

which is rendered more familiar with $r = \sin \theta$. Finally, we need the combination of adjoint actions that enter into the PL sigma models:

$$\Pi = \begin{pmatrix} 0 & 2r^2 & -2r\sqrt{1-r^2} \sin(x-\phi) \\ -2r^2 & 0 & -2r\sqrt{1-r^2} \cos(x-\phi) \\ 2r\sqrt{1-r^2} \sin(x-\phi) & 2r\sqrt{1-r^2} \cos(x-\phi) & 0 \end{pmatrix}. \tag{G.8}$$

G.2 $\mathfrak{su}(1, 1) \oplus \mathfrak{e}_3$

We work with the following basis of generators

$$\begin{aligned} T_1 &= \frac{1}{2}\sigma_3 \otimes \sigma_2 & T_2 &= \frac{1}{2}\sigma_3 \otimes \sigma_1 & T_3 &= \frac{i}{2}\mathbb{I} \otimes \sigma_3 \\ \tilde{T}^1 &= -\frac{1}{2}\sigma_3 \otimes \sigma_1 - \frac{i}{2}\mathbb{I} \otimes \sigma_2 & \tilde{T}^2 &= \frac{1}{2}\sigma_3 \otimes \sigma_2 - \frac{i}{2}\mathbb{I} \otimes \sigma_1 & \tilde{T}^3 &= -\frac{1}{2}\sigma_3 \otimes \sigma_3. \end{aligned} \quad (\text{G.9})$$

and realise the inner-product again as (G.3).

Recall that in its defining representation $SU(1, 1)$ consists of complex matrices of unit determinant that satisfy

$$g^\dagger \omega g = \omega, \quad \omega = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (\text{G.10})$$

and such a group element can be parameterized as

$$g = \begin{pmatrix} e^{-it}\sqrt{\rho^2+1} & e^{-i\psi}\rho \\ e^{i\psi}\rho & e^{it}\sqrt{\rho^2+1} \end{pmatrix}. \quad (\text{G.11})$$

In the 4×4 representation used for the Drinfel'd double we have

$$G = \begin{pmatrix} g & 0 \\ 0 & \text{ad}_\omega g \end{pmatrix}. \quad (\text{G.12})$$

The left-invariant one forms are given as

$$\frac{1}{2}(e^1 \pm ie^2) = \frac{e^{\pm i(t-\psi)}}{\sqrt{1+\rho^2}}(\rho(1+\rho^2)d(t+\psi) \pm id\rho), \quad \frac{e^3}{2} = -(1+\rho^2)dt - \rho^2 d\psi. \quad (\text{G.13})$$

The combination of adjoint actions that enter into the PL sigma models is

$$\Pi = \begin{pmatrix} 0 & -2\rho^2 & 2\rho\sqrt{\rho^2+1}\sin(t-\psi) \\ 2\rho^2 & 0 & -2\rho\sqrt{\rho^2+1}\cos(t-\psi) \\ -2\rho\sqrt{\rho^2+1}\sin(t-\psi) & 2\rho\sqrt{\rho^2+1}\cos(t-\psi) & 0 \end{pmatrix}. \quad (\text{G.14})$$

The metric on AdS_3 is obtained as

$$\begin{aligned} \frac{1}{4}\text{Tr}(G^{-1}dGG^{-1}dG) &= -(1+\rho^2)dt^2 + \frac{d\rho^2}{1+\rho^2} + \rho^2 d\psi^2 \\ &= -\cosh^2 \sigma dt^2 + d\sigma^2 + \sinh^2 \sigma d\psi^2, \end{aligned} \quad (\text{G.15})$$

with $\rho = \sinh \sigma$. This follows from the embedding

$$-1 = -X_0^2 + X_1^2 + X_2^2 - X_3^2 \quad (\text{G.16})$$

with

$$X_0 + iX_3 = e^{it} \cosh \sigma, \quad X_1 + iX_2 = e^{-i\psi} \sinh \sigma. \quad (\text{G.17})$$

Although we shall not directly need it we note for completeness the isomorphism to $SL(2, \mathbb{R})$ is made by defining

$$g_{SL(2)} = \begin{pmatrix} X_0 + X_1 & X_2 + X_3 \\ X_2 - X_3 & X_0 - X_1 \end{pmatrix}. \quad (\text{G.18})$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] X.C. de la Ossa and F. Quevedo, *Duality symmetries from nonAbelian isometries in string theory*, *Nucl. Phys. B* **403** (1993) 377 [[hep-th/9210021](#)] [[INSPIRE](#)].
- [2] C. Klimčík and P. Ševera, *Dual nonAbelian duality and the Drinfeld double*, *Phys. Lett. B* **351** (1995) 455 [[hep-th/9502122](#)] [[INSPIRE](#)].
- [3] C. Klimčík and P. Ševera, *Poisson-Lie T duality and loop groups of Drinfeld doubles*, *Phys. Lett. B* **372** (1996) 65 [[hep-th/9512040](#)] [[INSPIRE](#)].
- [4] Y. Lozano, *NonAbelian duality and canonical transformations*, *Phys. Lett. B* **355** (1995) 165 [[hep-th/9503045](#)] [[INSPIRE](#)].
- [5] K. Sfetsos, *Poisson-Lie T duality and supersymmetry*, *Nucl. Phys. Proc. Suppl. B* **56** (1997) 302 [[hep-th/9611199](#)] [[INSPIRE](#)].
- [6] K. Sfetsos, *Canonical equivalence of nonisometric σ -models and Poisson-Lie T duality*, *Nucl. Phys. B* **517** (1998) 549 [[hep-th/9710163](#)] [[INSPIRE](#)].
- [7] M. Roček and E.P. Verlinde, *Duality, quotients and currents*, *Nucl. Phys. B* **373** (1992) 630 [[hep-th/9110053](#)] [[INSPIRE](#)].
- [8] B. Fraser, D. Manolopoulos and K. Sfetsos, *Non-Abelian T-duality and Modular Invariance*, *Nucl. Phys. B* **934** (2018) 498 [[arXiv:1805.03657](#)] [[INSPIRE](#)].
- [9] K. Sfetsos and D.C. Thompson, *On non-abelian T-dual geometries with Ramond fluxes*, *Nucl. Phys. B* **846** (2011) 21 [[arXiv:1012.1320](#)] [[INSPIRE](#)].
- [10] Y. Lozano, E. O Colgáin, K. Sfetsos and D.C. Thompson, *Non-abelian T-duality, Ramond Fields and Coset Geometries*, *JHEP* **06** (2011) 106 [[arXiv:1104.5196](#)] [[INSPIRE](#)].
- [11] G. Itsios, C. Núñez, K. Sfetsos and D.C. Thompson, *Non-Abelian T-duality and the AdS/CFT correspondence:new $N = 1$ backgrounds*, *Nucl. Phys. B* **873** (2013) 1 [[arXiv:1301.6755](#)] [[INSPIRE](#)].
- [12] Y. Lozano and C. Núñez, *Field theory aspects of non-Abelian T-duality and $\mathcal{N} = 2$ linear quivers*, *JHEP* **05** (2016) 107 [[arXiv:1603.04440](#)] [[INSPIRE](#)].
- [13] C. Klimčík, *Yang-Baxter σ -models and dS/AdS T duality*, *JHEP* **12** (2002) 051 [[hep-th/0210095](#)] [[INSPIRE](#)].
- [14] F. Delduc, M. Magro and B. Vicedo, *Derivation of the action and symmetries of the q-deformed $AdS_5 \times S^5$ superstring*, *JHEP* **10** (2014) 132 [[arXiv:1406.6286](#)] [[INSPIRE](#)].
- [15] F. Delduc, M. Magro and B. Vicedo, *An integrable deformation of the $AdS_5 \times S^5$ superstring action*, *Phys. Rev. Lett.* **112** (2014) 051601 [[arXiv:1309.5850](#)] [[INSPIRE](#)].
- [16] K. Sfetsos, *Integrable interpolations: From exact CFTs to non-Abelian T-duals*, *Nucl. Phys. B* **880** (2014) 225 [[arXiv:1312.4560](#)] [[INSPIRE](#)].
- [17] T.J. Hollowood, J.L. Miramontes and D.M. Schmidt, *An Integrable Deformation of the $AdS_5 \times S^5$ Superstring*, *J. Phys. A* **47** (2014) 495402 [[arXiv:1409.1538](#)] [[INSPIRE](#)].

- [18] B. Hoare and A.A. Tseytlin, *On integrable deformations of superstring σ -models related to $AdS_n \times S^n$ supercosets*, *Nucl. Phys. B* **897** (2015) 448 [[arXiv:1504.07213](#)] [[INSPIRE](#)].
- [19] K. Sfetsos, K. Siampos and D.C. Thompson, *Generalised integrable λ - and η -deformations and their relation*, *Nucl. Phys. B* **899** (2015) 489 [[arXiv:1506.05784](#)] [[INSPIRE](#)].
- [20] C. Klimčík, *Poisson-Lie T-duals of the bi-Yang-Baxter models*, *Phys. Lett. B* **760** (2016) 345 [[arXiv:1606.03016](#)] [[INSPIRE](#)].
- [21] A.A. Tseytlin, *Duality Symmetric Formulation of String World Sheet Dynamics*, *Phys. Lett. B* **242** (1990) 163 [[INSPIRE](#)].
- [22] A.A. Tseytlin, *Duality symmetric closed string theory and interacting chiral scalars*, *Nucl. Phys. B* **350** (1991) 395 [[INSPIRE](#)].
- [23] C.M. Hull, *Doubled Geometry and T-Folds*, *JHEP* **07** (2007) 080 [[hep-th/0605149](#)] [[INSPIRE](#)].
- [24] K. Lee and J.-H. Park, *Covariant action for a string in “doubled yet gauged” spacetime*, *Nucl. Phys. B* **880** (2014) 134 [[arXiv:1307.8377](#)] [[INSPIRE](#)].
- [25] M. Hatsuda and W. Siegel, *$O(D, D)$ gauge fields in the T-dual string Lagrangian*, [arXiv:1810.04761](#) [[INSPIRE](#)].
- [26] C.M. Hull and R.A. Reid-Edwards, *Non-geometric backgrounds, doubled geometry and generalised T-duality*, *JHEP* **09** (2009) 014 [[arXiv:0902.4032](#)] [[INSPIRE](#)].
- [27] D. Lüst and D. Osten, *Generalised fluxes, Yang-Baxter deformations and the $O(d, d)$ structure of non-abelian T-duality*, *JHEP* **05** (2018) 165 [[arXiv:1803.03971](#)] [[INSPIRE](#)].
- [28] V.E. Marotta, F. Pezzella and P. Vitale, *Doubling, T-duality and Generalized Geometry: a Simple Model*, *JHEP* **08** (2018) 185 [[arXiv:1804.00744](#)] [[INSPIRE](#)].
- [29] C. Klimčík, *η and λ deformations as \mathcal{E} -models*, *Nucl. Phys. B* **900** (2015) 259 [[arXiv:1508.05832](#)] [[INSPIRE](#)].
- [30] A. Stern, *Hamiltonian approach to Poisson Lie T-duality*, *Phys. Lett. B* **450** (1999) 141 [[hep-th/9811256](#)] [[INSPIRE](#)].
- [31] C. Hull and B. Zwiebach, *Double Field Theory*, *JHEP* **09** (2009) 099 [[arXiv:0904.4664](#)] [[INSPIRE](#)].
- [32] S.D. Avramis, J.-P. Derendinger and N. Prezas, *Conformal chiral boson models on twisted doubled tori and non-geometric string vacua*, *Nucl. Phys. B* **827** (2010) 281 [[arXiv:0910.0431](#)] [[INSPIRE](#)].
- [33] K. Sfetsos, K. Siampos and D.C. Thompson, *Renormalization of Lorentz non-invariant actions and manifest T-duality*, *Nucl. Phys. B* **827** (2010) 545 [[arXiv:0910.1345](#)] [[INSPIRE](#)].
- [34] J. Scherk and J.H. Schwarz, *Spontaneous Breaking of Supersymmetry Through Dimensional Reduction*, *Phys. Lett. B* **82** (1979) 60 [[INSPIRE](#)].
- [35] J. Scherk and J.H. Schwarz, *How to Get Masses from Extra Dimensions*, *Nucl. Phys. B* **153** (1979) 61 [[INSPIRE](#)].
- [36] M. Graña and D. Marques, *Gauged Double Field Theory*, *JHEP* **04** (2012) 020 [[arXiv:1201.2924](#)] [[INSPIRE](#)].
- [37] D. Geissbuhler, *Double Field Theory and $N = 4$ Gauged Supergravity*, *JHEP* **11** (2011) 116 [[arXiv:1109.4280](#)] [[INSPIRE](#)].

- [38] G. Aldazabal, W. Baron, D. Marques and C. Núñez, *The effective action of Double Field Theory*, *JHEP* **11** (2011) 052 [Erratum *ibid.* **11** (2011) 109] [[arXiv:1109.0290](#)] [[INSPIRE](#)].
- [39] D.S. Berman, E.T. Musaev, D.C. Thompson and D.C. Thompson, *Duality Invariant M-theory: Gauged supergravities and Scherk-Schwarz reductions*, *JHEP* **10** (2012) 174 [[arXiv:1208.0020](#)] [[INSPIRE](#)].
- [40] E.T. Musaev, *Gauged supergravities in 5 and 6 dimensions from generalised Scherk-Schwarz reductions*, *JHEP* **05** (2013) 161 [[arXiv:1301.0467](#)] [[INSPIRE](#)].
- [41] R. Blumenhagen, F. Hassler and D. Lüüst, *Double Field Theory on Group Manifolds*, *JHEP* **02** (2015) 001 [[arXiv:1410.6374](#)] [[INSPIRE](#)].
- [42] R. Blumenhagen, P. du Bosque, F. Hassler and D. Lüüst, *Generalized Metric Formulation of Double Field Theory on Group Manifolds*, *JHEP* **08** (2015) 056 [[arXiv:1502.02428](#)] [[INSPIRE](#)].
- [43] P. du Bosque, F. Hassler and D. Lüüst, *Flux Formulation of DFT on Group Manifolds and Generalized Scherk-Schwarz Compactifications*, *JHEP* **02** (2016) 039 [[arXiv:1509.04176](#)] [[INSPIRE](#)].
- [44] F. Hassler, *Poisson-Lie T-duality in Double Field Theory*, [arXiv:1707.08624](#) [[INSPIRE](#)].
- [45] P. du Bosque, F. Hassler and D. Lüüst, *Generalized parallelizable spaces from exceptional field theory*, *JHEP* **01** (2018) 117 [[arXiv:1705.09304](#)] [[INSPIRE](#)].
- [46] M. Graña, R. Minasian, M. Petrini and D. Waldram, *T-duality, Generalized Geometry and Non-Geometric Backgrounds*, *JHEP* **04** (2009) 075 [[arXiv:0807.4527](#)] [[INSPIRE](#)].
- [47] K. Lee, C. Strickland-Constable and D. Waldram, *Spheres, generalised parallelisability and consistent truncations*, *Fortsch. Phys.* **65** (2017) 1700048 [[arXiv:1401.3360](#)] [[INSPIRE](#)].
- [48] G. Arutyunov, S. Frolov, B. Hoare, R. Roiban and A.A. Tseytlin, *Scale invariance of the η -deformed $AdS_5 \times S^5$ superstring, T-duality and modified type-II equations*, *Nucl. Phys. B* **903** (2016) 262 [[arXiv:1511.05795](#)] [[INSPIRE](#)].
- [49] B. Hoare, R. Roiban and A.A. Tseytlin, *On deformations of $AdS_n \times S^n$ supercosets*, *JHEP* **06** (2014) 002 [[arXiv:1403.5517](#)] [[INSPIRE](#)].
- [50] R. Borsato and L. Wulff, *Target space supergeometry of η and λ -deformed strings*, *JHEP* **10** (2016) 045 [[arXiv:1608.03570](#)] [[INSPIRE](#)].
- [51] Y. Sakatani, S. Uehara and K. Yoshida, *Generalized gravity from modified DFT*, *JHEP* **04** (2017) 123 [[arXiv:1611.05856](#)] [[INSPIRE](#)].
- [52] A. Baguet, M. Magro and H. Samtleben, *Generalized IIB supergravity from exceptional field theory*, *JHEP* **03** (2017) 100 [[arXiv:1612.07210](#)] [[INSPIRE](#)].
- [53] T. Araujo, E. Ó Colgáin, J. Sakamoto, M.M. Sheikh-Jabbari and K. Yoshida, *I in generalized supergravity*, *Eur. Phys. J. C* **77** (2017) 739 [[arXiv:1708.03163](#)] [[INSPIRE](#)].
- [54] I. Bakhmatov, E. Ó Colgáin, M.M. Sheikh-Jabbari and H. Yavartanoo, *Yang-Baxter Deformations Beyond Coset Spaces (a slick way to do TsT)*, *JHEP* **06** (2018) 161 [[arXiv:1803.07498](#)] [[INSPIRE](#)].
- [55] T. Araujo, E. Ó Colgáin and H. Yavartanoo, *Embedding the modified CYBE in Supergravity*, *Eur. Phys. J. C* **78** (2018) 854 [[arXiv:1806.02602](#)] [[INSPIRE](#)].
- [56] P. Ševera and F. Valach, *Courant algebroids, Poisson-Lie T-duality and type-II supergravities*, [arXiv:1810.07763](#) [[INSPIRE](#)].

- [57] C. Klimčík, *On integrability of the Yang-Baxter σ -model*, *J. Math. Phys.* **50** (2009) 043508 [[arXiv:0802.3518](#)] [[INSPIRE](#)].
- [58] I. Kawaguchi, T. Matsumoto and K. Yoshida, *Jordanian deformations of the $AdS_5 \times S^5$ superstring*, *JHEP* **04** (2014) 153 [[arXiv:1401.4855](#)] [[INSPIRE](#)].
- [59] T. Matsumoto and K. Yoshida, *Yang-Baxter σ -models based on the CYBE*, *Nucl. Phys. B* **893** (2015) 287 [[arXiv:1501.03665](#)] [[INSPIRE](#)].
- [60] B. Hoare and A.A. Tseytlin, *Homogeneous Yang-Baxter deformations as non-abelian duals of the AdS_5 σ -model*, *J. Phys. A* **49** (2016) 494001 [[arXiv:1609.02550](#)] [[INSPIRE](#)].
- [61] D. Osten and S.J. van Tongeren, *Abelian Yang-Baxter deformations and TsT transformations*, *Nucl. Phys. B* **915** (2017) 184 [[arXiv:1608.08504](#)] [[INSPIRE](#)].
- [62] O. Hohm, C. Hull and B. Zwiebach, *Generalized metric formulation of double field theory*, *JHEP* **08** (2010) 008 [[arXiv:1006.4823](#)] [[INSPIRE](#)].
- [63] O. Hohm, S.K. Kwak and B. Zwiebach, *Double Field Theory of Type II Strings*, *JHEP* **09** (2011) 013 [[arXiv:1107.0008](#)] [[INSPIRE](#)].
- [64] C. Klimčík and P. Ševera, *NonAbelian momentum winding exchange*, *Phys. Lett. B* **383** (1996) 281 [[hep-th/9605212](#)] [[INSPIRE](#)].
- [65] F. Hassler, *The Topology of Double Field Theory*, *JHEP* **04** (2018) 128 [[arXiv:1611.07978](#)] [[INSPIRE](#)].
- [66] Y. Kosmann-Schwarzbach, *Quasi, twisted, and all that... in Poisson geometry and Lie algebroid theory*, [math/0310359](#).
- [67] F. Hassler and F. Rudolph, *Poisson-Lie σ -models and Para-Hermitian Geometries*, work in progress.
- [68] L. Freidel, F.J. Rudolph and D. Svoboda, *A Unique Connection for Born Geometry*, [arXiv:1806.05992](#) [[INSPIRE](#)].
- [69] V.E. Marotta and R.J. Szabo, *Para-Hermitian Geometry, Dualities and Generalized Flux Backgrounds*, [arXiv:1810.03953](#) [[INSPIRE](#)].
- [70] D. Svoboda, *Algebroid Structures on Para-Hermitian Manifolds*, *J. Math. Phys.* **59** (2018) 122302 [[arXiv:1802.08180](#)] [[INSPIRE](#)].
- [71] O. Hohm, S.K. Kwak and B. Zwiebach, *Unification of Type II Strings and T-duality*, *Phys. Rev. Lett.* **107** (2011) 171603 [[arXiv:1106.5452](#)] [[INSPIRE](#)].
- [72] J.-i. Sakamoto, Y. Sakatani and K. Yoshida, *Weyl invariance for generalized supergravity backgrounds from the doubled formalism*, *PTEP* **2017** (2017) 053B07 [[arXiv:1703.09213](#)] [[INSPIRE](#)].
- [73] B. Jurčo and J. Vysoky, *Poisson-Lie T-duality of string effective actions: A new approach to the dilaton puzzle*, *J. Geom. Phys.* **130** (2018) 1 [[arXiv:1708.04079](#)] [[INSPIRE](#)].
- [74] K. Hori, *D-branes, T duality and index theory*, *Adv. Theor. Math. Phys.* **3** (1999) 281 [[hep-th/9902102](#)] [[INSPIRE](#)].
- [75] E. Gevorgyan and G. Sarkissian, *Defects, Non-abelian T-duality and the Fourier-Mukai transform of the Ramond-Ramond fields*, *JHEP* **03** (2014) 035 [[arXiv:1310.1264](#)] [[INSPIRE](#)].
- [76] G. Georgiou, K. Sfetsos and K. Siampos, *Double and cyclic λ -deformations and their canonical equivalents*, *Phys. Lett. B* **771** (2017) 576 [[arXiv:1704.07834](#)] [[INSPIRE](#)].

- [77] G. Georgiou and K. Sfetsos, *A new class of integrable deformations of CFTs*, *JHEP* **03** (2017) 083 [[arXiv:1612.05012](#)] [[INSPIRE](#)].
- [78] F. Delduc, M. Magro and B. Vicedo, *Integrable double deformation of the principal chiral model*, *Nucl. Phys. B* **891** (2015) 312 [[arXiv:1410.8066](#)] [[INSPIRE](#)].
- [79] S. Demulder, S. Driezen, A. Sevrin and D.C. Thompson, *Classical and Quantum Aspects of Yang-Baxter Wess-Zumino Models*, *JHEP* **03** (2018) 041 [[arXiv:1711.00084](#)] [[INSPIRE](#)].
- [80] M.J. Duff, *Kaluza-Klein theory in perspective*, in *The Oskar Klein centenary. Proceedings, Symposium*, Stockholm, Sweden, September 19–21, 1994, pp. 22–35 (1994), [[hep-th/9410046](#)] [[INSPIRE](#)].
- [81] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Consistent group and coset reductions of the bosonic string*, *Class. Quant. Grav.* **20** (2003) 5161 [[hep-th/0306043](#)] [[INSPIRE](#)].
- [82] S. Demulder, F. Hassler and D.C. Thompson, *Doubled aspects of generalised dualities: the coset case*, work in progress.
- [83] S.F. Hassan, *T duality, space-time spinors and RR fields in curved backgrounds*, *Nucl. Phys. B* **568** (2000) 145 [[hep-th/9907152](#)] [[INSPIRE](#)].
- [84] Ö. Kelekci, Y. Lozano, N.T. Macpherson and E.Ó. Colgáin, *Supersymmetry and non-Abelian T-duality in type-II supergravity*, *Class. Quant. Grav.* **32** (2015) 035014 [[arXiv:1409.7406](#)] [[INSPIRE](#)].
- [85] R. Minasian and G.W. Moore, *K theory and Ramond-Ramond charge*, *JHEP* **11** (1997) 002 [[hep-th/9710230](#)] [[INSPIRE](#)].
- [86] E. Witten, *D-branes and k-theory*, *JHEP* **12** (1998) 019 [[hep-th/9810188](#)] [[INSPIRE](#)].
- [87] O. Hohm and B. Zwiebach, *Double field theory at order α'* , *JHEP* **11** (2014) 075 [[arXiv:1407.3803](#)] [[INSPIRE](#)].
- [88] O. Hohm and B. Zwiebach, *Green-Schwarz mechanism and α' -deformed Courant brackets*, *JHEP* **01** (2015) 012 [[arXiv:1407.0708](#)] [[INSPIRE](#)].
- [89] A. Deser, *Star products on graded manifolds and α' -corrections to Courant algebroids from string theory*, *J. Math. Phys.* **56** (2015) 092302 [[arXiv:1412.5966](#)] [[INSPIRE](#)].
- [90] B. Vicedo, *Deformed integrable σ -models, classical R-matrices and classical exchange algebra on Drinfel'd doubles*, *J. Phys. A* **48** (2015) 355203 [[arXiv:1504.06303](#)] [[INSPIRE](#)].
- [91] J. Shelton, W. Taylor and B. Wecht, *Nongeometric flux compactifications*, *JHEP* **10** (2005) 085 [[hep-th/0508133](#)] [[INSPIRE](#)].
- [92] K. Sfetsos and D.C. Thompson, *Spacetimes for λ -deformations*, *JHEP* **12** (2014) 164 [[arXiv:1410.1886](#)] [[INSPIRE](#)].