

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,6,8-Tribromoquinoline

 İsmail Çelik,^a Mehmet Akkurt,^{b*} Salih Ökten,^c Osman Çakmak^c and Santiago García-Granda^d

^aDepartment of Physics, Faculty of Arts and Sciences, Cumhuriyet University, 58140 Sivas, Turkey, ^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^cDepartment of Chemistry, Faculty of Art and Science, Gaziosmanpaşa University, 60240 Tokat, Turkey, and ^dDepartamento Química Física y Analítica, Facultad de Química, Universidad Oviedo, C/ Julián Clavería, 8, 33006 Oviedo (Asturias), Spain
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 28 October 2010; accepted 5 November 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.054; wR factor = 0.150; data-to-parameter ratio = 15.4.

The title molecule, $\text{C}_9\text{H}_4\text{Br}_3\text{N}$, is almost planar, the maximum deviation being 0.110 (1) Å. The crystal structure is stabilized by weak aromatic π - π interactions [centroid-centroid distance = 3.802 (4) Å] between the pyridine and benzene rings of the quinoline ring systems of adjacent molecules.

Related literature

For background to the synthesis of natural biologically active quinoline derivatives and for the synthesis of the title compound, see: Şahin *et al.* (2008). For the structure of 6,8-dibromoquinoline, see: Çelik *et al.* (2010).



Experimental

Crystal data

 $\text{C}_9\text{H}_4\text{Br}_3\text{N}$
 $M_r = 365.83$

Monoclinic, $P2_1/n$
 $a = 3.9810$ (2) Å
 $b = 12.4176$ (4) Å
 $c = 19.7419$ (6) Å
 $\beta = 92.827$ (3)°
 $V = 974.74$ (7) Å³

$Z = 4$
 Cu $K\alpha$ radiation
 $\mu = 14.93$ mm⁻¹
 $T = 296$ K
 $0.51 \times 0.06 \times 0.03$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby Gemini CCD detector
 Absorption correction: multi-scan (*CrysAlis PRO*; Oxford)

Diffraction, 2010)
 $T_{\min} = 0.049$, $T_{\max} = 0.663$
 3688 measured reflections
 1816 independent reflections
 1484 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.060$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.150$
 $S = 1.05$
 1816 reflections

118 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 1.45$ e Å⁻³
 $\Delta\rho_{\min} = -0.80$ e Å⁻³

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1999); software used to prepare material for publication: *WinGX* (Farrugia, 1997) and *PLATON* (Spek, 2009).

The authors thank the Cumhuriyet University Research Foundation (CUBAP grant No. 2009/ F-266) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2350).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
 Çelik, İ., Akkurt, M., Çakmak, O., Ökten, S. & García-Granda, S. (2010). *Acta Cryst.* **E66**, o2997–o2998.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
 Şahin, A. O., Çakmak, O., Demirtaş, I., Ökten, S. & Tutar, A. (2008). *Tetrahedron*, **64**, 10068–10074.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2010). E66, o3133 [doi:10.1107/S1600536810045484]

3,6,8-Tribromoquinoline

İsmail Çelik, Mehmet Akkurt, Salih Ökten, Osman Çakmak and Santiago García-Granda

S1. Comment

The presence of quinoline skeleton in the framework of pharmacologically active compounds and natural products has spurred on the development of different strategies for their synthesis. The lithium–halogen exchange reaction of the title compound (I) may serve for the synthesis of natural biologically active quinoline derivatives, such as quinine, pentaquine, and plasmoquine (Şahin *et al.*, 2008). In this paper we report a one pot synthesis of (I) with high yield (90%) and its crystal structure.

The title molecule is almost planar, with the maximum and minimum deviations from the mean plane being 0.110 (1) and -0.001 (6) Å for Br2 and C4, respectively. Its crystal structure is stabilized by weak π – π stacking interactions between the pyridine and benzene rings of the quinoline ring systems of the adjacent molecules [$Cg1 \cdots Cg2^i = 3.802$ (4) Å; symmetry code: (i) $1 + x, y, z$; Cg1 and Cg2 are centroids of the N1/C1/C6–C9 pyridine and C1–C6 benzene rings of the quinoline ring system, respectively].


The crystal structure of 6,8-dibromoquinoline has been reported recently Çelik *et al.* (2010).

S2. Experimental

6,8-Dibromo-1,2,3,4-tetrahydroquinoline was synthesized according to the literature method (Şahin *et al.*, 2008). To a solution of 6,8-dibromo-1,2,3,4-tetrahydroquinoline (0.5 g, 3.75 mmol, 1 eq) in $CHCl_3$ (20 ml) was dropped bromine (1.8 g, 11.25 mmol, 3 eq) in $CHCl_3$ (10 ml) over 5 min in the dark and at room temperature. After completion of the reaction (bromine consumed completely, 3 days), the solid was dissolved in $CHCl_3$ (35 ml) and the organic layer was washed with 5% $NaHCO_3$ solution (3x20 ml) and dried over Na_2SO_4 . After evaporation of the solvent, the crude material (1.32 g) was passed through a short alumina column eluting with EtOAc–hexane (1:12, 75 ml) (hexane/ethyl acetate, 9:1, $R_f = 0.65$). Colourless solid residue was obtained. The mixture was recrystallized from the solvent (benzene) in a freezer (263 K) to give pure 3,6,8-tribromoquinoline in 90% yield (1.24 g) if the form of colourless needle shaped crystals; m.p. 441–443 K.

S3. Refinement

H atoms were included in geometric positions with $C-H = 0.93$ Å and refined by using a riding model [$U_{iso}(H) = 1.2U_{eq}(C)$]. The highest peak in the final difference map was located 0.92 Å from Br2, while the deepest hole was located 1.05 Å from Br3.

**Figure 1**

The title molecule with the atom numbering scheme. Displacement ellipsoids for have been drawn at the 50% probability level.

3,6,8-Tribromoquinoline

Crystal data

$C_9H_4Br_3N$

$M_r = 365.83$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 3.9810(2)\ \text{\AA}$

$b = 12.4176(4)\ \text{\AA}$

$c = 19.7419(6)\ \text{\AA}$

$\beta = 92.827(3)^\circ$

$V = 974.74(7)\ \text{\AA}^3$

$Z = 4$

$F(000) = 680$

$D_x = 2.493\ \text{Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.5418\ \text{\AA}$

Cell parameters from 2025 reflections

$\theta = 3.6\text{--}70.4^\circ$

$\mu = 14.93\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Needle, colourless

$0.51 \times 0.06 \times 0.03\ \text{mm}$

Data collection

Oxford Diffraction Xcalibur

diffractometer with a Ruby Gemini CCD

detector

Radiation source: Enhance (Cu) X-ray Source

Graphite monochromator

Detector resolution: $10.2673\ \text{pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Oxford Diffraction, 2010)

$T_{\min} = 0.049$, $T_{\max} = 0.663$

3688 measured reflections

1816 independent reflections

1484 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.060$

$\theta_{\max} = 70.6^\circ$, $\theta_{\min} = 5.7^\circ$

$h = -4 \rightarrow 4$

$k = -9 \rightarrow 15$

$l = -22 \rightarrow 24$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.150$
 $S = 1.05$
 1816 reflections
 118 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1066P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.45 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.80 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.6380 (2)	0.29099 (6)	0.90189 (4)	0.0568 (3)
Br2	0.11453 (19)	-0.12590 (6)	0.92944 (4)	0.0525 (3)
Br3	0.7497 (2)	0.08258 (7)	0.56672 (4)	0.0608 (3)
N1	0.7203 (14)	0.2168 (5)	0.7553 (3)	0.0440 (17)
C1	0.5666 (16)	0.1396 (5)	0.7921 (3)	0.0396 (17)
C2	0.5134 (15)	0.1576 (5)	0.8619 (3)	0.0404 (17)
C3	0.3758 (16)	0.0808 (5)	0.9010 (3)	0.0428 (17)
C4	0.2822 (16)	-0.0184 (5)	0.8725 (3)	0.0425 (17)
C5	0.3078 (15)	-0.0384 (5)	0.8042 (3)	0.0405 (17)
C6	0.4583 (15)	0.0399 (5)	0.7636 (3)	0.0387 (17)
C7	0.5126 (16)	0.0209 (5)	0.6946 (3)	0.0426 (17)
C8	0.6665 (16)	0.0996 (5)	0.6597 (3)	0.0429 (17)
C9	0.7713 (17)	0.1961 (6)	0.6919 (3)	0.0462 (17)
H3	0.34400	0.09420	0.94660	0.0520*
H5	0.22720	-0.10250	0.78520	0.0480*
H7	0.44560	-0.04330	0.67380	0.0510*
H9	0.88170	0.24730	0.66670	0.0550*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0752 (6)	0.0456 (4)	0.0506 (5)	-0.0125 (3)	0.0119 (4)	-0.0120 (3)
Br2	0.0622 (5)	0.0498 (5)	0.0464 (4)	-0.0097 (3)	0.0107 (3)	0.0057 (3)
Br3	0.0798 (6)	0.0652 (5)	0.0385 (4)	0.0047 (4)	0.0132 (3)	-0.0003 (3)
N1	0.053 (3)	0.040 (3)	0.039 (3)	-0.003 (2)	0.003 (2)	0.002 (2)

C1	0.043 (3)	0.032 (3)	0.044 (3)	0.002 (2)	0.003 (2)	0.002 (2)
C2	0.044 (3)	0.037 (3)	0.040 (3)	0.002 (2)	0.000 (2)	-0.004 (2)
C3	0.042 (3)	0.050 (3)	0.037 (3)	-0.003 (3)	0.007 (2)	-0.006 (3)
C4	0.046 (3)	0.039 (3)	0.043 (3)	0.000 (3)	0.006 (2)	0.003 (2)
C5	0.045 (3)	0.038 (3)	0.038 (3)	0.000 (2)	-0.004 (2)	0.000 (2)
C6	0.039 (3)	0.038 (3)	0.039 (3)	0.006 (2)	0.000 (2)	0.003 (2)
C7	0.048 (3)	0.041 (3)	0.039 (3)	0.007 (3)	0.005 (2)	-0.003 (2)
C8	0.045 (3)	0.047 (3)	0.037 (3)	0.010 (3)	0.004 (2)	0.003 (2)
C9	0.053 (3)	0.047 (3)	0.039 (3)	-0.002 (3)	0.006 (3)	0.004 (2)

Geometric parameters (Å, °)

Br1—C2	1.891 (6)	C4—C5	1.380 (8)
Br2—C4	1.888 (6)	C5—C6	1.412 (9)
Br3—C8	1.893 (6)	C6—C7	1.410 (8)
N1—C1	1.366 (9)	C7—C8	1.359 (9)
N1—C9	1.303 (8)	C8—C9	1.410 (9)
C1—C2	1.422 (8)	C3—H3	0.9300
C1—C6	1.418 (9)	C5—H5	0.9300
C2—C3	1.359 (9)	C7—H7	0.9300
C3—C4	1.397 (9)	C9—H9	0.9300
Br1...N1	3.070 (6)	C7...C8 ⁱⁱ	3.542 (9)
Br1...Br3 ⁱ	3.6969 (12)	C7...Br1 ^{iv}	3.738 (6)
Br1...C7 ⁱ	3.738 (6)	C8...C7 ^{viii}	3.542 (9)
Br2...C4 ⁱⁱ	3.696 (6)	C9...Br2 ^{ix}	3.553 (7)
Br2...C9 ⁱⁱⁱ	3.553 (7)	C9...C6 ^{viii}	3.587 (9)
Br3...Br1 ^{iv}	3.6969 (12)	C5...H9 ^{iv}	2.9800
Br3...Br3 ^v	3.8186 (12)	H3...Br2 ^{vii}	3.1500
Br1...H7 ⁱ	3.0800	H3...Br2 ^{vi}	3.2000
Br2...H9 ⁱⁱⁱ	3.1000	H5...H7	2.5100
Br2...H3 ^{vi}	3.2000	H5...H9 ^{iv}	2.5800
Br2...H9 ^{iv}	3.2400	H7...H5	2.5100
Br2...H3 ^{vii}	3.1500	H7...Br1 ^{iv}	3.0800
N1...Br1	3.070 (6)	H9...Br2 ^{ix}	3.1000
C4...Br2 ^{viii}	3.696 (6)	H9...Br2 ⁱ	3.2400
C5...C6 ⁱⁱ	3.572 (8)	H9...C5 ⁱ	2.9800
C6...C5 ^{viii}	3.572 (8)	H9...H5 ⁱ	2.5800
C6...C9 ⁱⁱ	3.587 (9)		
C1—N1—C9	117.9 (6)	C5—C6—C7	121.5 (6)
N1—C1—C2	119.8 (6)	C6—C7—C8	117.7 (6)
N1—C1—C6	122.5 (6)	Br3—C8—C7	121.1 (5)
C2—C1—C6	117.7 (5)	Br3—C8—C9	118.0 (5)
Br1—C2—C1	119.6 (5)	C7—C8—C9	120.9 (6)
Br1—C2—C3	118.8 (5)	N1—C9—C8	123.0 (6)
C1—C2—C3	121.6 (6)	C2—C3—H3	120.00
C2—C3—C4	119.8 (6)	C4—C3—H3	120.00

Br2—C4—C3	118.6 (4)	C4—C5—H5	120.00
Br2—C4—C5	120.0 (5)	C6—C5—H5	121.00
C3—C4—C5	121.4 (6)	C6—C7—H7	121.00
C4—C5—C6	119.0 (6)	C8—C7—H7	121.00
C1—C6—C5	120.3 (5)	N1—C9—H9	119.00
C1—C6—C7	118.1 (5)	C8—C9—H9	118.00
C9—N1—C1—C2	178.4 (6)	C2—C3—C4—Br2	176.6 (5)
C9—N1—C1—C6	-0.9 (9)	C2—C3—C4—C5	-3.9 (10)
C1—N1—C9—C8	1.9 (10)	Br2—C4—C5—C6	-175.1 (5)
N1—C1—C2—Br1	2.9 (8)	C3—C4—C5—C6	5.4 (9)
N1—C1—C2—C3	-176.9 (6)	C4—C5—C6—C1	-3.0 (9)
C6—C1—C2—Br1	-177.8 (4)	C4—C5—C6—C7	175.5 (6)
C6—C1—C2—C3	2.5 (9)	C1—C6—C7—C8	0.1 (9)
N1—C1—C6—C5	178.5 (6)	C5—C6—C7—C8	-178.4 (6)
N1—C1—C6—C7	-0.1 (9)	C6—C7—C8—Br3	-179.7 (5)
C2—C1—C6—C5	-0.9 (9)	C6—C7—C8—C9	0.8 (9)
C2—C1—C6—C7	-179.5 (6)	Br3—C8—C9—N1	178.6 (5)
Br1—C2—C3—C4	-179.9 (5)	C7—C8—C9—N1	-1.9 (10)
C1—C2—C3—C4	-0.2 (9)		

Symmetry codes: (i) $-x+3/2, y+1/2, -z+3/2$; (ii) $x-1, y, z$; (iii) $-x+1/2, y-1/2, -z+3/2$; (iv) $-x+3/2, y-1/2, -z+3/2$; (v) $-x+1, -y, -z+1$; (vi) $-x+1, -y, -z+2$; (vii) $-x, -y, -z+2$; (viii) $x+1, y, z$; (ix) $-x+1/2, y+1/2, -z+3/2$.