

Phosphane-functionalized heavier tetrylenes: Synthesis of silylene- and germylene-decorated phosphanes and their reactions with Group 10 metal complexes†

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Javier A. Cabeza,^{a*} Pablo García-Álvarez,^a Carlos J. Laglera-Gándara^a and Enrique Pérez-Carreño^b

The stable phosphane-functionalized heavier tetrylenes E(^tBu₂bzam)pyrmP^tBu₂ (E = Si (**1**_{Si}), Ge (**1**_{Ge}); ^tBu₂bzam = *N,N'*-ditertbutylbenzamidinate; HpyrmP^tBu₂ = ditertbutyl(2-pyrrolylmethyl)phosphane) have been prepared by reacting the amidinatotetrylenes E(^tBu₂bzam)Cl (E = Si, Ge) with LipyrmP^tBu₂. The reactions of **1**_{Si} and **1**_{Ge} with selected M⁰ and M^{II} (M = Ni, Pd, Pt) metal precursors have allowed the synthesis of square-planar [MCl₂{κ²E,*P*-E(^tBu₂bzam)pyrmP^tBu₂}] (M = Ni, Pd, Pt; E = Si, Ge), tetrahedral [Ni(κ²E,*P*-E(^tBu₂bzam)pyrmP^tBu₂)(cod)] (E = Si, Ge; cod = 1,5-cyclooctadiene) and triangular [M(κ²E,*P*-E(^tBu₂bzam)pyrmP^tBu₂)(PPh₃)] (M = Pd, Pt; E = Si, Ge) complexes, showing that **1**_{Si} and **1**_{Ge} are excellent *Si*,*P*- and *Ge*,*P*-chelating ligands that, due to their large steric bulk, are able to stabilize three-coordinate Pd⁰ and Pt⁰ complexes.

Introduction

The heavier analogues of carbenes, *i.e.* silylenes, germylenes, stannylenes and plumbylens, also known as heavier tetrylenes (HTs),¹ have been recognized as very strong electron-donating ligands, particularly if they are stabilized by donor groups,² and some of their transition metal (TM) complexes have already demonstrated high efficiency in various catalytic reactions.^{3,4}

Stimulated by the above-mentioned facts, the synthesis and coordination chemistry of potentially bidentate ligands containing HTs as donor groups have recently started to be investigated.^{4a,5-9} However, despite the excellent coordination and catalytic properties of phosphanes, prior to the start of this work (May 2019), the known stable (isolated) phosphane-functionalized HTs that had proven to lead to bidentate κ²*P*,*E*-ligands (E = HT tetrel atom) upon reactions with transition metal complexes were restricted to Wesemann's very bulky stannylene compounds **A–C** (Fig. 1). **A**⁶ and **B**^{6,7} have been reported to react with Ni⁰ and Pd⁰ metal precursors to give products that result from the insertion of the metal atom into the Sn–P bond, while anionic stannylene **C** replaces the nitrile ligands of [PtCl₂(MeCN)₂] and undergoes transmetalation with [Rh₂(μ-Cl)₂(CO)₄] to give the corresponding square-planar complexes.⁸ In some occasions, the reactions of **A**,⁷ **B**⁷ and other⁹ phosphane-functionalized HTs with TM complexes lead




Fig. 1 The currently known stable phosphane-functionalized HTs that have led to bidentate κ²*P*,*E*-ligands (E = HT tetrel atom) upon reactions with TM complexes.

to metal products that no longer contain an HT fragment (it has been transformed into a tetryl ligand by insertion of the HT E atom into an M–X bond).

The absence of phosphane-functionalized silylenes and germylenes having unengaged the lone pairs of their HT and phosphane functionalities and our previous experience on transition metal chemistry of amidinatotetrylenes^{1f,2e,4b,4g,10} and PEP pincer (E = HT tetrel atom) compounds,^{11,12} including those involving pyrrolylphosphane groups,¹² led us to investigate the synthesis of such compounds by combining amidinatotetrylenes with pyrrolylphosphanes. As amidinatotetrylenes are donor-stabilized tetrylenes, they are not prone to be transformed into tetryl ligands upon reactions with TM complexes containing polar M–X bonds.^{1f}

^aCentro de Innovación en Química Avanzada (ORFEO-CINQA) and Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, 33071 Oviedo, Spain.

^bDepartamento de Química Física y Analítica, Universidad de Oviedo, 33071 Oviedo, Spain.

*E-mail: jac@uniovi.es (J.A.C.).


†Electronic Supplementary Information (ESI) available: CCDC 2001500–2001503. NMR spectra and complementary XRD and DFT data. See DOI: 10.1039/x0xx00000x.

We now report that such a synthetic approach has allowed the preparation of a silylene-phosphane and a germylene-phosphane (**1_{Si}** and **1_{Ge}**, respectively; Fig. 1) and that both compounds behave as genuine bidentate κ^2P,E -ligands (E = Si, Ge) in their reactions with Group 10 metal (0 and II) complexes.

When we were finishing the labwork described in this article, Tan, Zhang, Huang et al. reported the synthesis of a germylene-phosphane (**D** in Fig. 1) that can also behave as a bidentate κ^2P,Ge -ligand.¹³

Results and discussion

The reactions of *tert*-butyl(2-pyrrolylmethyl)phosphane^{12a} (HpyrmP^{*t*}Bu₂) first with *n*-butyl lithium and then with E(^{*t*}Bu₂bzam)Cl (E = Si,¹⁴ Ge;¹⁵ ^{*t*}Bu₂bzam = *N,N'*-*tert*-butylbenzamidinate) cleanly led to the silylene- and germylene-decorated phosphanes E(^{*t*}Bu₂bzam)pyrmP^{*t*}Bu₂ (**1_E**; E = Si, Ge) (Scheme 1). Both compounds are very air-sensitive and very soluble in organic solvents (including aliphatic hydrocarbons).



Scheme 1 Synthesis of the heavier tetrelene-phosphane compounds **1_{Si}** and **1_{Ge}**.

The ¹H and ¹³C{¹H} NMR spectra of **1_{Si}** and **1_{Ge}** are very similar and indicate equivalence of their *N*^{*t*}Bu groups, denoting an average C₅ symmetry in solution. The chemical shifts of their ³¹P{¹H} NMR signals, 28.2 (**1_{Si}**) and 28.3 (**1_{Ge}**) ppm, which differ from that of HpyrmP^{*t*}Bu₂ (22.8 ppm) by only 5 ppm,^{12b} hint to the absence of direct P...E interactions. The X-ray diffraction (XRD) structure of silylene **1_{Si}** (Fig. 2) shows that the Si atom is in the apex of a trigonal pyramid, attached to both benzamidinate N atoms and to the pyrrolyl fragment of the pyrmP^{*t*}Bu₂ group. The very long Si1...P1 distance, 4.795(1) Å, discards any interaction between these atoms.




Fig. 2 Molecular structure of compound **1_{Si}** (50% displacement ellipsoids, H atoms omitted for clarity). Selected bond lengths (Å) and angles (°): Si1–N1 1.873(1), Si1–N2 1.869(1), Si1–N3 1.809(1), C4–N1 1.483(2), C5–N1 1.341(2), C5–C6 1.492(2), C5–N2 1.337(2), C12–N2 1.480(2), C19–N3 1.397(2), C19–C20 1.505(2), C20–P1 1.868(1), C21–P1 1.895(1), C25–P1 1.892(2), N1–Si1–N2 69.56(5), N1–Si1–N3 100.47(6), N2–Si1–N3 101.52(5), N1–C5–N2 105.7(1).

A theoretical DFT/NBO analysis (at the wB97XD/cc-pVDZ level) of **1_{Si}** (Table 1) showed that the energy of the orbital containing the lone pair (LP) of the silylene fragment lies 42.66 kcal mol^{−1} above the orbital that contains the LP of the phosphane

fragment, indicating that the basicity of the amidinatosilylene fragment is considerably higher than that of the trialkylphosphane scaffold.^{2c} However, in the case of **1_{Ge}**, the basicity of the amidinatosilylene fragment is slightly smaller than that of the trialkylphosphane scaffold as the germylene LP lies 16.75 kcal mol^{−1} below the phosphane LP (Table 2).^{2a} In addition, the silylene LP has a smaller *s* (and a greater *p*) character than the germylene LP (Tables 1 and 2), also pointing to a stronger basicity of the silylene. The filled molecular orbitals (HOMO, HOMO−1, etc.) that are higher in energy than those containing the tetrelene and phosphane LPs are not relevant to the coordination chemistry of these molecules because they are not LPs (they are π -type orbitals of the amidinate or pyrrole fragments; ESI, Fig. S50).

Table 1 Selected orbital properties of silylene-phosphane **1_{Si}**


Orbital	<i>E</i> (hartree)	ΔE (kcal mol ^{−1})	Composition (%)
LP Si (HOMO−7)	−0.36111	42.66	65.46 (<i>s</i>), 34.45 (<i>p</i>)
LP P (HOMO−11)	−0.42909	0	49.01 (<i>s</i>), 50.98 (<i>p</i>)

Table 2 Selected orbital properties of germylene-phosphane **1_{Ge}**

Orbital	<i>E</i> (hartree)	ΔE (kcal mol ^{−1})	Composition (%)
LP P (HOMO−10)	−0.43002	0	49.07 (<i>s</i>), 50.92 (<i>p</i>)
LP Ge (HOMO−11)	−0.45671	−16.75	81.6 (<i>s</i>), 18.39 (<i>p</i>)

In order to verify and compare the coordination ability of HT-phosphanes **1_{Si}** and **1_{Ge}**, they were reacted with Group 10 metal complexes in +2 and 0 oxidation states.

Both **1_{Si}** and **1_{Ge}** reacted at room temperature with Group 10 metal(II) complexes containing easy displaceable ligands, [NiCl₂(dme)] (dme = 1,2-dimethoxyethane), [PdCl₂(MeCN)₂] and [PtCl₂(cod)] (cod = 1,5-cyclooctadiene), to give the corresponding [MCl₂{ κ^2E,P -E(^{*t*}Bu₂bzam)pyrmP^{*t*}Bu₂}] (**2_{M-E}**; M = Ni, Pd, Pt; E = Si, Ge) reaction products (Scheme 2).



Scheme 2 Reactions of **1_{Si}** and **1_{Ge}** with Group 10 metal dichlorido complexes.

All compounds of type **2_{M-E}** are diamagnetic species with very similar ¹H and ¹³C{¹H} NMR spectra that are compatible with C₅ molecular symmetry in solution. In all cases, the ³¹P resonances appear at higher frequencies than those of the free tetrelenes-phosphanes. Large *J*_{Pt-P}, 3701.2 (**2_{Pt-Si}**) and 3553.2 (**2_{Pt-Ge}**) Hz, and *J*_{Pt-Si}, 2298.4 (**2_{Pt-Si}**) Hz, coupling constants were observed in the ³¹P{¹H} and ²⁹Si{¹H} spectra of the platinum complexes, corroborating the direct attachment of the Pt atom to the Si and P atoms. The XRD molecular structure of **2_{Pd-Si}** (Fig. 3) confirms the square-planar coordination geometry and the chelating ability of the silylene-phosphane ligand. The notably different Pd–Cl1 and Pd–Cl2 bond lengths, 2.361(1) and 2.408(1) Å, respectively, demonstrates that the *trans* influence of the silylene is much greater than that of the phosphane. A strong *trans* influence has been previously observed for monodentate amidinatosilylenes.¹⁶

Regarding reactions of **1_{Si}** and **1_{Ge}** with Group 10 metals in zero oxidation state, both HT-phosphanes reacted at room temperature with [Ni(cod)₂], displacing one of the cod ligands,

to give complexes of the type $[\text{Ni}\{\kappa^2E,P\text{-}E(\text{tBu}_2\text{bzam})\text{pyrmp}^i\text{Bu}_2\}(\text{cod})]$ ($\mathbf{3}_{\text{Ni-E}}$; E = Si, Ge) as single reaction products (Scheme 3).





Fig. 3 XRD molecular structure of complex $\mathbf{2}_{\text{Pd-Si}}$ (30% displacement ellipsoids, H atoms omitted for clarity). Selected bond lengths (Å) and angles ($^\circ$): Pd1–C1 2.361(1), Pd1–C12 2.408(1), Pd1–P1 2.2654(9), Pd1–Si1 2.228(1), Si1–N1 1.807(3), Si1–N2 1.818(3), Si1–N3 1.760(3), C4–N1 1.488(5), C5–N1 1.338(5), C5–C6 1.483(5), C5–N2 1.335(5), C12–N2 1.494(5), C19–N3 1.400(5), C19–C20 1.489(5), C20–P1 1.837(4), C21–P1 1.885(4), C25–P1 1.899(4), Cl1–Pd1–Cl2 91.20(4), Cl1–Pd1–P1 170.93(4), Cl1–Pd1–Si1 83.42(4), Cl2–Pd1–P1 93.95(4), Cl2–Pd1–Si1 161.67(4), P1–Pd1–Si1 93.86(3), N1–Si1–N2 102.9(2), N1–Si1–N3 106.8(2), N2–Si1–N3 102.9(2), N1–Si1–Pd1 125.8(1), N2–Si1–Pd1 119.2(1), N3–Si1–Pd1 119.1(1), N1–C5–N2 106.7(3).



Scheme 3 Reactions of $\mathbf{1}_{\text{Si}}$ and $\mathbf{1}_{\text{Ge}}$ with Group 10 metal(0) complexes.




Fig. 4 XRD molecular structure of complex $\mathbf{3}_{\text{Ni-Ge}}$ (40% displacement ellipsoids, H atoms omitted for clarity). Ni1–C29 2.126(4), Ni1–C30 2.131(4), Ni1–C33 2.119(4), Ni1–C34 2.145(4), Ni1–P1 2.222(1), Ni1–Ge1 2.2439(7), Ge1–N1 2.013(3), Ge1–N2 2.004(3), Ge1–N3 1.921(3), C4–N1 1.474(5), C5–N1 1.330(5), C5–C6 1.495(5), C5–N2 1.344(4), C12–N2 1.492(5), C19–N3 1.383(4), C19–C20 1.491(5), C20–P1 1.869(4), C21–P1 1.899(4), C25–P1 1.900(4), {C29–C30_{centroid}–Ni1–{C33–C34_{centroid}}} 90.32(2), {C29–C30_{centroid}–Ni1–Ge1} 114.37(3), {C33–C34_{centroid}–Ni1–Ge1} 113.79(2), {C29–

C30_{centroid}–Ni1–P1 122.28(3), {C33–C34_{centroid}–Ni1–P1 122.72(4), Ge1–Ni1–P1 95.00(4), N1–Ge1–N2 65.50(1), N1–Ge1–N3 97.7(1), N2–Ge1–N3 97.2(1), N1–Ge1–Ni1 130.0(1), N2–Ge1–Ni1 131.9(1), N3–Ge1–Ni1 120.0(1), N1–C5–N2 108.7(3).

NMR spectroscopy confirmed that $\mathbf{3}_{\text{Ni-Si}}$ and $\mathbf{3}_{\text{Ni-Ge}}$ contain one cod and one HT-phosphane ligand and that they behave as C_5 symmetric molecules in solution. The molecular structure of $\mathbf{3}_{\text{Ni-Ge}}$ was determined by XRD. Fig. 4 shows that the Ni atom is in a distorted tetrahedral coordination environment. The chelating bite angle of the HT-phosphane ligand is a quite narrow for a tetrahedral complex, Ge1–Ni1–P1 95.00(4) $^\circ$ (ideal angle = 109.5 $^\circ$), being only 1.2 $^\circ$ wider than the P1–Pd1–Si1 angle of $\mathbf{2}_{\text{Pd-Si}}$, 93.86(3) $^\circ$ (ideal angle = 90.0 $^\circ$).

Both $\mathbf{1}_{\text{Si}}$ and $\mathbf{1}_{\text{Ge}}$ reacted cleanly at room temperature with $[\text{M}(\text{PPh}_3)_4]$ (M = Pd, Pt) to give the three-coordinated complexes $[\text{M}\{\kappa^2E,P\text{-}E(\text{tBu}_2\text{bzam})\text{pyrmp}^i\text{Bu}_2\}(\text{PPh}_3)]$ ($\mathbf{4}_{\text{M-E}}$; M = Pd, Pt; E = Si, Ge) (Scheme 3). The use of an excess of HT-phosphane reagent had no effect on the outcome of these reactions. Probably the HT-phosphanes $\mathbf{1}_{\text{E}}$ ate too large to allow an appropriate fitting of two of them around the metal atom. The isolation of pure products from these reactions required several recrystallizations (in order to get rid of the accompanying PPh_3) and this reduced the final yields to 30–40%. The similar solubility of PPh_3 and $\mathbf{4}_{\text{Pd-Ge}}$ avoided the isolation of a PPh_3 -free sample of the metal complex.

NMR spectroscopy confirmed that, in solution, the compounds of type $\mathbf{4}_{\text{M-E}}$ behave as symmetric molecules (C_5) and that they contain one PPh_3 per HT-phosphane ligand; for example, the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of $\mathbf{4}_{\text{Pt-Si}}$ consists of two doublets ($J_{\text{P-P}} = 118.4$ Hz) with ^{195}Pt satellites ($J_{\text{P-Pt}} = 3521.7$ and 4287.0 Hz). An XRD study on a crystal of $\mathbf{4}_{\text{Pd-Si}}$ (Fig. 5) established that the Pd atom is in a distorted triangular coordination environment. Again, the chelating bite angle of the HT-phosphane ligand, Si1–Pd1–P1 98.77(2) $^\circ$, is narrower than expected (ideal angle for a planar triangular complex: 120.0 $^\circ$).




Fig. 5 XRD molecular structure of complex $\mathbf{4}_{\text{Pd-Si}}$ (30% displacement ellipsoids, H atoms omitted for clarity). Selected bond lengths (Å) and angles ($^\circ$): Pd1–P1 2.3108(4), Pd1–P2 2.2881(4), Pd1–Si1 2.2622(4), Si1–N1 1.862(1), Si1–N2 1.858(1), Si1–N3 1.804(1), C4–N1 1.476(2), C5–N1 1.337(2), C5–C6 1.489(2), C5–N2 1.338(2), C12–N2 1.483(2), C19–N3 1.393(2), C19–C20 1.499(2), C20–P1 1.866(2), C21–P1 1.885(2), C25–P1 1.895(2), P1–Pd1–P2 131.06(2), P1–Pd1–Si1 98.77(2), P2–Pd1–Si1 130.18(2), N1–Si1–N2 70.06(6), N1–Si1–N3 101.21(6), N2–Si1–N3 99.70(6), N1–Si1–Pd1 130.31(5), N2–Si1–Pd1 128.86(5), N3–Si1–Pd1 116.06(5), N1–C5–N2 105.9(1).

Conclusions

In summary, this work describes the syntheses of the first silylene-phosphane (**1_{Si}**) and the second¹³ germylene-phosphane (**1_{Ge}**) whose coordination abilities have been proven by reactions with transition-metal complexes. The reactivity studies have indicated that both **1_{Si}** and **1_{Ge}** are excellent chelating ligands for square-planar and octahedral complexes (their bite angle is close to 90°) and that they can also participate as chelating ligands in tetrahedral and triangular-planar complexes, leading to slightly distorted coordination geometries. No doubt, the offering of **1_{Si}** and **1_{Ge}** to the chemical community will open up new pathways toward TM complexes for which interesting structural, bonding and catalytic properties can be anticipated.

Experimental section

General data

All reactions and product manipulations were carried out under argon in a drybox or using Schlenk-vacuum line techniques. Solvents were dried over appropriate desiccating reagents and were distilled under argon before use. The compounds HpyrmP^tBu₂,^{12b} E(^tBu₂bzam)Cl (E = Si,¹⁴ Ge;¹⁵) [PdCl₂(MeCN)₂],¹⁶ [PtCl₂(cod)]¹⁷ and [Pt(PPh₃)₄]¹⁸ were prepared following published procedures. All remaining reagents were purchased from commercial sources and were stored under argon in a drybox. All reaction products were vacuum-dried for several hours prior to being weighted and analyzed. NMR spectra were run on Bruker NAV-400 and AC-300 instruments, using as standards the residual protic solvent resonance for ¹H [δ(CHCl₃) 7.26 ppm; δ(C₆HD₅) 7.16 ppm; δ(CHDCl₂) 5.32 ppm], the solvent resonance for ¹³C [δ(CDCl₃) 77.16 ppm; δ(C₆D₆) 128.10 ppm; δ(CD₂Cl₂) 54.00 ppm], external 85% H₃PO₄ for ³¹P (δ 0.00 ppm) and external SiMe₄ in CDCl₃ for ²⁹Si (δ = 0.00 ppm). Microanalyses were obtained with a FlashEA112 (Thermo-Finnigan) microanalyzer. High-resolution mass spectra (HRMS) were obtained with a Bruker Impact II mass spectrometer operating in the ESI-Q-ToF positive mode; data given refer to the most abundant isotopomer of the species with the greatest mass. CHN microanalysis and/or mass spectra were not obtained for the products that were very air- and/or moisture-sensitive.

Synthetic procedures and characterization data

Si(^tBu₂bzam)pyrmP^tBu₂ (1_{Si}**):** LiⁿBu (0.75 mL, 1.6 M in hexanes, 1.2 mmol) was dropwise added to a cold (−78 °C) solution of HpyrmP^tBu₂ (2.5 mL, 0.4 M in toluene, 1.0 mmol) in diethyl ether (2 mL). The resulting bright yellow solution was allowed to reach the room temperature, was stirred for 2 h and was added to a suspension of Si(^tBu₂bzam)Cl (0.295 g, 1.0 mmol) in diethyl ether (2 mL). The resulting orange suspension was stirred for 18 h. Solvents were removed under vacuum and the residue was extracted into hexane (4 x 5 mL). A white solid was separated by decantation from a clear yellow solution that was evaporated to dryness to give **1_{Si}** as a pale yellow solid (0.436 g, 90 %). ¹H NMR (C₆D₆, 300.1 MHz, 298 K): δ 7.03–6.92 (m, 6 H, CHs of Ph and 1 CH of pyrrole), 6.58 (s, 1 H, 1 CH of

pyrrole), 6.48 (s, 1 H, 1 CH of pyrrole), 3.63 (s, 2 H, 1 CH₂ of PCH₂), 1.26 (d, *J*_{H-P} = 10.2 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.04 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 164.3 (s, NCN), 137.8 (d, *J*_{C-P} = 15.4 Hz, pyrrole C²), 133.8 (s, C_{ipso} of Ph), 130.1–128.0 (m, CHs of Ph), 120.5 (s, pyrrole CH), 110.5 (d, *J*_{C-P} = 9.1 Hz, pyrrole CH), 109.8 (s, 1 pyrrole CH), 53.2 (s, 2 C of 2 N^tBu), 32.0 (d, *J*_{C-P} = 25.8 Hz, 2 C of P^tBu₂), 31.5 (s, 6 CH₃ of 2 N^tBu), 30.4 (d, *J*_{C-P} = 13.1 Hz, 6 CH₃ of P^tBu₂), 21.6 (d, *J*_{C-P} = 24.1 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 121.5 MHz, 298 K): δ 28.2 (s) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ −18.8 (d, *J*_{Si-P} = 14.2 Hz) ppm.

Ge(^tBu₂bzam)pyrmP^tBu₂ (1_{Ge}**):** This compound was prepared as a pale orange solid following the procedure described above for **1_{Si}**. Reagents: HpyrmP^tBu₂ (2.5 mL, 0.4 M in toluene, 1.0 mmol), BuLi (0.75 mL, 1.6 M in hexanes, 1.2 mmol), Ge(^tBu₂bzam)Cl (0.340 g, 1.0 mmol). Yield: 0.286 g, 54 %. ¹H NMR (C₆D₆, 300.1 MHz, 298 K): δ 7.25 (s, 1 H, pyrrole CH), 7.09–6.92 (m, 5 H, CHs of Ph), 6.58 (s, 1 H, pyrrole CH), 6.47 (s, 1 H, pyrrole CH), 3.46 (s, 2 H, PCH₂), 1.24 (d, *J*_{H-P} = 9.0 Hz, 18 H, 6 CH₃ de P^tBu₂), 0.99 (s, 18 H, 6 CH₃ de 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 169.8 (s, NCN), 136.0 (d, *J*_{C-P} = 11.1 Hz, pyrrole C²), 135.7 (C_{ipso} of Ph), 129.8–127.7 (m, CHs of Ph), 122.2 (s, pyrrole CH), 109.4 (d, *J*_{C-P} = 5.0 Hz, pyrrole CH), 108.5 (s, pyrrole CH), 53.1 (s, 2 C of 2 N^tBu), 32.1 (d, *J*_{C-P} = 26.2 Hz, 2 C of P^tBu₂), 31.9 (s, 6 CH₃ of 2 N^tBu), 30.2 (d, *J*_{C-P} = 12.1 Hz, 6 CH₃ de P^tBu₂), 22.2 (d, *J*_{C-P} = 24.2 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 121.5 MHz, 298 K): δ 28.3 (s) ppm.

[NiCl₂{κ²P,Si-Si(^tBu₂bzam)pyrmP^tBu₂}] (2_{Ni-Si}**):** Silylene **1_{Si}** (0.030 g, 0.06 mmol) was added to a solution of [NiCl₂(dme)] (0.013 g, 0.06 mmol) in toluene (3 mL). After stirring for 12 h, the solvent was removed under reduced pressure and the residue was washed with hexane (2 mL) to give **2_{Ni-Si}** as a pale brown solid (0.025 g, 69 %). Anal. (%) Calcd. for C₂₈H₄₆Cl₂N₃NiPSi (M_w = 613.34 amu): C, 54.83; H, 7.56; N, 6.85; found: C, 54.92; H, 7.65; N, 6.82. (+)-ESI HRMS: *m/z* 576.1960, calcd. for C₂₈H₄₆ClN₃NiPSi [M – Cl]⁺: 576.2241. ¹H NMR (C₆D₆, 400.5 MHz, 298 K): δ 7.57 (d, *J*_{H-P} = 9.0 Hz, 1 CH of Ph), 7.09 (s, 1 H, pyrrole CH), 6.93–6.79 (m, 4 H, 4 CH of Ph), 6.38 (s, 1 H, pyrrole CH), 6.04 (s, 1 H, pyrrole CH), 2.93 (d, *J*_{H-P} = 9.0 Hz, 2 H, CH₂P), 1.52 (d, *J*_{H-P} = 15.0 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.18 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 176.6 (s, NCN), 132.7 (s, pyrrole C²), 130.9–127.6 (m, CHs + C_{ipso} of Ph), 122.2 (s, pyrrole CH), 111.7 (d, *J*_{C-P} = 5.0 Hz, pyrrole CH), 111.2 (s, pyrrole CH), 54.9 (s, 2 C of 2 N^tBu), 37.2 (d, *J*_{C-P} = 14.1 Hz, 2 C of P^tBu₂), 31.0 (s, 6 CH₃ of 2 N^tBu), 30.5 (s, 6 Me of P^tBu₂), 22.3 (d, *J*_{C-P} = 21.8 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 49.1 (s) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ −27.7 (d, *J*_{Si-P} = 81.3 Hz) ppm.

[NiCl₂{κ²P,Ge-Ge(^tBu₂bzam)pyrmP^tBu₂}] (2_{Ni-Ge}**):** Germylene **1_{Ge}** (0.032 g, 0.06 mmol) was added to a solution of [NiCl₂(dme)] (0.013 g, 0.06 mmol) in toluene (3 mL). After stirring for 12 h, the solvent was removed under reduced pressure and the residue was washed with hexane (2 mL) to give **2_{Ni-Ge}** as a purple solid (0.025 g, 63 %). Anal. (%) Calcd. for C₂₈H₄₆Cl₂GeN₃NiP (M_w = 657.88 amu): C, 51.12; H, 7.05; N, 6.39; found: C, 51.20; H, 7.13; N, 6.37. (+)-ESI HRMS: *m/z* 680.1135, calcd. for C₂₈H₄₆Cl₂GeN₃NaNiP [M + Na]⁺: 680.1269. ¹H NMR (CD₂Cl₂, 400.5 MHz, 298 K): δ 7.59–7.49 (m, 5 H, 5 CH of Ph), 7.05 (s, 1 H, pyrrole CH), 6.20 (s, 1 H, pyrrole CH), 6.03 (s, 1 H, pyrrole

CH), 3.33 (s, br, 2 H, CH₂P), 1.57 (s, br, 18 H, 6 CH₃ of P^tBu₂), 1.23 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (CD₂Cl₂, 100.6 MHz, 298 K): δ 175.9 (s, NCN), 132.4 (s, pyrrole C²), 132.1 (s, C_{ipso} of Ph), 131.3–128.6 (m, CHs of Ph), 123.1 (s, pyrrole CH), 111.4 (s, pyrrole CH), 110.3 (s, pyrrole CH), 55.5 (s, 2 C of 2 N^tBu), 39.3 (s, 2 C of P^tBu₂), 31.8 (s, 6 CH₃ of N^tBu), 30.0 (s, 6 CH₃ of P^tBu₂), 22.1 (d, J_{C-P} = 22.1 Hz, PCH₂) ppm. ³¹P{¹H} NMR (CD₂Cl₂, 162.1 MHz, 298 K): δ 63.4 (s) ppm.

[PdCl₂{κ²P,Si-Si(^tBu₂bzam)pyrmP^tBu₂}] (2_{Pd-Si}): Silylene **1_{Si}** (0.030 g, 0.06 mmol) was added to a solution of [PdCl₂(MeCN)₂] (0.016 g, 0.06 mmol) in toluene (3 mL). After stirring for 12 h, the solvent was removed under reduced pressure and the residue was washed with hexane (2 mL) to give **2_{Pd-Si}** as a pale brown solid (0.034 g, 86 %). Anal. (%) Calcd. for C₂₈H₄₆Cl₂N₃PPdSi (M_w = 661.07 amu): C, 50.87; H, 7.01; N, 6.36; found: C, 50.95; H, 7.08; N, 6.29. (+)-ESI HRMS: *m/z* 626.1644, calcd. for C₂₈H₄₆ClN₃PPdSi [M – Cl]⁺: 626.1921. ¹H NMR (C₆D₆, 400.5 MHz, 298 K): δ 7.95 (d, J_{H-P} = 7.8 Hz, 1 CH of Ph), 7.13 (dt, J_{H-H} = 2.9 Hz; 1.4 Hz, 1 H, pyrrole CH), 7.03–6.89 (m, 4 H, 4 CH of Ph), 6.39 (q, J_{H-H} = 2.9 Hz, 1 H, pyrrole CH), 6.08 (m, 1 H, pyrrole CH), 3.06 (d, J_{H-P} = 9.4 Hz, 2 H, CH₂P), 1.43 (d, J_{H-P} = 14.4 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.10 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 177.7 (s, NCN), 133.4 (s, pyrrole C²), 131.1–127.2 (m, CHs of Ph), 123.3 (s, pyrrole CH), 112.6 (d, J_{C-P} = 5.4 Hz, pyrrole CH), 111.2 (s, pyrrole CH), 55.0 (s, 2 C of 2 N^tBu), 38.1 (d, J_{C-P} = 17.5 Hz, 2 C of P^tBu₂), 31.0 (s, 6 CH₃ of 2 N^tBu), 30.3 (s, 6 CH₃ of P^tBu₂), 22.3 (d, J_{C-P} = 22.1 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 54.9 (s) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ –0.46 (d, J_{Si-P} = 11.7 Hz) ppm.

[PdCl₂{κ²P,Ge-Ge(^tBu₂bzam)pyrmP^tBu₂}] (2_{Pd-Ge}): Germylene **1_{Ge}** (0.032 g, 0.06 mmol) was added to a solution of [PdCl₂(MeCN)₂] (0.016 g, 0.06 mmol) in toluene (3 mL). After stirring for 12 h a solid was precipitated. The solution was removed and the solid was dried under vacuum to give **2_{Pd-Ge}** as a yellow solid (0.026 g, 62 %). Anal. (%) Calcd. for C₂₈H₄₆Cl₂GeN₃PPd (M_w = 705.62 amu): C, 47.66; H, 6.57; N, 5.96; found: C, 47.74; H, 6.65; N, 5.89. (+)-ESI HRMS: *m/z* 728.0630, calcd. for C₂₈H₄₆Cl₂GeN₃NaPPd [M + Na]⁺: 728.0960. ¹H NMR (CD₂Cl₂, 400.5 MHz, 298 K): δ 7.78 (m, 1 CH of Ph), 7.59–7.49 (m, 4 H, 4 CH of Ph), 7.13 (td, J_{H-H} = 3.0, 1.7 Hz, 1 H, pyrrole CH), 6.26 (td, J_{H-H} = 3.0, 1.7 Hz, 1 H, pyrrole CH), 6.08 (m, 1 H, pyrrole CH), 3.33 (d, J_{H-P} = 9.4 Hz, 2 H, CH₂P), 1.47 (d, J_{H-P} = 14.7 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.17 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (CD₂Cl₂, 100.6 MHz, 298 K): δ 176.6 (s, NCN), 131.8 (s, pyrrole C²), 131.5 (s, CH of Ph), 131.2 (s, C_{ipso} of Ph), 130.7 (s, CH of Ph), 129.0 (s, CH of Ph), 128.7 (s, CH of Ph), 128.65 (s, CH of Ph), 123.8 (s, pyrrole CH), 112.4 (d, J_{C-P} = 5.1 Hz, pyrrole CH), 110.6 (s, pyrrole CH), 55.8 (s, 2 C of 2 N^tBu), 39.23 (d, J_{C-P} = 16.4 Hz, 2 C of P^tBu₂), 31.5 (s, 6 CH₃ of 2 N^tBu), 30.4 (s, 6 CH₃ of P^tBu₂), 22.7 (d, J_{C-P} = 22.1 Hz, PCH₂) ppm. ³¹P{¹H} NMR (CD₂Cl₂, 162.1 MHz, 298 K): δ 56.2 (s) ppm.

[PtCl₂{κ²P,Si-Si(^tBu₂bzam)pyrmP^tBu₂}] (2_{Pt-Si}): Silylene **1_{Si}** (0.030 g, 0.06 mmol) was added to a solution of [PtCl₂(cod)] (0.022 g, 0.06 mmol) in toluene (3 mL). After stirring for 12 h, the solvent was removed under reduced pressure and the residue was washed with hexane (2 mL) to give **2_{Pt-Si}** as a brown solid (0.042 g, 90 %). Anal. (%) Calcd. for C₂₈H₄₆Cl₂N₃PPTSi (M_w = 749.73 amu): C, 44.86; H, 6.18; N, 5.61; found: C, 44.93; H, 6.23; N 5.57. (+)-ESI HRMS: *m/z* 715.2221, calcd. for C₂₈H₄₇ClN₃PPTSi [M – Cl + H]⁺: 715.2611. ¹H NMR (C₆D₆, 400.5 MHz, 298 K): δ 7.92 (m, 1 CH of Ph), 7.12 (dt, J_{H-H} = 2.9, 1.5 Hz, 1 H, pyrrole CH), 7.03–6.90 (m, 4 H, 4 CH of Ph), 6.38 (td, J_{H-H} = 2.9,

1.5 Hz, 1 H, pyrrole CH), 6.06 (m, 1 H, pyrrole CH), 3.28 (dd, J_{H-P} = 10.3 Hz, J_{H-H} = 1.0 Hz, 2 H, CH₂P), 1.42 (d, J_{H-P} = 14.2 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.18 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 178.2 (s, NCN), 133.4 (d, J_{C-P} = 5.4 Hz, pyrrole C²), 131.2–127.5 (m, C_{ipso} + CHs of Ph), 123.2 (s, pyrrole CH), 112.8 (d, J_{C-P} = 6.1 Hz, pyrrole CH), 111.3 (s, pyrrole CH), 54.9 (s, 2 C of 2 N^tBu), 37.7 (d, J_{C-P} = 26.2 Hz, 2 C of P^tBu₂), 30.8 (s, 6 CH₃ of 2 N^tBu), 30.3 (s, 6 CH₃ of P^tBu₂), 22.4 (d, J_{C-P} = 31.3 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 32.8 (s, sat, J_{P-Pt} = 3701.2 Hz) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ –16.8 (d, sat, J_{Si-P} = 13.1 Hz, J_{Si-Pt} = 2298.3 Hz) ppm.

[PtCl₂{κ²P,Ge-Ge(^tBu₂bzam)pyrmP^tBu₂}] (2_{Pt-Ge}): Germylene **1_{Ge}** (0.032 g, 0.06 mmol) was added to a solution of [PtCl₂(cod)] (0.022 g, 0.06 mmol) in toluene (3 mL). After stirring for 12 h, the solvent was removed under reduced pressure and the residue was washed with hexane (2 mL) to give **2_{Pt-Ge}** as a pale pink solid (0.038 g, 90 %). Anal. (%) Calcd. for C₂₈H₄₆Cl₂GeN₃PPT (M_w = 794.28 amu): C, 42.34; H, 5.84; N, 5.29; found: C, 42.46; H, 5.93; N 5.24. (+)-ESI HRMS: *m/z* 816.1184, calcd. for C₂₈H₄₆Cl₂GeN₃NaPPT [M + Na]⁺: 816.1562. ¹H NMR (CD₂Cl₂, 400.5 MHz, 298 K): δ 7.77 (m, 1 CH of Ph), 7.65–7.34 (m, 4 H, 4 CH of Ph), 7.10 (m, 1 H, pyrrole CH), 6.21 (m, 1 H, pyrrole CH), 6.09 (m, 1 H, pyrrole CH), 3.53 (d, J_{H-P} = 10.0 Hz, 2 H, CH₂P), 1.45 (d, J_{H-P} = 14.4 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.17 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (CD₂Cl₂, 100.6 MHz, 298 K): δ 176.9 (s, NCN), 132.0 (d, J_{C-P} = 6.0 Hz, pyrrole C²), 131.5 (s, CH of Ph), 131.3 (s, C_{ipso} of Ph), 130.8 (s, CH of Ph), 129.0 (s, CH of Ph), 128.8 (s, CH of Ph), 128.6 (s, CH of Ph), 123.7 (s, pyrrole CH), 112.8 (d, J_{C-P} = 6.0 Hz, pyrrole CH), 110.3 (s, pyrrole CH), 55.7 (s, 2 C of 2 N^tBu), 38.4 (d, J_{C-P} = 25.2 Hz, 2 C of P^tBu₂), 31.4 (s, 6 CH₃ of 2 N^tBu), 30.5 (d, J_{C-P} = 2.2 Hz, 6 CH₃ of P^tBu₂), 22.5 (d, J_{C-P} = 30.8 Hz, PCH₂) ppm. ³¹P{¹H} NMR (CD₂Cl₂, 162.1 MHz, 298 K): δ 28.80 (s, sat, J_{P-Pt} = 3553.2 Hz) ppm.

[Ni{κ²P,Si-Si(^tBu₂bzam)pyrmP^tBu₂}(cod)] (3_{Ni-Si}): Silylene **1_{Si}** (0.030 g, 0.06 mmol) was added to a solution of [Ni(cod)₂] (0.017 g, 0.06 mmol) in toluene (3 mL). After stirring for 4 h, the solvent was removed under reduced pressure and the residue was washed with hexane (1 mL) to give **3_{Ni-Si}** as an orange solid (0.014 g, 40 %). Anal. (%) Calcd. for C₃₆H₅₈N₃NiPSi (M_w = 650.62 amu): C, 66.46; H, 8.98; N, 6.46; found: C, 66.53; H, 9.04; N, 6.43. (+)-ESI HRMS: *m/z* 649.3510, calcd. for C₃₆H₅₈N₃NiPSi [M]⁺: 649.3491. ¹H NMR (C₆D₆, 300.1 MHz, 298 K): δ 7.39 (s, 1 H, CH of Ph), 6.98 (m, 5 H, 4 CH of Ph and 1 pyrrole CH), 6.58 (s, 1 H, pyrrole CH), 6.29 (s, 1 H, pyrrole CH), 4.70 (m, 2 H, 2 CH of cod), 4.48 (m, 2 H, 2 CH of cod), 3.22 (s, 2 H, CH₂P), 2.87–2.10 (m, 8 H, 4 CH₂ of cod), 1.29 (d, J_{H-P} = 15.0 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.08 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 171.3 (s, NCN), 135.6 (s, pyrrole C²), 132.7 (s, C_{ipso} of Ph), 130.4–127.6 (m, 5 CHs of Ph), 121.2 (s, pyrrole CH), 110.7 (d, J_{C-P} = 5.3 Hz, pyrrole CH), 108.7 (s, pyrrole CH), 80.1 (d, J_{C-P} = 8.0 Hz, 2 CH of cod), 78.2 (d, J_{C-P} = 4.1 Hz, 2 CH of cod), 53.3 (s, 2 C of 2 N^tBu), 35.6 (s, 2 C of P^tBu₂), 32.9 (d, J_{C-P} = 6.0 Hz, 2 CH₂ of cod), 32.5 (s, 2 CH₂ of cod), 30.9 (s, 6 CH₃ of 2 N^tBu), 30.4 (s, 6 CH₃ of P^tBu₂), 26.5 (d, J_{C-P} = 14.8 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 121.5 MHz, 298 K): δ 55.0 (s) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ 56.8 (d, J_{Si-P} = 26.3 Hz) ppm.

[Ni{κ²P,Ge-Ge(^tBu₂bzam)pyrmP^tBu₂}(cod)] (3_{Ni-Ge}): Germylene **1_{Ge}** (0.032 g, 0.06 mmol) was added to a solution of [Ni(cod)₂] (0.017 g, 0.06 mmol) in toluene (3 mL). After stirring for 4 h, the solvent was removed under reduced pressure and the residue was washed with

hexane (1 mL) to give **5_{Ge}** as a pale red solid (0.025 g, 59 %). Anal. (%) Calcd. for C₃₆H₅₈GeN₃NiP (M_w = 695.16 amu): C, 62.20; H, 8.41; N, 6.05; found: C, 62.38; H, 8.48; N 6.01. (+)-ESI HRMS: *m/z* 696.2691, calcd. for C₃₆H₅₉GeN₃NiP [M + H]⁺: 696.3011. ¹H NMR (C₆D₆, 400.5 MHz, 298 K): δ 7.38–6.94 (m, 6 H, 5 CH of Ph + pyrrole CH), 6.64 (s, 1 H, pyrrole CH), 6.32 (s, 1 H, pyrrole CH), 4.77 (m, 2 H, 2 CH of cod), 4.63 (m, 2 H, 2 CH of cod), 3.25 (s, 2 H, CH₂P), 2.72–2.40 (m, 8 H, 4 CH₂ of cod), 1.27 (d, *J*_{H-P} = 12.0 Hz, 6 CH₃ of P^tBu₂), 1.05 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 172.5 (s, NCN), 134.6 (s, pyrrole C²), 134.4 (s, C_{ipso} of Ph), 130.3–127.6 (m, CHs of Ph), 121.7 (s, pyrrole CH), 110.3 (d, *J*_{C-P} = 4.2 Hz, pyrrole CH), 108.0 (s, pyrrole CH), 80.3 (d, *J*_{C-P} = 7.2 Hz, 2 CH of cod), 80.0 (d, *J*_{C-P} = 3.4 Hz, 2 CH of cod), 53.3 (s, 2 C of 2 N^tBu), 35.2 (s, 2 C of P^tBu₂), 33.0 (s, 2 CH₂ of cod), 32.8 (d, *J*_{C-P} = 5.2 Hz, 2 CH₂ of cod), 31.3 (s, 6 CH₃ of 2 N^tBu), 30.1 (s, 3 CH₃ of P^tBu₂), 29.2 (s, 3 CH₃ of P^tBu), 26.4 (d, *J*_{C-P} = 12.3 Hz, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 51.8 (s) ppm.

[Pd{κ²P,Si-Si(^tBu₂bzam)pyrmP^tBu₂}(PPh₃)] (4_{Pd-Si}): Silylene **1_{Si}** (0.030 g, 0.06 mmol) was added to a solution of [Pd(PPh₃)₄] (0.069 g, 0.06 mmol) in toluene (3 mL). After stirring for 4 h, the solvent was removed under reduced pressure and the residue was recrystallized from toluene to give **4_{Pd-Si}** as a pale orange solid (0.014 g, 40 %). Anal. (%) Calcd. for C₄₆H₆₁N₃P₂PdSi (M_w = 852.45 amu): C, 64.81; H, 7.21; N, 4.93; found: C, 64.87; H, 7.29; N 4.89. (+)-ESI HRMS: *m/z* 923.2355, calcd. for C₄₉H₆₇N₄OP₂PdSi [M + CH₃CN + CH₃OH – H]⁺: 923.3611. ¹H NMR (C₆D₆, 300.1 MHz, 298 K): δ 8.00 (t, *J*_{H-P} = 8.3 Hz, 6 H, 6 CH of 3 Ph), 7.39–6.96 (m, 15 H, 14 CH of 4 Ph + 1 pyrrole CH), 6.60 (s, 1 H, pyrrole CH), 6.39 (s, 1 H, pyrrole CH), 3.52 (s, 2 H, CH₂P), 1.25 (d, *J*_{H-P} = 11.6 Hz, 18 H, 6 CH₃ of P^tBu₂), 1.05 (s, 18 H, 6 CH₃ of 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 165.3 (s, NCN), 141.9 (m, pyrrole C²), 131.0–127.8 (m, C_{ipso}s + CHs of Phs), 122.7 (s, pyrrole CH), 111.8 (d, *J*_{C-P} = 5.5 Hz, pyrrole CH), 108.5 (s, pyrrole CH), 53.4 (s, 2 C of 2 N^tBu), 34.7 (d, *J*_{C-P} = 9.1 Hz, 2 C of P^tBu₂), 31.6 (6 CH₃ of 2 N^tBu), 30.2 (d, *J*_{C-P} = 10.9 Hz, 6 CH₃ of P^tBu₂), 26.4 (s, PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 49.3 (d, *J*_{P-P} = 59.2 Hz), 35.2 (d, *J*_{P-P} = 59.2 Hz) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ 19.3 (dd, *J*_{Si-P} = 69.7, 42.9 Hz) ppm.

Attempted synthesis of [Pd{κ²P,Ge-Ge(^tBu₂bzam)pyrmP^tBu₂}(PPh₃)] (4_{Pd-Ge}): Germylene **1_{Ge}** (0.032 g, 0.06 mmol) was added to a solution of [Pd(PPh₃)₄] (0.069 g, 0.06 mmol) in toluene (3 mL). After stirring for 4 h, the solvent was removed under reduced pressure and the residue, which contained a mixture of **4_{Pd-Ge}** and PPh₃ (NMR analysis), was recrystallized several times from hexane, but **4_{Pd-Ge}** could not be obtained free of PPh₃. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 54.0 (s) ppm.

[Pt{κ²P,Si-Si(^tBu₂bzam)pyrmP^tBu₂}(PPh₃)] (4_{Pt-Si}): Silylene **1_{Si}** (0.030 g, 0.06 mmol) was added to a solution of [Pt(PPh₃)₄] (0.075 g, 0.06 mmol) in toluene (3 mL). After stirring for 4 h, the solvent was removed under reduced pressure and the residue was recrystallized several times from hexane to give **4_{Pt-Si}** as an orange solid (0.018 g, 32 %). Anal. (%) Calcd. for C₄₆H₆₁N₃P₂PtSi (M_w = 941.12 amu): C, 58.71; H, 6.53; N, 4.46; found C, 58.76; H, 6.60; N, 4.38. (+)-ESI HRMS: *m/z* 941.3685, calcd. for C₄₆H₆₂N₃P₂PtSi [M + H]⁺: 941.3839. ¹H NMR (C₆D₆, 300.1 MHz, 298 K): δ 8.01 (t, *J*_{H-P} = 8.9 Hz, 6 H of 3 Ph), 7.39–6.98 (m, 15 H, 14 CH of 4 Ph + pyrrole CH), 6.60 (s, 1 H, pyrrole CH), 6.37 (s, 1

H, pyrrole CH), 3.73 (s, 2 H, CH₂ de PCH₂), 1.27 (d, *J*_{H-P} = 12.0 Hz, 18 H, 6 CH₃ de P^tBu₂), 1.04 (s, 18 H, 6 CH₃ de 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 167.6 (s, NCN), 144.2 (m, 3 C_{ipso} of PPh₃), 136.9 (s, pyrrole C²), 133.2 (s, C_{ipso} of Ph), 130.5–127.8 (m, CHs of Phs), 123.2 (s, pyrrole CH), 112.2 (d, *J*_{C-P} = 6.0 Hz, pyrrole CH), 108.4 (s, pyrrole CH), 53.5 (s, 2 C de 2 N^tBu), 37.5 (s, sat, *J*_{C-Pt} = 40.3 Hz, 2 C de P^tBu₂), 31.6 (s, 6 CH₃ de 2 N^tBu), 30.2 (d, *J*_{C-P} = 8.5 Hz, 6 CH₃ de P^tBu₂), 26.4 (d, sat, *J*_{C-Pt} = 16.7 Hz, *J*_{C-P} = 11.0 Hz, CH₂ de PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 88.1 (d, sat, *J*_{P-P} = 118.4 Hz, *J*_{P-Pt} = 3521.7), 66.2 (d, sat, *J*_{P-P} = 118.4 Hz, *J*_{P-Pt} = 4287.0 Hz) ppm. ²⁹Si{¹H} NMR (C₆D₆, 79.5 MHz, 298 K): δ 70.3 (dd, *J*_{Si-P} = 105.7, 65.9 Hz) ppm.

[Pt{κ²P,Ge-Ge(^tBu₂bzam)pyrmP^tBu₂}(PPh₃)] (4_{Pt-Ge}): Germylene **1_{Ge}** (0.032 g, 0.06 mmol) was added to a solution of [Pt(PPh₃)₄] (0.075 g, 0.06 mmol) in toluene (3 mL). After stirring for 4 h, the solvent was removed under reduced pressure and the residue was recrystallized several times from hexane to give **4_{Pt-Ge}** as an orange solid (0.021 g, 36 %). Anal. (%) Calcd. for C₄₆H₆₁GeN₃P₂Pt (M_w = 985.66 amu): C, 56.05; H, 6.24; N, 4.26; found C, 56.17; H, 6.33; N 4.21. (+)-ESI HRMS: *m/z* 963.3692, calcd. for C₄₁H₆₀GeN₃NaOP₂Pt [M – Ph + MeOH + Na]⁺: 963.2979. ¹H NMR (C₆D₆, 300.1 MHz, 298 K): δ 8.01 (t, *J*_{H-P} = 8.3 Hz, 6 H of 3 Ph), 7.39–6.96 (m, 15 H, 14 CH of 4 Ph + pyrrole CH), 6.66 (s, 1 H, pyrrole CH), 6.39 (s, 1 H, pyrrole CH), 3.76 (s, 2 H, CH₂ de PCH₂), 1.29 (d, *J*_{H-P} = 12.0 Hz, 18 H, 6 CH₃ de P^tBu₂), 1.01 (s, 18 H, 6 CH₃ de 2 N^tBu) ppm. ¹³C{¹H} NMR (C₆D₆, 100.6 MHz, 298 K): δ 169.9 (s, NCN), 153.0 (m, 3 C_{ipso} of PPh₃), 137.9 (s, pyrrole C²), 134.9 (s, C_{ipso} of Ph), 129.9–127.8 (m, CHs of Phs), 123.8 (s, pyrrole CH), 111.9 (d, *J*_{C-P} = 5.4 Hz, pyrrole CH), 107.5 (s, pyrrole CH), 53.7 (s, 2 C de 2 N^tBu), 36.8 (m, 2 C de P^tBu₂), 31.8 (s, 6 CH₃ de 2 N^tBu), 30.0 (d, *J*_{C-P} = 8.3 Hz, 6 CH₃ de P^tBu₂), 25.5 (d, *J*_{C-P} = 14.2 Hz, CH₂ de PCH₂) ppm. ³¹P{¹H} NMR (C₆D₆, 162.1 MHz, 298 K): δ 79.5 (d, sat, *J*_{P-P} = 181.3 Hz, *J*_{P-Pt} = 4677.0 Hz), 60.9 (d, sat, *J*_{P-P} = 181.3 Hz, *J*_{P-Pt} = 5788.5 Hz) ppm.

X-Ray diffraction analyses

Crystals of **1_{Si}**, **2_{Pd-Si}**·THF, **3_{Ni-Ge}** and **4_{Pd-Si}** were analyzed by X-ray diffraction. A selection of crystal, measurement and refinement data is given in Table S1. Diffraction data were collected on a Bruker D8 Venture (for **1_{Si}**; MoK α radiation), an Oxford Diffraction Xcalibur Rubi Gemini (for **3_{Ni-Ge}**; MoK α radiation) and an Oxford Diffraction Xcalibur Onyx Nova (for **2_{Pd-Si}**·THF and **4_{Pd-Si}**; CuK α radiation) single crystal diffractometers. Empirical absorption corrections were applied using the SCALE3 ABSPACK algorithm as implemented in SADABS-2016/2¹⁹ (for **1_{Si}**) and CrysAlisPro RED²⁰ (for **2_{Pd-Si}**·THF, **3_{Ni-Ge}** and **4_{Pd-Si}**). The structures were solved using SIR-97.²²¹ Isotropic and full matrix anisotropic least square refinements were carried out using SHELXL.²² All non-H atoms were refined anisotropically. H atoms were set in calculated positions and were refined riding on their parent atoms. The WINGX program system²³ was used throughout the structure determinations. The molecular plots were made with MERCURY.²⁴ CCDC deposition numbers: 2001500 (**1_{Si}**), 2001501 (**2_{Pd-Si}**·THF), 2001502 (**3_{Ni-Ge}**) and 2001503 (**4_{Pd-Si}**).

Computational details

Structure optimizations were performed with the Gaussian09²⁵ suite of programs, using the wb97XD²⁶ functional, which includes the second generation of Grimme's dispersion

interaction correction,²⁷ and the cc-pVDZ²⁸ basis set (for all atoms). Frequency calculations confirmed the optimized structures as energy minima (zero imaginary eigenvalues). Orbital calculations were performed with the NBO²⁹ package.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements


This work was supported by MINECO-FEDER projects CTQ2016-75218-P and RED2018-102387-T.

References

- For general reviews on coordination chemistry of HTs, see: (a) M. Saito, *Acc. Chem. Res.*, 2018, **51**, 160; (b) M. C. Lipke, A. L. Liberman-Martin and T. D. Tilley, *Angew. Chem. Int. Ed.*, 2017, **56**, 2260; (c) R. Tacke and T. Ribbeck, *Dalton Trans.*, 2017, **46**, 13628; (d) J. Y. Corey, *Chem. Rev.* 2016, **116**, 11291; (e) L. Álvarez-Rodríguez, J. A. Cabeza, P. García-Álvarez and D. Polo, *Coord. Chem. Rev.*, 2015, **300**, 1; (f) J. Baumgartner and C. Marschner, *Rev. Inorg. Chem.*, 2014, **34**, 119; (g) B. Blom, M. Stoelzel and M. Driess, *Chem. Eur. J.*, 2013, **19**, 40; (h) R. Waterman, P. G. Hayes and T. D. Tilley, *Acc. Chem. Res.*, 2007, **40**, 712; (i) M. Okazaki, H. Tobita and H. Ogino, *Dalton Trans.*, 2003, 493; (j) M. F. Lappert and R. S. Rowe, *Coord. Chem. Rev.*, 1990, **100**, 267; (k) W. Petz, *Chem. Rev.*, 1986, **86**, 1019; (l) M. F. Lappert and P. P. Power, *J. Chem. Soc., Dalton Trans.*, 1985, 548.
- (a) Z. Benedek and T. Szilvási, *Organometallics*, 2017, **36**, 1591; (b) A. Rosas-Sánchez, I. Alvarado-Beltrán, A. Baceiredo, N. Saffon-Merceron, S. Massou, V. Ranchadell and T. Kato, *Angew. Chem. Int. Ed.*, 2017, **56**, 10549; (c) Z. Benedek and T. Szilvási, *RSC Adv.*, 2015, **5**, 5077; (d) G. Tan, S. Enthaler, S. Inoue, B. Blom and M. Driess, *Angew. Chem. Int. Ed.*, 2015, **54**, 2214; (e) J. A. Cabeza, P. García-Álvarez, E. Pérez-Carreño and D. Polo, *Chem. Eur. J.*, 2014, **20**, 8654.
- For a review on silylene TM complexes in catalysis, see: Y.-P. Zhou and M. Driess, *Angew. Chem. Int. Ed.*, 2019, **58**, 3715.
- See, for example: (a) Y.-P. Zhou, Z. Mo, M.-P. Luecke and M. Driess, *Chem. Eur. J.*, 2018, **24**, 4780; (b) J. A. Cabeza, P. García-Álvarez and L. González-Álvarez, *Chem. Commun.*, 2017, **53**, 10275; (c) T. Iimura, N. Akasaka, T. Kosai and T. Iwamoto, 2017, **46**, 8868; (d) H. Ren, Y.-P. Zhou, Y. Bai, C. Cui and M. Driess, *Chem. Eur. J.*, 2017, **23**, 5663; (e) Iimura, N. Akasaka and T. Iwamoto, 2016, **35**, 4071; (f) Y. Wu, C. Shan, Y. Sun, P. Chen, J. Ying, J. Zhu, L. Liu and Y. Zhao, *Chem. Commun.*, 2016, **52**, 13799; (g) L. Álvarez-Rodríguez, J. A. Cabeza, J. M. Fernández-Colinas, P. García-Álvarez and D. Polo, *Organometallics*, 2016, **35**, 2516; (h) Y.-P. Zhou, M. Karni, S. Yao, Y. Apeloig and M. Driess, *Angew. Chem. Int. Ed.*, 2016, **55**, 15096; (i) Y.-P. Zhou, S. Raoufoghaddam, T. Szilvási and M. Driess, *Angew. Chem. Int. Ed.*, 2016, **55**, 12868; (j) T. T. Metsänen, D. Gallego, T. Szilvási, M. Driess and M. Oestreich, *Chem. Sci.*, 2015, **6**, 7143; (k) D. Gallego, S. Inoue, B. Blom and M. Driess, *Organometallics*, 2014, **33**, 6885; (l) D. Gallego, A. Brück, E. Irran, F. Meier, F. Kaupp and M. Driess, *J. Am. Chem. Soc.*, 2013, **135**, 15617; (m) A. Brück, D. Gallego, W. Wang, E. Irran, M. Driess and J. F. Hartwig, *Angew. Chem. Int. Ed.*, 2012, **51**, 11478.
- See, for example: (a) X. Qi, T. Zheng, J. Zhou, Y. Dong, X. Zuo, X. Li, H. Sun, O. Fuhr and D. Fenske, *Organometallics*, 2019, **38**, 268; (b) A. N. Paesch, A.-K. Kreyenschmidt, R. Herbst-Irmer and D. Stalke, *Inorg. Chem.*, 2019, **58**, 7000; (c) W. Wang, S. Inoue, S. Enthaler and M. Driess, *Angew. Chem. Int. Ed.*, 2012, **51**, 6167; (d) M.-P. Luecke, D. Porwal, A. Kostenko, Y.-P. Zhou, S. Yao, M. Keck, C. Limberg, M. Oestreich and M. Driess, *Dalton Trans.*, 2017, **46**, 16412; (e) Y. Wang, A. Kostenko, S. Yao and M. Driess, *J. Am. Chem. Soc.*, 2017, **139**, 13499; (f) Z. Mo, A. Kostenko, Y.-P. Zhou, S. Yao and M. Driess, *Chem. Eur. J.*, 2018, **24**, 14608; (g) Y.-P. Zhou, S. Raoufoghaddam, T. Szilvási and M. Driess, *Angew. Chem. Int. Ed.*, 2016, **55**, 12868.
- K. M. Krebs, S. Freitag, H. Schubert, B. Gerke, R. Pöttgen and L. Wesemann, *Chem. Eur. J.*, 2015, **21**, 4628.
- K. M. Krebs, S. Freitag, J.-J. Maudrich, H. Schubert, P. Sirsch and L. Wesemann, *Dalton Trans.*, 2018, **47**, 83.
- T. Wütz, F. Diab and L. Wesemann, *Eur. J. Inorg. Chem.*, 2017, 4645.
- J. M. García, E. Ocando-Narváez, T. Kato, D. S. Coll, A. Briceño, N. Saffon-Merceron and A. Baceiredo, *Inorg. Chem.*, 2012, **52**, 8187.
- (a) J.A. Cabeza, J. M. Fernández-Colinas, P. García-Álvarez, L. González-Álvarez and E. Pérez-Carreño, *Organometallics*, 2020, **39**, DOI: 10.1021/acs.organomet.0c00188; (b) J. A. Cabeza, P. García-Álvarez, M. Gómez-Gallego, L. González-Álvarez, A. D. Merinero and M. A. Sierra, *Chem. Eur. J.*, 2019, **25**, 8635; (c) L. Álvarez-Rodríguez, J. A. Cabeza, P. García-Álvarez, M. Gómez-Gallego, L. González-Álvarez, A. D. Merinero and M. A. Sierra, *Chem. Eur. J.*, 2019, **25**, 1588-1594; (d) J. A. Cabeza, P. García-Álvarez and L. González-Álvarez, *Chem. Commun.*, 2017, **53**, 10275; (e) L. Álvarez-Rodríguez, J.A. Cabeza, P. García-Álvarez, M. Gómez-Gallego, A. D. Merinero and M. A. Sierra, *Chem. Eur. J.*, 2017, **23**, 4287; (f) J.A. Cabeza, P. García-Álvarez, R. Gobetto, L. González-Álvarez, C. Nervi, E. Pérez-Carreño and D. Polo, *Organometallics*, 2016, **35**, 1761; (g) J. A. Cabeza, P. García-Álvarez and D. Polo, *Eur. J. Inorg. Chem.*, 2016, 10; (h) L. Álvarez-Rodríguez, J. A. Cabeza, P. García-Álvarez and D. Polo, *Organometallics*, 2015, **34**, 5479; (i) J. A. Cabeza, J. M. Fernández-Colinas, P. García-Álvarez, E. Pérez-Carreño and D. Polo, *Inorg. Chem.*, 2015, **54**, 4850; (j) L. Álvarez-Rodríguez, J. A. Cabeza, P. García-Álvarez, E. Pérez-Carreño and D. Polo, *Inorg. Chem.*, 2015, **54**, 2983; (k) J. A. Cabeza, P. García-Álvarez, E. Pérez-Carreño and D. Polo, *Inorg. Chem.*, 2014, **53**, 8735; (l) J. A. Cabeza, P. García-Álvarez, E. Pérez-Carreño and D. Polo, *Chem. Eur. J.*, 2014, **20**, 8654; (m) J. A. Cabeza, P. García-Álvarez and D. Polo, *Dalton Trans.*, 2013, **42**, 1329.
- (a) L. Álvarez-Rodríguez, J. Brugos, J. A. Cabeza, P. García-Álvarez, E. Pérez-Carreño and D. Polo, *Chem. Commun.*, 2017, **53**, 893; (b) L. Álvarez-Rodríguez, J. Brugos, J. A. Cabeza, P. García-Álvarez and E. Pérez-Carreño, *Chem. Eur. J.*, 2017, **23**, 15107; (c) J. Brugos, J. A. Cabeza, P. García-Álvarez and E. Pérez-Carreño, *Organometallics*, 2018, **37**, 1507; (d) J. Brugos, J. A. Cabeza, P. García-Álvarez, E. Pérez-Carreño and D. Polo *Dalton Trans.*, 2018, **477**, 4534;
- (a) J. A. Cabeza, I. Fernández, J. M. Fernández-Colinas, P. García-Álvarez and C. J. Laglera-Gándara, *Dalton Trans.*, 2019, **48**, 13273; (b) J. A. Cabeza, I. Fernández, J. M. Fernández-Colinas, P. García-Álvarez and C. J. Laglera-Gándara, *Chem. Eur. J.*, 2019, **25**, 12423; (c) J. A. Cabeza, P. García-Álvarez and C. J. Laglera-Gándara, *Eur. J. Inorg. Chem.*, 2020, 784.
- Z. Feng, Y. Jiang, H. Ruan, Y. Zhao, G. Tan, L. Zhang and X. Wang, *Dalton Trans.*, 2019, **48**, 14975.
- S. S. Sen, H. W. Roesky, D. Stern, J. Henn and D. Stalke, *J. Am. Chem. Soc.*, 2010, **132**, 1123.
- S. Nagendran, S. S. Sen, H. W. Roesky, D. Koley, H. Grubmüller, A. Pal and R. Herbst-Irmer, *Organometallics*, 2008, **27**, 5459.
- C. J. Mathews, P. J. Smith and T. Welton, *J. Mol. Catal. A. Chem.*, 2003, **206**, 77.
- D. Drew and J. R. Doyle, *Inorg. Synth.*, 1972, **13**, 48.
- R. Ugo, F. Cariati and G. La Monica, *Inorg. Synth.*, 1990, **28**, 123.
- SADABS-2016/2: L. Krause, R. Herbst-Irmer, G. M. Sheldrick and D. Stalke, *J. Appl. Crystallogr.*, 2015, **48**, 3.
- CrysAlisPro RED, version 1.171.37.35: Oxford Diffraction Ltd., Oxford, UK, 2014.
- SIR-97: A. Altomare, M. C. Burla, M. Camalli, G. L. Casciarano, C. Giacovazzo, A. Guagliardi, A. G. C. Moliterni, G. Polidori and R. Spagna, *J. Appl. Crystallogr.*, 1999, **32**, 115.
- SHELXL-2014: G. M. Sheldrick, *Acta Cryst.*, 2008, **A64**, 112.
- WINGX, version 2013.3: J. Farrugia, *J. Appl. Crystallogr.*, 2012, **45**, 849.
- MERCURY, version 2020.1 (build 280191): Cambridge Crystallographic Data Centre, Cambridge, UK, 2018.

- 25 *Gaussian 09*, revision D.01: M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, N. J. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, Gaussian, Inc., Wallingford, CT, 2009.
- 26 J.-D. Chai and M. Head-Gordon, *Phys. Chem. Chem. Phys.*, 2008, **10**, 6615.
- 27 (a) S. Ehrlich, J. Moellmann and S. Grimme, *Acc. Chem. Res.*, 2013, **46**, 916; (b) S. Grimme, *Comp. Mol. Sci.*, 2011, **1**, 211; (c) T. Schwabe and S. Grimme, *Acc. Chem. Res.*, 2008, **41**, 569.
- 28 T. H. Dunning, *J. Chem. Phys.*, 1989, **90**, 1007.
- 29 (a) J. P. Foster and F. Weinhold, *J. Am. Chem. Soc.*, 1980, **102**, 7211; (b) A. E. Reed and F. Weinhold, *J. Chem. Phys.*, 1985, **83**, 1736; (c) A. E. Reed, R. B. Weinstock and F. Weinhold, *J. Chem. Phys.*, 1985, **83**, 735; (d) A. E. Reed, L. A. Curtiss and F. Weinhold, *Chem. Rev.*, 1988, **88**, 899.

Figure and Text for the Table of Contents



The silylene- and germylene-decorated phosphanes **1_{Si}** and **2_{Ge}** have been prepared and their coordination abilities have been proven by reactions with transition-metal complexes.