

Proceedings
of the
XXVI Congreso de Ecuaciones
Diferenciales y Aplicaciones
XVI Congreso de Matemática Aplicada

Gijón (Asturias), Spain

June 14-18, 2021



SēMA
Sociedad Española
de Matemática Aplicada



Universidad de Oviedo

Editors:

Rafael Gallego, Mariano Mateos

Esta obra está bajo una licencia Reconocimiento- No comercial- Sin Obra Derivada 3.0 España de Creative Commons. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-nc-nd/3.0/es/> o envíe una carta a Creative Commons, 171 Second Street, Suite 300, San Francisco, California 94105, USA.



Reconocimiento- No Comercial- Sin Obra Derivada (by-nc-nd): No se permite un uso comercial de la obra original ni la generación de obras derivadas.



Usted es libre de copiar, distribuir y comunicar públicamente la obra, bajo las condiciones siguientes:



Reconocimiento – Debe reconocer los créditos de la obra de la manera especificada por el licenciador:

Coordinadores: Rafael Gallego, Mariano Mateos (2021), Proceedings of the XXVI Congreso de Ecuaciones Diferenciales y Aplicaciones / XVI Congreso de Matemática Aplicada. Universidad de Oviedo.

La autoría de cualquier artículo o texto utilizado del libro deberá ser reconocida complementariamente.



No comercial – No puede utilizar esta obra para fines comerciales.



Sin obras derivadas – No se puede alterar, transformar o generar una obra derivada a partir de esta obra.

© 2021 Universidad de Oviedo

© Los autores

Universidad de Oviedo

Servicio de Publicaciones de la Universidad de Oviedo

Campus de Humanidades. Edificio de Servicios. 33011 Oviedo (Asturias)

Tel. 985 10 95 03 Fax 985 10 95 07

[http: www.uniovi.es/publicaciones](http://www.uniovi.es/publicaciones)

servipub@uniovi.es

ISBN: 978-84-18482-21-2

Todos los derechos reservados. De conformidad con lo dispuesto en la legislación vigente, podrán ser castigados con penas de multa y privación de libertad quienes reproduzcan o plagien, en todo o en parte, una obra literaria, artística o científica, fijada en cualquier tipo de soporte, sin la preceptiva autorización.

Foreword

It is with great pleasure that we present the Proceedings of the 26th Congress of Differential Equations and Applications / 16th Congress of Applied Mathematics (XXVI CEDYA / XVI CMA), the biennial congress of the Spanish Society of Applied Mathematics SĒMA, which is held in Gijón, Spain from June 14 to June 18, 2021.

In this volume we gather the short papers sent by some of the almost three hundred and twenty communications presented in the conference. Abstracts of all those communications can be found in the abstract book of the congress. Moreover, full papers by invited lecturers will shortly appear in a special issue of the SĒMA Journal.

The first CEDYA was celebrated in 1978 in Madrid, and the first joint CEDYA / CMA took place in Málaga in 1989. Our congress focuses on different fields of applied mathematics: Dynamical Systems and Ordinary Differential Equations, Partial Differential Equations, Numerical Analysis and Simulation, Numerical Linear Algebra, Optimal Control and Inverse Problems and Applications of Mathematics to Industry, Social Sciences, and Biology. Communications in other related topics such as Scientific Computation, Approximation Theory, Discrete Mathematics and Mathematical Education are also common.

For the last few editions, the congress has been structured in mini-symposia. In Gijón, we will have eighteen minis-symposia, proposed by different researchers and groups, and also five thematic sessions organized by the local organizing committee to distribute the individual contributions. We will also have a poster session and ten invited lectures. Among all the mini-symposia, we want to highlight the one dedicated to the memory of our colleague Francisco Javier “Pancho” Sayas, which gathers two plenary lectures, thirty-six talks, and more than forty invited people that have expressed their wish to pay tribute to his figure and work.

This edition has been deeply marked by the COVID-19 pandemic. First scheduled for June 2020, we had to postpone it one year, and move to a hybrid format. Roughly half of the participants attended the conference online, while the other half came to Gijón. Taking a normal conference and moving to a hybrid format in one year has meant a lot of efforts from all the parties involved. Not only did we, as organizing committee, see how much of the work already done had to be undone and redone in a different way, but also the administration staff, the scientific committee, the mini-symposia organizers, and many of the contributors had to work overtime for the change.

Just to name a few of the problems that all of us faced: some of the already accepted mini-symposia and contributed talks had to be withdrawn for different reasons (mainly because of the lack of flexibility of the funding agencies); it became quite clear since the very first moment that, no matter how well things evolved, it would be nearly impossible for most international participants to come to Gijón; reservations with the hotels and contracts with the suppliers had to be cancelled; and there was a lot of uncertainty, and even anxiety could be said, until we were able to confirm that the face-to-face part of the congress could take place as planned.

On the other hand, in the new open call for scientific proposals, we had a nice surprise: many people that would have not been able to participate in the original congress were sending new ideas for mini-symposia, individual contributions and posters. This meant that the total number of communications was about twenty percent greater than the original one, with most of the new contributions sent by students.

There were almost one hundred and twenty students registered for this CEDYA / CMA. The hybrid format allows students to participate at very low expense for their funding agencies, and this gives them the opportunity to attend different conferences and get more merits. But this, which can be seen as an advantage, makes it harder for them to obtain a full conference experience. Alfréd Rényi said: “a mathematician is a device for turning coffee into theorems”. Experience has taught us that a congress is the best place for a mathematician to have a lot of coffee. And coffee cannot be served online.

In Gijón, June 4, 2021

The Local Organizing Committee from the Universidad de Oviedo

Scientific Committee

- Juan Luis Vázquez, Universidad Autónoma de Madrid
- María Paz Calvo, Universidad de Valladolid
- Laura Grigori, INRIA Paris
- José Antonio Langa, Universidad de Sevilla
- Mikel Lezaun, Euskal Herriko Unibersitatea
- Peter Monk, University of Delaware
- Ira Neitzel, Universität Bonn
- José Ángel Rodríguez, Universidad de Oviedo
- Fernando de Terán, Universidad Carlos III de Madrid

Sponsors

- Sociedad Española de Matemática Aplicada
- Departamento de Matemáticas de la Universidad de Oviedo
- Escuela Politécnica de Ingeniería de Gijón
- Gijón Convention Bureau
- Ayuntamiento de Gijón

Local Organizing Committee from the Universidad de Oviedo

- Pedro Alonso Velázquez
- Rafael Gallego
- Mariano Mateos
- Omar Menéndez
- Virginia Selgas
- Marisa Serrano
- Jesús Suárez Pérez del Río

Contents

On numerical approximations to diffuse-interface tumor growth models Acosta-Soba D., Guillén-González F. and Rodríguez-Galván J.R.	8
An optimized sixth-order explicit RKN method to solve oscillating systems Ahmed Demba M., Ramos H., Kumam P. and Watthayu W.	15
The propagation of smallness property and its utility in controllability problems Apraiz J.	23
Theoretical and numerical results for some inverse problems for PDEs Apraiz J., Doubova A., Fernández-Cara E. and Yamamoto M.	31
Pricing TARN options with a stochastic local volatility model Arregui I. and Ráfales J.	39
XVA for American options with two stochastic factors: modelling, mathematical analysis and numerical methods Arregui I., Salvador B., Ševčovič D. and Vázquez C.	44
A numerical method to solve Maxwell's equations in 3D singular geometry Assous F. and Raichik I.	51
Analysis of a SEIRS metapopulation model with fast migration Atienza P. and Sanz-Lorenzo L.	58
Goal-oriented adaptive finite element methods with optimal computational complexity Becker R., Gantner G., Innerberger M. and Praetorius D.	65
On volume constraint problems related to the fractional Laplacian Bellido J.C. and Ortega A.	73
A semi-implicit Lagrange-projection-type finite volume scheme exactly well-balanced for 1D shallow-water system Caballero-Cárdenas C., Castro M.J., Morales de Luna T. and Muñoz-Ruiz M.L.	82
SEIRD model with nonlocal diffusion Calvo Pereira A.N.	90
Two-sided methods for the nonlinear eigenvalue problem Campos C. and Roman J.E.	97
Fractionary iterative methods for solving nonlinear problems Candelario G., Cordero A., Torregrosa J.R. and Vassileva M.P.	105
Well posedness and numerical solution of kinetic models for angiogenesis Carpio A., Cebrián E. and Duro G.	109
Variable time-step modal methods to integrate the time-dependent neutron diffusion equation Carreño A., Vidal-Ferrándiz A., Ginestar D. and Verdú G.	114

Homoclinic bifurcations in the unfolding of the nilpotent singularity of codimension 4 in R^4 Casas P.S., Drubi F. and Ibáñez S.	122
Different approximations of the parameter for low-order iterative methods with memory Chicharro F.I., Garrido N., Sarría I. and Orcos L.	130
Designing new derivative-free memory methods to solve nonlinear scalar problems Cordero A., Garrido N., Torregrosa J.R. and Triguero P.	135
Iterative processes with arbitrary order of convergence for approximating generalized inverses Cordero A., Soto-Quirós P. and Torregrosa J.R.	141
FCF formulation of Einstein equations: local uniqueness and numerical accuracy and stability Cordero-Carrión I., Santos-Pérez S. and Cerdá-Durán P.	148
New Galilean spacetimes to model an expanding universe De la Fuente D.	155
Numerical approximation of dispersive shallow flows on spherical coordinates Escalante C. and Castro M.J.	160
New contributions to the control of PDEs and their applications Fernández-Cara E.	167
Saddle-node bifurcation of canard limit cycles in piecewise linear systems Fernández-García S., Carmona V. and Teruel A.E.	172
On the amplitudes of spherical harmonics of gravitational potencial and generalised products of inertia Floría L.	177
Turing instability analysis of a singular cross-diffusion problem Galiano G. and González-Tabernero V.	184
Weakly nonlinear analysis of a system with nonlocal diffusion Galiano G. and Velasco J.	192
What is the humanitarian aid required after tsunami? González-Vida J.M., Ortega S., Macías J., Castro M.J., Michelini A. and Azzarone A.	197
On Keller-Segel systems with fractional diffusion Granero-Belinchón R.	201
An arbitrary high order ADER Discontinuous Galerking (DG) numerical scheme for the multilayer shallow water model with variable density Guerrero Fernández E., Castro Díaz M.J., Dumbser M. and Morales de Luna T.	208
Picard-type iterations for solving Fredholm integral equations Gutiérrez J.M. and Hernández-Verón M.A.	216
High-order well-balanced methods for systems of balance laws based on collocation RK ODE solvers Gómez-Bueno I., Castro M.J., Parés C. and Russo G.	220
An algorithm to create conservative Galerkin projection between meshes Gómez-Molina P., Sanz-Lorenzo L. and Carpio J.	228
On iterative schemes for matrix equations Hernández-Verón M.A. and Romero N.	236
A predictor-corrector iterative scheme for improving the accessibility of the Steffensen-type methods Hernández-Verón M.A., Magreñán A.A., Martínez E. and Sukhjit S.	242

CONTENTS

Recent developments in modeling free-surface flows with vertically-resolved velocity profiles using moments Koellermeier J.	247
Stability of a one degree of freedom Hamiltonian system in a case of zero quadratic and cubic terms Lanchares V. and Bardin B.	253
Minimal complexity of subharmonics in a class of planar periodic predator-prey models López-Gómez J., Muñoz-Hernández E. and Zanolin F.	258
On a non-linear system of PDEs with application to tumor identification Maestre F. and Pedregal P.	265
Fractional evolution equations in discrete sequences spaces Miana P.J.	271
KPZ equation approximated by a nonlocal equation Molino A.	277
Symmetry analysis and conservation laws of a family of non-linear viscoelastic wave equations Márquez A. and Bruzón M.	284
Flux-corrected methods for chemotaxis equations Navarro Izquierdo A.M., Redondo Nebel M.V. and Rodríguez Galván J.R.	289
Ejection-collision orbits in two degrees of freedom problems Ollé M., Álvarez-Ramírez M., Barrabés E. and Medina M.	295
Teaching experience in the Differential Equations Semi-Virtual Method course of the Tecnológico de Costa Rica Oviedo N.G.	300
Nonlinear analysis in lorentzian geometry: the maximal hypersurface equation in a generalized Robertson-Walker spacetime Pelegrín J.A.S.	307
Well-balanced algorithms for relativistic fluids on a Schwarzschild background Pimentel-García E., Parés C. and LeFloch P.G.	313
Asymptotic analysis of the behavior of a viscous fluid between two very close mobile surfaces Rodríguez J.M. and Taboada-Vázquez R.	321
Convergence rates for Galerkin approximation for magnetohydrodynamic type equations Rodríguez-Bellido M.A., Rojas-Medar M.A. and Sepúlveda-Cerda A.	325
Asymptotic aspects of the logistic equation under diffusion Sabina de Lis J.C. and Segura de León S.	332
Analysis of turbulence models for flow simulation in the aorta Santos S., Rojas J.M., Romero P., Lozano M., Conejero J.A. and García-Fernández I.	339
Overdetermined elliptic problems in unduloid-type domains with general nonlinearities Wu J.	344

Pricing TARN options with a stochastic local volatility model

Iñigo Arregui^{1,2}, Jonatan Ráfales^{1,2}

1. Depto. de Matemáticas, Universidad de A Coruña, A Coruña, Spain
 2. CITIC, A Coruña, Spain

Abstract

Target Accumulation Redemption Notes (TARNs) are financial derivatives which give their holders the right to receive periodic coupons until the accumulated sum of those ones reaches an agreed target. In this work, we solve a partial differential equations (PDEs) model for pricing TARN options by implementing an alternating-direction implicit finite difference method (ADI method). We combine the numerical solution with a stochastic local volatility (SLV) technique and show the numerical results for a particular example.

1. Introduction

European and American options are the most well-known derivative products, and have been widely studied from the financial and mathematical points of view. For these options, jointly known as vanilla options, their price depends mainly on the value of the underlying asset.

On the opposite side, exotic options present different features which also have an effect on the price. For example, Bermudan options offer multiple exercise dates, the pay-off of Asian options depends on the average price of the underlying asset and the pay-off of barrier options depends on whether or not the price of the underlying asset reaches an agreed value during the option's lifetime. Other examples of exotic products are the target redemption products, whose notional amount increases until a certain target is reached [5]. In particular, the value of a Target Accumulation Redemption Note (TARN) depends on an accumulated amount: if the sum of coupons reaches an agreed target before the maturity date, the holder of the note receives a final payment, also known as knockout, and the contract terminates. These products are usually traded in foreign exchange (FX) markets.

2. Mathematical model

We propose a stochastic local volatility model based on Heston model [4]:

$$\begin{cases} dS_t = (r_d(t) - r_f(t))S_t dt + L(S_t, t)\sqrt{V_t}S_t dW_t^1 \\ dV_t = \kappa(\theta - V_t) dt + \lambda\sqrt{V_t} dW_t^2 \\ dW_t^1 dW_t^2 = \rho dt, \end{cases} \quad (2.1)$$

where L is the leverage function, which represents the contribution of the local volatility and will be calibrated with the help of market data, S_t is the underlying asset, V_t is the stochastic variance, W_t^1 and W_t^2 are two Brownian motions, r_d and r_f are the domestic and foreign interest rates, respectively, and κ , θ , λ and ρ are the Heston parameters.

Let $r(t) = r_d(t) - r_f(t)$. Let us assume $2\kappa\theta \leq \lambda^2$ and $V_0 > 0$. Then, Feller condition states that $V_t > 0$ for every $t > 0$. Moreover, we will assume S_0 , κ , θ and λ are strictly positive, $-1 < \rho < 1$ and the leverage function is positive and bounded. Under these hypotheses, it is proven [4] that there exists a unique solution of model (2.1) and there also exists a function $p := p(S, V, t)$, called transition probability function, solution of the Fokker-Planck (FP) equation:

$$\frac{\partial p}{\partial t} = -\frac{\partial}{\partial S}(r(t)Sp) - \frac{\partial}{\partial V}(\kappa(\theta - V)p) + \frac{1}{2}\frac{\partial^2}{\partial S^2}(L^2S^2Vp) + \frac{\partial^2}{\partial S\partial V}(\lambda\rho LSVp) + \frac{1}{2}\frac{\partial^2}{\partial V^2}(\lambda^2Vp), \quad (2.2)$$

such that the leverage function can be written as [4]:

$$L(S, t) = \sigma_{LV}(S, t) \sqrt{\frac{\int_{\mathbb{R}} p(S, V, t) dV}{\int_{\mathbb{R}} V p(S, V, t) dV}}. \quad (2.3)$$

Let $S(t)$ be the FX rate at time t , t_0 the actual date and t_1, t_2, \dots, t_K the so-called fixing dates. Moreover, let E be the strike, U the target accrual level and $A(t)$ the accumulated amount at time t [3]. On each fixing date, t_k , there is a cash flow payment:

$$\tilde{C}_k \equiv \beta(S(t_k) - E) \times 1_{\beta \times S(t_k) \geq \beta \times E},$$

where β is a strategy on foreign exchange ($\beta = 1$ for a call option and $\beta = -1$ for a put option), until the accumulated amount A reaches the target U . Let $t_{\tilde{K}}$ be the first fixing date when the target is breached:

$$\tilde{K} = \min_{k=1, \dots, K} \{k : A(t_k) \geq U\}$$

and let $\tilde{K} = K$ if the target is not breached, the payment can be written as:

$$C_k(S, A) = \begin{cases} \tilde{C}_k \times \left(1_{A(t_{k-1}) + \tilde{C}_k < U} + W_k \times 1_{A(t_{k-1}) + \tilde{C}_k \geq U} \right), & \text{if } t_k \leq t_{\tilde{K}}, \\ 0, & \text{if } t_k > t_{\tilde{K}}, \end{cases} \quad (2.4)$$

where $A(t_{k-1})$ is the accumulated amount immediately after t_{k-1} . This magnitude is given by a piecewise constant function:

$$A(t) = \begin{cases} A(t_{k-1}), & \text{if } t_{k-1} \leq t < t_k, \\ A(t_{k-1}) + C_k(S(t_k), A(t_{k-1})), & \text{if } t = t_k. \end{cases}$$

Moreover, W_k is a weight that depends on the knockout when the target is breached. We will consider three types of knockout, such that the weight can be written as:

$$W_k = \begin{cases} 1, & \text{in the case of full gain,} \\ \frac{U - A(t_{k-1})}{\beta \times (S(t_k) - E)}, & \text{in the case of part gain,} \\ 0, & \text{in the case of no gain.} \end{cases}$$

Finally, let $u := u(S, V, t, A)$ be the value of a TARN option, the following SLV option pricing PDE:

$$\frac{\partial u}{\partial t} + r(t)S \frac{\partial u}{\partial S} + (\kappa(\theta - V)) \frac{\partial u}{\partial V} + \frac{1}{2}L^2VS^2 \frac{\partial^2 u}{\partial S^2} + \rho L\lambda SV \frac{\partial^2 u}{\partial S \partial V} + \frac{1}{2}\lambda^2V \frac{\partial^2 u}{\partial V^2} - r_d(t)u = 0$$

is valid between fixing dates. Furthermore, for each fixing date we can pose:

$$u(S, V, t_k^-, A(t_k^-)) = u(S, V, t_k, A(t_k^-)) + C_k(S, A(t_k^-)),$$

where C_k is given by (2.4) and t_k^- is the time infinitesimally before t_k .

2.1. Numerical methods

We compute the optimal Heston parameters $(\kappa, \theta, \lambda, \rho)$ for the calibration of the SLV model. For each maturity, the COS method is developed to price European options under the Heston model, $[w^{\text{Hes}}]_i^m := w^{\text{Hes}}(E_i, T^m)$ and the Levenberg-Marquardt non-linear least squares algorithm is applied to find the optimal parameters by minimizing

$$\min_{\kappa, \theta, \lambda, \rho} \sum_{i=1}^{N_E} ([w^{\text{Hes}}(\kappa, \theta, \lambda, \rho)]_i^m - w_i^m)^2,$$

where $w_i^m := w(E_i, T^m)$ are the market data for different strikes and maturities.

As Feller condition does not always hold in real markets, we propose a logarithmic change of variable [2]. Thus, let S_0 and V_0 be the initial values of the underlying and the variance, and let

$$X_t = \log(S_t/S_0), \quad Z_t = \log(V_t/V_0).$$

In the new domain $(-\infty, \infty) \times (-\infty, \infty)$, model (2.1) is rewritten as:

$$\begin{cases} dX_t = (r(t) - \frac{1}{2}L(X_t, t)^2V_0e^{Z_t})dt + L(X_t, t)\sqrt{V_0e^{Z_t}}dW_t^1 \\ dZ_t = ((\kappa\theta - \frac{1}{2}\lambda^2)\frac{1}{V_0e^{Z_t}} - \kappa)dt + \lambda\frac{1}{\sqrt{V_0e^{Z_t}}}dW_t^2 \\ dW_t^1 dW_t^2 = \rho dt. \end{cases}$$

A previous step to apply numerical methods is the truncation of the new unbounded domain to a bounded one. Thus, we consider the fixed domain $\Omega = (X_{\min}, X_{\max}) \times (Z_{\min}, Z_{\max})$, which is finer around $(X_0 = 0, Z_0 = 0)$, and pose the FP equation (2.2):

$$\begin{aligned} \frac{\partial p}{\partial t} = & -\frac{\partial}{\partial X} \left((r(t) - \frac{1}{2}L^2V_0e^Z)p \right) - \frac{\partial}{\partial Z} \left(((\kappa\theta - \frac{1}{2}\lambda^2)\frac{1}{V_0e^Z} - \kappa)p \right) \\ & + \frac{1}{2}\frac{\partial^2}{\partial X^2} (L^2V_0e^Z p) + \frac{\partial^2}{\partial X \partial Z} (\lambda \rho L p) + \frac{1}{2}\frac{\partial^2}{\partial Z^2} \left(\lambda^2 \frac{1}{V_0e^Z} p \right), \end{aligned} \quad (2.5)$$

with the initial condition:

$$p(X, Z, 0) = \delta(X)\delta(Z), \quad (2.6)$$

where δ is the Dirac function. Additionally, we impose the boundary conditions:

$$\frac{\partial^2 p}{\partial X^2}(X_{\min}, Z, t) = 0, \quad \frac{\partial^2 p}{\partial X^2}(X_{\max}, Z, t) = 0, \quad \frac{\partial^2 p}{\partial Z^2}(X, Z_{\min}, t) = 0, \quad \frac{\partial^2 p}{\partial Z^2}(X, Z_{\max}, t) = 0. \quad (2.7)$$

Moreover, the leverage function (2.3) is given by:

$$L(X, t) = \sigma_{LV}(X, t) \sqrt{\frac{\int_{\mathbb{R}^+} p(X, Z, t) dZ}{\int_{\mathbb{R}^+} V_0 e^Z p(X, Z, t) dZ}}. \quad (2.8)$$

As we can see in (2.5) and (2.8), the computing of the transition probability function needs the leverage function and reciprocally. Therefore, we propose a fixed point scheme to solve the problem at each time step, in which an alternating directions implicit (ADI) method is developed to solve the FP problem and the trapezoidal rule is applied to approximate the leverage function. The modified Douglas ADI scheme at each step can be written as:

$$A = p^{n-1} + \Delta t^n [F_0(p^{n-1}, t^{n-1}) + F_1(p^{n-1}, t^{n-1}) + F_2(p^{n-1}, t^{n-1})], \quad (2.9)$$

$$B - \alpha \Delta t^n F_1(B, t^n) = A - \alpha \Delta t^n F_1(p^{n-1}, t^{n-1}),$$

$$C - \alpha \Delta t^n F_2(C, t^n) = B - \alpha \Delta t^n F_2(p^{n-1}, t^{n-1}), \quad (2.10)$$

$$p^n = C,$$

for $n = 1, 2, \dots, N_T$, where

$$F_0(p, t) = \frac{\partial^2}{\partial X \partial Z} (\lambda \rho L p),$$

$$F_1(p, t) = -\frac{\partial}{\partial Z} \left((\kappa \theta - \frac{1}{2} \lambda^2) \frac{1}{V_0 e^Z} - \kappa \right) p + \frac{1}{2} \frac{\partial^2}{\partial Z^2} \left(\frac{\lambda^2}{V_0 e^Z} p \right),$$

$$F_2(p, t) = -\frac{\partial}{\partial X} \left((r(t) - \frac{1}{2} L^2 V_0 e^Z) p \right) + \frac{1}{2} \frac{\partial^2}{\partial X^2} (L^2 V_0 e^Z p).$$

In addition, a mixing fraction parameter, η , is applied to the volatility of the volatility, λ :

$$dV_t = \kappa(\theta - V_t)dt + \eta \lambda \sqrt{V_t} dW_t^2.$$

For each maturity, the ADI method is developed to price options under the SLV model, $[y^{\text{SLV}}]_i^m := y^{\text{SLV}}(E_i, T^m)$ and the golden section search algorithm is applied to determine the optimal parameter by minimizing

$$\min_{\eta} \sum_{i=1}^{N_E} ([y^{\text{SLV}}(\eta)]_i^m - y_i^m)^2,$$

where $y_i^m := y(E_i, T^m)$ are the market data for different strikes and maturities.

Finally, we introduce the time-to-maturity variable ($\tau = T - t$) and deduce the TARN price PDE in terms of X and Z :

$$\begin{aligned} \frac{\partial u}{\partial \tau} = & (r(\tau) - \frac{1}{2} L^2 V_0 e^Z) \frac{\partial u}{\partial X} + \frac{1}{2} L^2 V_0 e^Z \frac{\partial^2 u}{\partial X^2} + \lambda \eta \rho L \frac{\partial^2 u}{\partial X \partial Z} \\ & + ((\kappa \theta - \frac{1}{2} \lambda^2 \eta^2) \frac{1}{V_0 e^Z} - \kappa) \frac{\partial u}{\partial Z} + \frac{1}{2} \frac{\lambda^2 \eta^2}{V_0 e^Z} \frac{\partial^2 u}{\partial Z^2} - r_d(\tau) u, \end{aligned} \quad (2.11)$$

with the initial condition:

$$u(X, Z, 0, A) = 0. \quad (2.12)$$

Additionally, we assume the boundary conditions [4]:

$$\begin{aligned} \frac{1}{S_0^2} e^{-2X} \left(\frac{\partial^2 u}{\partial X^2}(X_{\min}, Z, t) - \frac{\partial u}{\partial X}(X_{\min}, Z, t) \right) = 0, \quad \frac{1}{V_0^2} e^{-2Z} \left(\frac{\partial^2 u}{\partial Z^2}(X, Z_{\min}, t) - \frac{\partial u}{\partial Z}(X, Z_{\min}, t) \right) = 0, \\ \frac{1}{S_0^2} e^{-2X} \left(\frac{\partial^2 u}{\partial X^2}(X_{\max}, Z, t) - \frac{\partial u}{\partial X}(X_{\max}, Z, t) \right) = 0, \quad \frac{1}{V_0^2} e^{-2Z} \left(\frac{\partial^2 u}{\partial Z^2}(X, Z_{\max}, t) - \frac{\partial u}{\partial Z}(X, Z_{\max}, t) \right) = 0. \end{aligned} \quad (2.13)$$

As for the numerical solution of the FP problem, we also propose the use of the ADI algorithm to solve (2.11 – 2.13), jointly with the jump condition for each fixing date τ_k :

$$u(X, Z, \tau_k^+, A(\tau_k^+)) = u(X, Z, \tau_k, A(\tau_k^+)) + C_k(X, A(\tau_k^+)) + C_k(X, A(\tau_k^+)),$$

where τ_k^+ is the time infinitesimally after τ_k . Fig. 1 shows a sketch of the scheme.




Fig. 1 Jump condition

2.2. Numerical results

We present a numerical test for the valuation of a TARN call option in the frame of foreign exchange. For this aim, we have considered the US dollar and British pound as domestic and foreign currencies, respectively, an initial underlying $S_0 = 1.320$, an initial variance $V_0 = 0.004$ and a strike $E = 1.283$. Furthermore, the domestic and foreign interest rates are shown in Tab. 1 of [1], the maturity period is $T = 12$ months and the fixing dates are taken every 30 days, thus $K = 12$.

Fig. 2 shows the market implied volatility (left) and the local volatility (right), which is computed by means of Dupire's formula. Thus, we apply the previous techniques to compute the Heston parameters, which are shown in Tab. 3 (left) of [1].




Fig. 2 The implied volatility (left) and the local volatility (right)

Next, we have approximated the (X, Z) domain with a mesh similar to the one plotted in Fig. 3, consisting of 400×100 nodes. As we have previously detailed, the mesh is finer around the point $(X_0 = 0, Z_0 = 0)$. We have also refined the mesh for values close to Z_{\min} in order to minimize the errors arising from the fact that the Feller condition may be not accurate for these values of the volatility. Moreover, we have used 180 time steps and the parameter $\alpha = 0.5$ in the ADI method. Fig. 4 shows the computed solution of the FP problem at the maturity (left) and the leverage function (right).




Fig. 3 Mesh




Fig. 4 Numerical solution of the FP equation (left) and the leverage function (right)

Finally, we calibrate the mixing fraction parameter η , which is shown in Tab. 3 (right) of [1], and show the numerical approximation of the TARN price for different knockouts in Fig. 5. More details and results are available in [1].




Fig. 5 TARN option price: no gain (left), part gain (center) and full gain (right) knockouts for $U = 0.90$.

2.3. Conclusions

We solve a partial differential equations model to price TARN options. We have improved previous results by introducing a SLV technique in order to better reflect the market volatilities (taking advantage of local volatility methods) on a path dependent derivative product (for which stochastic volatility methods are more convenient). This SLV approach can be extended to other kinds of exotic options.

An alternating directions method (ADI) is implemented and the volatility surfaces, transition probability function, leverage function and option price are computed. Therefore, we have a tool to value this kind of exotic options.

Acknowledgements

Authors are financially supported by Spanish Ministerio de Ciencia e Innovación (grant PID2019-108584RB-100). Authors also acknowledge the support received from the Centro de Investigación de Galicia (CITIC), funded by Xunta de Galicia and the European Union (European Regional Development Fund- Galicia 2014-2020 Program), by grant ED431G 2019/01.

References

- [1] Iñigo Arregui and Jonatan Ráfales. A stochastic local volatility technique for TARN options. *International Journal of Computer Mathematics*, 97(5):1133–1149, 2020.
- [2] Fang Fang and Cornelis W. Oosterlee. A novel pricing method for European options based on Fourier-Cosine series expansions. *SIAM Journal on Scientific Computing*, 31(2):826–848, 2009.
- [3] Xiaolin Luo and Pavel V. Shevchenko. Pricing TARNs using a finite difference method. *The Journal of Derivatives*, 23(1):62–72, 2015.
- [4] Yu Tian. *The hybrid stochastic-local volatility model with applications in pricing FX options*. PhD thesis, Monash University, 2013.
- [5] Uwe Wystup. *FX options and Structured Products*. John Wiley & Sons, Chichester, 2017.