

2 **Non-invasive genetic study of the endangered Cantabrian brown**
3 **bear (*Ursus arctos*)**

4 **Trinidad Pérez · Fernando Vázquez · Javier Naves ·**
5 **Alberto Fernández · Ana Corao · Jesús Albornoz ·**
6 **Ana Domínguez**

7 Received: 18 October 2007 / Accepted: 22 March 2008
8 © Springer Science+Business Media B.V. 2008

9 **Abstract** The Brown Bear (*Ursus arctos*) population
10 present in the Cantabrian Mountains has suffered a dra-
11 matic decline in recent centuries and is now threatened
12 with extinction. This situation has led to the development
13 and implementation of a species recovery plan. To
14 accomplish this plan, we need to improve our knowledge
15 about the ecology, demography and genetics of this pop-
16 ulation. This paper presents the genetic analysis of the
17 Cantabrian brown bear population using non-invasive
18 samples (faeces and hairs) collected between 2004 and
19 2006. It was necessary to optimize a set of 18 microsatellite
20 loci and a sex marker (several new multiplex reactions
21 were developed) to obtain a suitable probability of identity
22 among genotypes to work with this small, deeply structured
23 population. Genotyping of 48 individuals was carried out
24 using a two-step PCR protocol to increase the quality of the
25 multilocus genotypes. Validation of genotypes was per-
26 formed using a multi-tube approach combined with
27 different software programmes to measure their error rate
28 and reliability. Diversity in the Cantabrian population was

low ($H_e = 0.51$) and the population was markedly subdivi- 29
divided into two subpopulations (western and eastern) 30
without current gene flow between them. The level of 31
divergence between the two subpopulations ($F_{st} = 0.41$) 32
and the extremely low diversity in the eastern group 33
($H_e = 0.25$) indicate that this has had an extremely low 34
effective population size and had been isolated from the 35
main group during the last century. Connectivity between 36
the two subpopulations will be of prime importance for the 37
long-term survival of this species in the Cantabrian 38
Mountains. 39

40
41 **Keywords** *Ursus arctos* · Microsatellites ·
42 Non-invasive samples · Individual identification ·
43 Population structure 44

45 **Introduction**


46 Brown bears were once found throughout Europe and even 46
inhabited the British Isles until the tenth century. However, 47
since the mid-1800's, populations in Europe have been 48
severely reduced due to habitat destruction and overexplo- 49
itation by humans (Servheen 1990). Four very small, isolated 50
populations, all of which are endangered, can still be found 51
in southern and western Europe (the Pyrenees, Southern 52
Alps, Cantabrian Mountains and Apennine Mountains), 53
representing the remnants of a once widespread brown bear 54
population (Zedrosser et al. 2001). The Cantabrian popula- 55
tion is found in two areas of the Spanish Cantabrian 56
Mountains separated by 30–50 km of mountainous terrain. 57
Interchange between both subpopulations is unlikely (Naves 58
and Nores 1997), due to unsuitable habitat and a high speed 59
railway and motorway that bisect the area (Fig. 1). The 60
population estimate for the western subpopulation is around 61

A1 T. Pérez (✉) · F. Vázquez · A. Corao · J. Albornoz ·
A2 A. Domínguez
A3 Departamento de Biología Funcional (Genética), Facultad de
A4 Medicina 7ª Planta, Universidad de Oviedo, C/Julián Clavería
A5 s/n, 33071 Oviedo, Spain
A6 e-mail: mtpmendez@yahoo.es

A7 J. Naves · A. Fernández
A8 Departamento de Biología de Organismos y Sistemas (Ecología),
A9 Universidad de Oviedo, C/Catedrático Rodrigo Uría s/n, 33071
A10 Oviedo, Spain

A11 J. Naves · A. Fernández
A12 Department of Applied Biology, Estación Biológica de Doñana,
A13 Consejo Superior de Investigaciones Científicas, Avda. de María
A14 Luisa s/n, Pabellón del Perú, 41013 Sevilla, Spain

Fig. 1 Distribution map of the brown bear in Europe: (a) present distribution of the brown bear in Europe, and (b) distribution of the brown bear in the Cantabrian Mountains (based on Naves et al. 2003) and location of the 133 samples analysed in this study. White squares represent samples amplified for more than 14 markers, black squares represent samples with less than 14 loci amplified



50–60 individuals (Wiegand et al. 1998). The eastern Cantabrian Mountains subpopulation is estimated to contain around 20 bears (Naves et al. 1999). Both Cantabrian bear populations face similar conservation problems.

Mitochondrial DNA studies have shown that the Cantabrian population belongs to the Iberian refugia clade of the western lineage of European brown bears. Only the populations of the Cantabrian and Pyrenean Mountains and small populations from the south of Sweden and Norway belong to this clade (Taberlet and Bouvet 1994). It is important to preserve this population if we wish to maintain the most ancient lineage of the European brown bear.

In order to design an effective conservation plan, it is necessary to understand the structure of threatened populations, particularly those which, like this one, exist in degraded or fragmented habitats (Lande and Barrowclough 1987; Simberloff 1988; Hanski and Gilpin 1997; Taylor and Dizon 1999; Kraaijeveld-Smith et al. 2005). It is well known that small populations are more vulnerable to genetic factors, demographic and environmental stochasticity, genetic drift and inbreeding and have an increased probability of extinction (Soulé 1987). When small populations become fragmented and migration between subpopulations decreases or is eliminated, consequent increases in inbreeding and loss of genetic diversity can have serious negative effects on the long-term viability of population fragments and, by extension, of the population as a whole (Keller et al. 1994; Lacy 1997; Bjijsma et al. 2000; Sherwin and Moritz 2000; Coulon et al. 2004).

To avoid the extinction of this species in the Cantabrian Mountains, the different regional governments from the areas where it is present have developed plans for the conservation of brown bears since 1989. These plans have been coordinated at a national level. In order to be effective, however, an action plan should be based on reliable biological data, such as trustworthy estimates of population size, population genetic status and connectivity with other populations (Bellemain et al. 2007).

To assess the genetic status of the Cantabrian population and provide guidelines for the conservation and management of this population, we used non-invasive genetic techniques which have been successfully used in other studies (Taberlet et al. 1997; Kohn et al. 1999; Woods et al. 1999; Waits et al. 2000; Frantz et al. 2003; Smith et al. 2006; Bellemain et al. 2007). However, this non-invasive approach has a major drawback, namely the problems associated with low DNA quantity and quality (Taberlet and Luikart 1999). Under these circumstances, the probability of critical genotyping errors (primarily allelic dropout and false alleles) is high (Taberlet and Luikart 1999; Taberlet et al. 1996, 1999). The solutions proposed in these studies for addressing and reducing the severity of genotyping errors and increasing the reliability of genotypes include protocols for replicating amplification and programmes for determining the magnitude of error in a dataset and for calculating the reliability of genotype data (Taberlet et al. 1996; Miller et al. 2002; Bonin et al. 2004; McKelvey and Schwartz 2004).

120 The goal of the present study was to ascertain the current
 121 genetic status of the Cantabrian brown bear population
 122 using non-invasive genotyping techniques. The results of
 123 this work will help us to provide guidelines for the con-
 124 servation and management of this population. To achieve
 125 this goal, we optimized a set of suitable loci microsatellite
 126 markers for carrying out population and individual identi-
 127 fication studies in this population.

128 **Methods**

129 **Sampling**

130 This study presents the genetic data for the Cantabrian
 131 brown bear population using non-invasive samples (faeces
 132 and hairs) collected in the field between 2004 and 2006
 133 using two sampling methods: opportunistic and systematic.
 134 Most of the samples ($n = 106$) were collected in an
 135 opportunistic manner from daily routine field work of
 136 rangers and field biologists. Some samples ($n = 27$) were
 137 collected through systematic surveys carried out seasonally
 138 (summer: mid August–mid September; autumn: mid
 139 October–mid November) over a grid (2.5 km^2) in the
 140 western subpopulation in an area of roughly 750 km^2
 141 covering around 630 km of on-foot surveys each season.
 142 Although we were mainly looking for scats, some hair
 143 samples were also collected. Since we wished to cover the
 144 maximum area of distribution possible, we had to include
 145 many opportunistic samples because the systematic surveys
 146 only cover a partial area of the species distribution.
 147 Figure 1b shows where the samples were collected.

148 Faeces samples were dry stored with silica after ethanol
 149 soaking upon collection following Nsubuga et al. (2004)
 150 and Roeder et al. (2004). Hairs were placed in individual
 151 envelopes, no further manipulation being necessary until
 152 extraction. A total of 133 samples were analysed: 88 from
 153 the western subpopulation and 45 from the eastern sub-
 154 population. Together with these non-invasive samples, 13
 155 tissue/blood samples belonging to either dead or captured
 156 individuals were used (all from the western subpopulation,
 157 eight of which were found dead between 1989 and 2002).
 158 The numbers from each type of sample used for the study
 159 are listed in Table 1.

160 **DNA extraction and typing**

161 *DNA extraction*

162 DNA was extracted from faeces samples using the ‘Qiamp
 163 DNA stool kit’ (Qiagen, Hilden, Germany) specially
 164 developed for this type of material and following the
 165 manufacturer’s instructions. The only change introduced

Table 1 Number of samples analysed and percent of the total (between parentheses)

	Faeces	Hair	Skin/ blood	Total
No. DNA	25 (27.2%)	0 (0%)	1 (7.7%)	26 (17.8%)
Discarded genotypes ^a	27 (29.3%)	6 (14.6%)	3 (23.1%)	36 (24.7%)
Accepted genotypes ^b	40 (43.5%)	35 (85.4%)	9 (69.2%)	84 (57.5%)
Total number samples	92	41	13	146

^a Include all the incomplete genotypes (with less than 14 loci amplified)

^b Genotypes with more than 14 loci amplified (all the hair, skin/blood samples and 24 of the faeces samples in this category were genotyped for the 18 loci)

166 was the incubation of the samples in ASL buffer ON at
 167 25°C . For hair DNA extraction, 5–10 hair roots were used;
 168 these were incubated ON with PCR buffer and proteinase
 169 K (Allen et al. 1998; Vigilant 1999). Finally, tissue/blood
 170 samples were extracted using the ‘DNeasy Tissue kit’
 171 (Qiagen, Hilden, Germany) following the manufacturer’s
 172 instructions. All hair and faeces extractions were carried
 173 out in a separate room used exclusively for processing
 174 samples of this type.

175 *Genotyping*

176 For the genotyping, we used a total of 18 microsatellite
 177 markers (G1A, G1D, G10B, G10C, G10J, G10L, G10O,
 178 G10P, G10X (Paetkau and Strobeck 1994; Paetkau et al.
 179 1995) and MU05, MU09, MU10, MU23, MU50, MU51,
 180 MU59, MU61, MU64 (Taberlet et al. 1997)), as well as the
 181 sex marker SRY (Bellemain and Taberlet 2004), all of
 182 which were selected from markers previously used in
 183 genetic tracking of Pyrenean brown bears (Taberlet et al.
 184 1997).

185 DNA amplifications were performed in a two-step PCR
 186 following Taberlet et al. (1997) for faeces samples and
 187 using only one PCR for hair and tissue/blood samples. To
 188 avoid the problem of running out of DNA template
 189 (because of the low amount of DNA available) before
 190 completing the genotyping for all the microsatellite
 191 markers, the amplification was carried out following the
 192 multiplex preamplification method (Piggot et al. 2004;
 193 Bellemain and Taberlet 2004). This method was optimized
 194 for six microsatellite loci and for the sex marker (MU10,
 195 MU23, MU50, MU51, MU59, G10L and SRY) by Belle-
 196 main and Taberlet (2004). We designed three new
 197 multiplex PCRs to amplify all the remaining loci
 198 (MU64 + G1A + G10C + G10P, MU61 + G10J + -
 199 G10O + G10X, MU05 + MU09 + G1D + G10B). A

Author Proof

total number of four-first-step PCRs, where we amplified between 4 and 6 loci simultaneously, and nine-second-step PCRs, where the number of loci amplified simultaneously varied between 1 and 3, were sufficient to amplify the 18 loci microsatellites and the sex marker (Table 2).

For faeces samples, the first-step PCRs or preamplifications were prepared in a 25 µl volume containing 5 µl template DNA, 0.01 µM of each primer and 12.5 µl of “Qiagen Multiplex PCR Kit” (Qiagen, Hilden, Germany). The second-step, PCRs or amplifications were prepared in a 13 µl volume containing 3 µl preamplified product, 0.1 mM of each dNTP, 0.5 µM of each primer, 2 mM MgCl₂, 0.5 U *Taq* DNA Polymerase from Qiagen (Qiagen, Hilden, Germany) and 1× *Taq* Quiagen buffer (Qiagen, Hilden, Germany). Amplifications were performed on a GeneAmp PCR 9600 (Applied Biosystems) under the following conditions: for the first-step PCR 15 min at 95°C, 40 cycles composed of 30 s denaturing at 94°C, 90 s annealing at 60°C, 1 min extension at 72°C, and as a final extension step, 30 min at 60°C. For the second-step PCR, 3 min at 94°C, 35 cycles composed of 30 s denaturing at 94°C, 30 s annealing at 60°C, 1 min extension at 72°C, and as a final extension step, 7 min at 72°C.

For hair and tissue/blood samples, nine-one-step PCRs were carried out using the primers of the second-step PCRs in a 20 µl volume containing 5 µl (2 µl for tissue/blood samples) template DNA, 0.5 µM of each primer and 10 µl of “Qiagen Multiplex PCR Kit” (Qiagen, Hilden, Germany). Amplifications were performed on a GeneAmp

PCR 9600 (Applied Biosystems) under the following conditions: 15 min at 95°C, 40 cycles composed of 30 s denaturing at 94°C, 90 s annealing at 60°C, 1 min extension at 72°C, and as a final extension step, 30 min at 60°C.

PCR products were checked in a 2% agarose gel and the product diluted between 0 and 100 times depending on the intensity of the signal. One microlitre of this product was added to a 12 µl mix of formamide and ROX 400HD (12:0.2) and then loaded on an automatic sequencer ABI310 (Applied Biosystems). Microsatellite patterns were examined both visually and using GENESCAN ANALYSIS 3.1 and GENOTYPER 2.5 software (Applied Biosystems).

Reliability of genotyping results

To test the reliability of our genotyping results and to reduce tracking error in our dataset, we followed different recommendations already proposed in different studies addressing non-invasive genotyping. The suggestions by Bonin et al. (2004) for limiting potential errors in the genotyping process were followed. All the genetic typing was performed using a combination of the multi-tube approach and software packages that assign a reliability value to each multilocus genotype. Three positive PCRs (for both types of non-invasive samples, hair and faeces) were first analysed, a consensus genotype was assigned using the GIMLET v.1.3.2 software (Valière 2002) and its reliability was tested using the RELIOTYPE software

Table 2 Observed number of alleles (*A*) with the number of single alleles of each subpopulation between parentheses (*Ae*), observed heterozygosity (*H_o*), Nei’s estimated heterozygosity (*H_e*) and deviations from Hardy–Weinberg equilibrium (HWE) by locus for each subpopulation with sample size in parentheses

Locus (i,j)	Eastern subpopulation (n = 8)			Western subpopulation (n = 39)		
	<i>A</i> (<i>Ae</i>)	<i>H_e</i>	<i>H_o</i>	<i>A</i> (<i>Ae</i>)	<i>H_e</i>	<i>H_o</i>
MU10 (1,1)	2 (0)	0.50	0.50	4 (2)	0.42	0.28*
G10L (1,1)	1 (0)	0.00	0.00	6 (5)	0.74	0.64*
U50 (1,2)	1 (0)	0.00	0.00	5 (4)	0.67	0.69
MU23 (1,2)	3 (1)	0.63	0.88	3 (1)	0.65	0.72
MU59 (1,3)	1 (0)	0.00	0.00	4 (3)	0.56	0.59
MU51 (1,3)	2 (0)	0.49	0.63	4 (2)	0.37	0.38
G10C (2,1)	1 (0)	0.00	0.00	3 (2)	0.50	0.41
MU64 (2,1)	1 (1)	0.00	0.00	3 (3)	0.46	0.33*
G1A (2,2)	2 (1)	0.30	0.38	3 (2)	0.51	0.64
G10P (2,2)	2 (1)	0.22	0.25	2 (1)	0.45	0.49
G10J (3,1)	2 (0)	0.38	0.50	3 (1)	0.65	0.69
G10X (3,2)	2 (1)	0.49	0.13*	4 (3)	0.29	0.26
MU61 (3,2)	1 (0)	0.00	0.00	2 (1)	0.44	0.49
G10O (3,2)	1 (0)	0.00	0.00	1 (0)	0.00	0.00
MU05 (4,1)	3 (1)	0.63	0.75	4 (2)	0.58	0.56
G1D (4,1)	2 (1)	0.30	0.38	1 (0)	0.00	0.00
MU09 (4,2)	2 (1)	0.50	0.75	4 (3)	0.66	0.51*
G10B (4,2)	1 (0)	0.00	0.00	3 (2)	0.17	0.13
Mean	1.67	0.25	0.28	3.33	0.45	0.44

Next to the locus name, between parentheses, the number of the preamplification (i) and amplification (j) PCR

**P*-value significant *P* < 0.05

256 (Miller et al. 2002). If we found a multilocus genotype
 257 with reliability lower than 95%, more repetitions were
 258 carried out until achieving said level of reliability. All the
 259 samples that could not be reliably typed for at least 14 out
 260 of the 18 loci after the entire process was completed were
 261 discarded. To further identify any genotyping errors and
 262 the relative magnitude of a problem within our multilocus
 263 scores, we performed the tests proposed by McKelvey and
 264 Schwartz (2004): Examining Bimodality (EB) and Differ-
 265 ence in Capture History (DCH). Finally, we determined
 266 genotypic mismatches between all scores.

267 Probability of identity

268 Using the software GIMLET version 1.3.2 (Valière 2002),
 269 we computed the probability of identity (P_{ID}), which is the
 270 probability that two individuals drawn at random from a
 271 given population share identical genotypes at all typed loci
 272 (Paetkau and Strobeck 1994). We also computed the
 273 probability of identity among siblings (P_{ID-Sib}) (Waits et al.
 274 2001). This value is the upper limit of the possible ranges
 275 for the probability of identity in a population and thus
 276 provides the most conservative number of loci required to
 277 resolve all bears, including relatives. These calculations
 278 were carried out for each subpopulation.

279 Population genetic parameters and structure

280 We ran population genetic analyses using the software
 281 programmes GENEPOP version 3.4 (Raymond and Rous-
 282 set 1995), GENETIX version 4.02 (Belkhir et al. 1996–
 283 2004) and STRUCTURE version 2.1 (Pritchard et al.
 284 2000). Nuclear genetic diversity was measured as the
 285 number of alleles per locus (A), the observed heterozy-
 286 gosity (H_o), as well as Nei’s unbiased expected
 287 heterozygosity (H_e) (Nei 1978). Deviations from Hardy–
 288 Weinberg equilibrium were tested using an exact test.
 289 Global tests across loci for heterozygote deficiency and
 290 heterozygote excess and pairwise tests for linkage dis-
 291 equilibrium were performed using Fisher’s method (Sokal
 292 and Rohlf 1994) with 10,000 batches and 10,000 iterations
 293 per batch. Correspondence among individual genotypes
 294 was studied by means of Factorial Correspondence Anal-
 295 ysis (FCA), performed with the GENETIX software.
 296 Population substructure was detected with the programme
 297 STRUCTURE, which uses a Markov chain Monte Carlo
 298 (MCMC) algorithm to cluster individuals into populations
 299 on the basis of multilocus genotype data (Pritchard et al.
 300 2000). We used different values of K , from one to five. For
 301 each K tested, we ran STRUCTURE 20 times for 100,000
 302 steps, after a burn-in period of 50,000 steps. The correct
 303 value of K was estimated following Evanno et al. (2005).
 304 The programme also calculates the fractional membership

of each individual in each cluster (Q). Quantification of 305
 variation among subpopulations was performed with 306
 GENEPOP using Wright’s F -statistics (Weir 1996). 307

Results 308

Reliability of non-invasive genotyping 309
 for the identification of the Cantabrian bear 310

A total number of 133 non-invasive samples were analysed 311
 (92 faeces samples and 41 hair samples). We managed to 312
 obtain some amplification for all the hair samples and for 313
 67 faeces samples (72.8%), but only those samples suc- 314
 cessfully amplified for more than 14 markers were included 315
 in the analysis (Table 1). Another nine genotypes were 316
 obtained from tissue/blood samples. For the genotyping, 317
 we initially tested 24 loci previously used in genetic 318
 tracking of Pyrenean bears (Taberlet et al. 1997; Bellemain 319
 and Taberlet 2004). Six out these 24 loci did not give a 320
 scorable product and so were discarded. Finally, we used 321
 the remaining 18 loci and the sex marker; all the loci were 322
 amplified using multiplex PCRs, three of which were 323
 specifically developed for this study (Table 2). The prob- 324
 ability of identity values were 3.28E09 (P_{ID}) and 1.16E04 325
 (P_{ID-Sib}) for the western subpopulation and 7.45E05 and 326
 8.64E03 for the eastern subpopulation. The P_{ID} and P_{ID-Sib} 327
 values for each marker and subpopulation are shown in 328
 Table 3. 329

Once we had completed three repetitions per sample, we 330
 found that out of the 67 faeces that gave a product, only 40 331
 (59.7%) worked for at least 14 markers. For hair samples, 332
 35 out of 41 (85.4%) gave a complete profile. The error 333
 rates calculated using the GIMLET programme show that 334
 both the number of failed PCRs and percentage of total 335
 error are higher for faeces samples than for hair samples 336
 (see Fig. 2). These results indicate that the DNA recovered 337
 from hair samples has better quality than that recovered 338
 from faeces samples. Results from the RELIOTYPE pro- 339
 gramme showed that 76% of the samples reach 95% 340
 reliability after three repetitions, 16% needed a fourth 341
 repetition to reach this level of reliability and 8% needed 342
 more than four repetitions to reach it (one of the samples 343
 had to be repeated up to seven times to achieve 95% 344
 reliability). 345

The results from the EB test, which examines the dis- 346
 tribution of the genetic differences between samples, and 347
 the DCH test, which determines whether the number of 348
 new individuals in the sample increased faster than would 349
 be expected when additional loci are added (McKelvey and 350
 Schwartz 2004), indicated that our dataset had a low rate of 351
 genotyping error (unimodal distribution for the EB test and 352
 no locus-added new individuals for the DHC test). The 353


Author Proof

Table 3 Probability of identity (P_{ID}) and Probability of identity for siblings (P_{ID-sib}) by locus for each subpopulation with sample size in parentheses and accumulated values

Locus (i,j)	Eastern subpopulation (n = 8)		Western subpopulation (n = 39)	
	P_{ID}	P_{ID-sib}	P_{ID}	P_{ID-sib}
MU10 (1,1)	3.75E-01	5.94E-01	3.73E-01	6.34E-01
G10L (1,1)	1.00E+00	1.00E+00	1.10E-01	4.10E-01
MU50 (1,2)	1.00E+00	1.00E+00	1.60E-01	4.56E-01
MU23 (1,2)	2.12E-01	4.87E-01	1.93E-01	4.71E-01
MU59 (1,3)	1.00E+00	1.00E+00	2.71E-01	5.40E-01
MU51 (1,3)	3.79E-01	5.99E-01	4.32E-01	6.71E-01
G10C (2,1)	1.00E+00	1.00E+00	3.61E-01	5.89E-01
MU64 (2,1)	1.00E+00	1.00E+00	3.54E-01	6.10E-01
G1A (2,2)	5.30E-01	7.30E-01	3.58E-01	5.84E-01
G10P (2,2)	6.34E-01	7.99E-01	4.02E-01	6.24E-01
G10 J (3,1)	4.61E-01	6.78E-01	1.96E-01	4.73E-01
G10X (3,2)	3.79E-01	5.99E-01	5.18E-01	7.33E-01
MU61 (3,2)	1.00E+00	1.00E+00	4.09E-01	6.32E-01
G10O (3,2)	1.00E+00	1.00E+00	1.00E+00	1.00E+00
MU05 (4,1)	2.12E-01	4.87E-01	2.38E-01	5.20E-01
G1D (4,1)	5.30E-01	7.30E-01	1.00E+01	1.00E+01
MU09 (4,2)	3.75E-01	5.94E-01	1.82E-01	4.68E-01
G10B (4,2)	1.00E+00	1.00E+00	7.07E-01	8.44E-01
Accumulated	7.45E-05	8.64E-03	3.28E-09	1.16E-04

Next to the locus name, between parentheses, the number of the preamplification (i) and amplification (j) PCR

Fig. 2 Percentage of errors (failed PCR, dropout and other types of error) and correct genotypes depending on the non-invasive sample type: (a) hair samples, and (b) faeces samples



354 presence of a bimodal structure (Fig. 3) in the distribution
 355 of the number of loci at which individuals differed (EB
 356 test) for the eastern subpopulation is not necessarily a result
 357 of error; in this case, it indicates that one of the individuals
 358 is highly different from the rest, showing that it could be a
 359 migrant. The DHC test could not be carried out on the
 360 eastern subpopulation, since all the polymorphic loci tested
 361 are included in the genetic tag.

362 The average number of loci at which individuals differed
 363 was 10.62 ± 1.78 for the western subpopulation and
 364 7.69 ± 4.2 for the eastern, although if we remove the
 365 individual that appears to be different from the rest, the
 366 average is lower (5.68 ± 1.94).

367 A final number of 31 individuals (16 females, 15 males)
 368 out of 45 samples were identified in the western subpopu-

369 lation and 9 (5 females, 4 males) out of 30 samples in the
 370 eastern subpopulation.

371 **Microsatellite diversity and population substructure**

372 The number of alleles for the total population was 67; of
 373 these, 45 were uniquely sampled from one of the two sub-
 374 populations (8 for the eastern subpopulation and 37 for the
 375 western one). The null hypothesis of uniform allelic and
 376 genotypic frequencies in the two subpopulations was rejected
 377 at 15 out of 17 polymorphic loci. The number of alleles
 378 per locus for the western subpopulation ranged between 1
 379 and 6, with an average of 3.33 ± 1.28 and between 1 and 3
 380 for the eastern subpopulation with an average of 1.67 ± 0.67
 381 (Table 2). Two loci were monomorphic for the western




Fig. 3 Number of loci at which one individual differed from all the other individuals from the same population based on 18 loci for both populations: (a) western and (b) eastern. Only individuals with completed genotypes were used in this calculation (25 for the western subpopulation and 9 for the eastern one). All the comparisons involving one of the individuals found in the eastern subpopulation (in a circle) differed in a higher number of loci from all the rest of the individuals from the same subpopulation

382 subpopulation (G10O and G1D), but up to eight loci were
 383 monomorphic for the eastern subpopulation (G10L, Mu50,
 384 Mu59, G10C, Mu64, Mu61, G10O, and G10B). The locus
 385 G10O was monomorphic for both subpopulations. The
 386 average observed and expected heterozygosities for the total
 387 population were 0.51 and 0.43, respectively, and were sig-
 388 nificantly different ($P < 0.001$). The partition of genetic
 389 diversity among subpopulations was 42.96%; the values of
 390 diversity found for each subpopulation were 0.45 for the
 391 western subpopulation and 0.25 for the eastern one
 392 (Table 2).

393 Global tests showed that both subpopulations are in
 394 Hardy–Weinberg equilibrium, although four loci (Mu10,
 395 G10L, Mu64, Mu09) for the western subpopulation and
 396 one locus (G10X) for the eastern subpopulation had a
 397 significant deficiency in heterozygotes at the $P < 0.05$
 398 level (Table 2). Statistical tests for linkage disequilibrium
 399 were computed for all pairs of loci, though none of these
 400 were significant.

401 The canonical analysis based on factorial correspon-
 402 dence analysis (FCA) divided all the samples in two clearly

403 differentiated groups (Fig. 4). Each group corresponded
 404 with the two subpopulations present in the Cantabrian
 405 Mountains (western and eastern subpopulations). One
 406 individual that was detected in samples collected in the
 407 eastern area groups together with the individuals from the
 408 western subpopulation. This is the same individual that
 409 showed a strange pattern when we checked the distribution
 410 of the number of loci at which two individuals differed for
 411 the eastern subpopulation (Fig. 3). Furthermore, the pro-
 412 gramme STRUCTURE detected that Cantabrian brown
 413 bears are structured in two subpopulations which corre-
 414 spond with the eastern–western groupings (Fig. 4). The
 415 mean Q value (cluster membership) is 0.998 for the wes-
 416 tern subpopulation and 0.888 for the eastern one. Once
 417 more, one individual is detected in the eastern subpopula-
 418 tion that is more similar to the individuals from the western
 419 group. The Q value of this individual is 0.010 for the
 420 eastern subpopulation and 0.998 for the western one,
 421 indicating that this individual (a male) is a migrant. If we
 422 remove this individual, the mean Q for the eastern sub-
 423 population is 0.998. For all the aforementioned reasons,
 424 this individual was removed from the basic statistical cal-
 425 culations regarding population differentiation.

426 The degree of genetic differentiation between the two
 427 subpopulations was considerable. The proportion of vari-
 428 ation in genetic frequencies, θ (F_{st}), accounted for by
 429 subdivision was 0.41, while Nei’s standard distance
 430 between both subpopulations was 0.47.

Discussion


Quality of the genetic data

431
 432
 433 In order to maximize the success of the non-invasive
 434 sample DNA extracts, we used storage and extraction
 435 techniques that have previously shown their effectiveness.
 436 We obtained 72.8% amplification success for faeces and
 437 100% for hairs; these values are in the range of values
 438 described in the literature (see Nsubuga et al. 2004; Chu
 439 et al. 2006). However, these success values dropped to
 440 43.48% for faeces and 85.37% for hairs when we consid-
 441 ered only those samples that gave us a multilocus profile
 442 for at least 14 out the 18 loci used.

443 Taberlet et al. (1999) suggest that non-invasive studies
 444 should include a pilot study to assess the confidence level
 445 of the final result by quantifying the genotyping error rate.
 446 The results of studies of this type are unique and cannot be
 447 transferred to another species or even to another population
 448 with different heterozygosity or sample quality. By com-
 449 bining several methodological and statistical methods for
 450 tracking and reducing error previously used in different
 451 non-invasive studies (Frantz et al. 2003; Bellemain and

Author Proof

Fig. 4 (a) Distribution of individuals according to genotype based on factorial correspondence analysis (FCA), and (b) Graphical representation of the STRUCTURE programme. In both cases individuals belonging to the eastern and western populations are indicated in white and black, respectively



452 Taberlet 2004; Smith et al. 2006), we were able to thor- 477
 453 oughly assess the level of error in our data set (2.55% for 478
 454 hair samples and 5.21% for faeces samples). These error 479
 455 values are similar to the lowest found in the literature for 480
 456 these types of samples (Bayes et al. 2000; Bellemain et al. 481
 457 2004; Smith et al. 2006). The reliability of the multilocus 482
 458 genotypes after three repeats was quite high (76% of the 483
 459 samples had >95% reliability). A fourth repetition increas- 484
 460 es this percentage up to 92%, which is in concordance 485
 461 with the result previously presented by Bellemain and 486
 462 Taberlet (2004) in the brown bear. 487

463 The low diversity found in the Cantabrian population, 488
 464 mainly in the eastern subpopulation, together with the fact 489
 465 that the most informative loci are not the same in the two 490
 466 subpopulations, makes it necessary to use a high number of 491
 467 markers if we wish to unambiguously identify individuals 492
 468 for future demographic studies. 493

469 Genetic status of the bear population in the Cantabrian 494 470 Mountains 495

471 The genetic diversity of the Cantabrian population is lower 498
 472 than in other bear populations considered to have a good 499
 473 conservation status (Paetkau et al. 1998; Waits et al. 1998) 500
 474 and is comparable with those found in Yellowstone 501
 475 ($H_e = 0.55$; Paetkau et al. 1998) and Deosai National Park 502
 476 (Pakistan) ($H_e = 0.55$; Bellemain et al. 2007). Yellowstone 503

and Cantabrian populations have had a very similar history. 477
 Both populations have gone from being embedded in a very 478
 large continuous population to being an isolated remnant, 479
 separated from other brown bears for 300 years in the case of 480
 the Cantabrian population (Naves et al. 1999) and 100 years 481
 for the Yellowstone population (Paetkau et al. 1998). In both 482
 cases, there is no prospect of renewed connections with other 483
 populations. The low heterozygosity values can be explained 484
 either by a founder effect, which is not the case for the 485
 Cantabrian brown bear, or for a sharp decline in population 486
 size. The whole population probably began to lose genetic 487
 diversity about 300 years ago, when it began to decline in 488
 size from a larger population. 489

The brown bear habitat in the Cantabrian Mountains has 490
 decreased considerably from $\sim 9,000 \text{ km}^2$ at the turn of the 491
 twentieth century to $\sim 5,000 \text{ km}^2$ at present (Naves and 492
 Nores 1997), while the population subdivided into two 493
 apparently isolated subpopulations, the western and the 494
 eastern. Genetic differentiation ($F_{st} = 0.41$, Nei's standard 495
 $D = 0.47$) between the two subpopulations was found to 496
 be very high. These values can be compared with the 497
 reported microsatellite-based estimates of differentiation 498
 among other bear populations. For example, the degree of 499
 genetic differentiation is considerably higher than the 500
 values reported for pairwise comparisons among subpop- 501
 ulations in Scandinavia, where F_{st} ranged between 0.01 and 502
 0.14 and Nei's standard distance ranged between 0.03 and 503

504 0.38. The genetic distance between the two Cantabrian
505 subpopulations, which are only 30 km apart, are compa-
506 rable with the values reported for the most distant areas
507 within the continuous distribution of brown bears in North
508 America, which are several thousands of kilometres apart
509 (Paetkau et al. 1998). This result also shows that habitat
510 discontinuities such as roads and farmland play a larger
511 role in genetic substructuring of population than linear
512 distance, which has also been seen in populations of Cross
513 River gorilla (Bergl and Vigilant 2007).

514 The huge genetic differentiation shown between Canta-
515 brian subpopulations may be related to total isolation
516 between them in conjunction with an extremely low popu-
517 lation size in the eastern subpopulation. Diversity in this
518 subpopulation is among the lowest found in the literature and
519 can only be compared with the value reported in the isolated
520 population of the Kodiak islands in Alaska ($H_e = 0.26$;
521 Paetkau et al. 1998). The eastern subpopulation, with a
522 diversity of 0.25, was detached from the main group a few
523 generations ago. Assuming that diversity at the time of the
524 split was comparable with the diversity of the western sub-
525 population, the relative loss of diversity in the eastern
526 subpopulation ($H_{\text{subpopulation}}/H_{\text{initial}}$) is 0.56. This leads to a
527 fixation index ($1 - H_{\text{subpopulation}}/H_{\text{initial}}$) equal to 0.44, quite
528 close to the estimated value of F_{st} . If we assume a continent-
529 island model (given that the relative effect of drift in the
530 western subpopulation is low) with no migration,
531 $F_{st}=1 - (1/2N_e)^t$, where N_e is the effective size and t the
532 time in generations. Considering that both subpopulations
533 have been isolated for 10 generations (50–75 years with a
534 generation time for the brown bear of 10–15 years, Allen-
535 dorf and Servheen 1986; Craighead et al. 1995), the N_e for
536 the eastern subpopulation that would explain the obtained F_{st}
537 would be 4.99 individuals per generation. These figures
538 show that the two subpopulations have probably been totally
539 isolated without any effective migration during the last few
540 generations. Although the data point towards a total lack of
541 genetic flow between the two subpopulations, a migrant
542 male from the western subpopulation into the eastern
543 grouping was identified. This could either be interpreted as
544 the first signal that connectivity between both subpopula-
545 tions is starting to occur after this long period of isolation, or
546 as the existence of a certain rate of migration, though not
547 effective gene flow. It will be necessary to check in the future
548 whether more migrants are present in the population and
549 whether or not this migration is effective in terms of genetic
550 flux. The main genetic consequence of migration would be a
551 drop in genetic differentiation between the two subpopula-
552 tions of the Cantabrian brown bear. If we once more assume
553 a continent–island model, the F_{ST} at equilibrium between
554 gene flow due to migration and genetic drift due to the small
555 population size is $\hat{F}_{ST} \approx 1/(4N_{em} + 1)$, where N_e is the
556 effective size and m is the migration rate per generation. This

implies that with one effective migrant per generation 557
($N_{em} = 1$), the equilibrium F_{ST} is 0.20, and 0.11 should there 558
be two effective migrants per generation. That would be 559
sufficient to prevent the huge effects of genetic drift in the 560
oriental subpopulation, despite its reduced population size. 561

562 Conclusions and recommendations 562

563 The results of this study show that the population of 563
Cantabrian brown bears is effectively split into two sub- 564
populations with a very high level of differentiation. 565
Applying this unique criteria, it could be thought that they 566
should be treated as separate management units (MU) 567
(Moritz 1994) However, bearing in mind their contiguous 568
distribution range, the reduced population size of both 569
subpopulations and the real possibility of migration, these 570
two subpopulations need to be managed as a unique unit. 571
The rate of inbreeding per generation in the eastern sub- 572
population is around 10%, a value far exceeding the 573
maximum tolerable rate of 1% given for domestic animals 574
(Franklin 1980). Therefore, connectivity of both subpopu- 575
lations should be the highest priority if we wish to maintain 576
the diversity afforded by the eastern nucleus which is in 577
risk of immediate extinction. 578

579 The western subpopulation shows a moderate level of 579
diversity in the lowest range of values found in the species, 580
probably due to a sharp decline in population size that began 581
around 300 years ago, when it was isolated from a larger 582
population. A population size of 50–60 individuals was 583
estimated for this nucleus on the basis of the number of 584
females with cubs (Wiegand et al. 1998). It was suggested 585
(Franklin 1980) that the minimum effective size for a popu- 586
lation to be viable in the short-term should be 50. Bearing in 587
mind that the ratio of N_e/N found in the brown bear from 588
Yellowstone was 0.27 (Miller and Waits 2003) and using this 589
value in our case, the minimum size of population to be 590
viable in the short-term is ~ 186 individuals, and even the 591
western subpopulation is far from this number. 592

593 If the whole population is far from the minimum number 593
to be viable over the short-term, an even more difficult 594
situation can be depicted in the context of long-term pro- 595
tection of adaptive potential where the minimum 596
effective population size that has been put forward is 597
between 500 and 5,000 (Franklin 1980; Frankham and 598
Franklin 1998; Lynch and Lande 1998). It will be neces- 599
sary to monitor the whole population to obtain a more 600
accurate estimate of population size and its trend. Habitat 601
loss and human-caused mortality should be avoided in 602
order to facilitate an increase in population size. 603

604 **Acknowledgements** This work was funded by Grant CN-05-030 604
from the “Consejería de Medio Ambiente y Ordenación del Territorio 605
e Infraestructuras del Principado de Asturias” We wish to thank the 606

- 607 “Junta de Castilla y León” for their economic support and for providing some of the samples; Miguel Rico, Juan Seijas and the
608 “Guardería Rural del Principado de Asturias”, who also provided
609 samples; Dr. Lissette Waits and the two anonymous reviewers who
610 helped to improve this manuscript with their valuable comments, and
611 Paul Barnes, who thoroughly reviewed the English.
612
- 613 **References**
- 614 Allen M, Engström AS, Meyers S et al (1998) Mitochondrial DNA
615 sequencing of shed hairs and saliva on robbery caps: sensitivity
616 and matching probabilities. *J Forensic Sci* 43:453–464
- 617 Allendorf FW, Servheen C (1986) Genetics and the conservation of
618 grizzly bears. *Trends Ecol Evol* 1:88–89
- 619 Bayes MK, Smith KL, Alberts SC, Brudford MW (2000) Testing the
620 reliability of microsatellite typing from faecal DNA in the
621 savannah baboon. *Conserv Genet* 1:173–176
- 622 Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F (1996–2004)
623 GENETIX 4.05, logiciel sous Windows TM pour la génétique
624 des populations. Laboratoire Génome, Populations, Interactions,
625 CNRS UMR 5000, Université de Montpellier II, Montpellier
626 (France)
- 627 Bellemain E, Taberlet P (2004) Improved noninvasive genotyping
628 method: application to brown bear (*Ursus arctos*) faeces. *Mol*
629 *Ecol Notes* 4:519–522
- 630 Bellemain E, Nawaz MA, Valentini A, Swenson JE, Taberlet P (2007)
631 Genetic tracking of the brown bear in northern Pakistan and
632 implications for conservation. *Biol Conserv* 134:537–547
- 633 Bergl RA, Vigilant L (2007) Genetic analysis reveals population
634 structure and recent migration within the highly fragmented
635 range of the Cross River gorilla (*Gorilla gorilla diehli*). *Mol*
636 *Ecol* 16:501–516
- 637 Bjilmsa R, Bundgaard J, Boerema A (2000) Does inbreeding affect
638 the extinction risk of small population? Predictions from
639 *Drosophila*. *J Evol Biol* 13:502–514
- 640 Bonin A, Bellemain E, Eidesen PB et al (2004) How to track and
641 assess genotyping errors in population genetics studies. *Mol Ecol*
642 13:3261–3273
- 643 Chu JH, Lin YS, Wu HY (2006) Applicability of non-invasive
644 sampling in population genetic study of Taiwanese macaques
645 (*Macaca cyclopis*). *Taiwania* 51:258–265
- 646 Coulon A, Cosson JF, Angibault JM et al (2004) Landscape
647 connectivity influences gene flow in a roe deer population
648 inhabiting a fragmented landscape: an individual-based
649 approach. *Mol Ecol* 13:2841–2850
- 650 Craighead L, Paetkau D, Reynolds HV, Vyse ER, Strobeck C (1995)
651 Microsatellite analysis of paternity and reproduction in Artic
652 grizzly bears. *J Hered* 86:255–261
- 653 Evanno G, Regnaut S, Goudet J (2005) Detecting the number of
654 clusters of individuals using the software STRUCTURE: a
655 simulation study. *Mol Ecol* 14:2611–2620
- 656 Flagstad Ø, Hedmark E, Landa A et al (2004) Colonization history
657 and noninvasive monitoring of a reestablished wolverine pop-
658 ulation. *Conserv Biol* 18:676–688
- 659 Frankham R (1995) Effective population size/adult population size
660 ratios in wildlife: a review. *Genet Res* 66:95–107
- 661 Frankham R, Franklin IR (1998) Response to Lynch and Lande. *Anim*
662 *Conserv* 1:73
- 663 Franklin IR (1980) Evolutionary change in small populations. In: Soulé
664 ME, Wilcox BA (eds) Conservation biology: an evolutionary-
665 ecological perspective. Sinauer, Sunderland, MA, pp 135–150
- 666 Frantz AC, Pope LC, Carpenter PJ et al (2003) Reliable microsatellite
667 genotyping of the Eurasian badger (*Meles meles*) using faecal
668 DNA. *Mol Ecol* 12:1649–1661
- Hanski IA, Gilpin M (1997) Metapopulation biology: ecology and
evolution. Academic Press, Toronto
- Keller L, Arcese P, Smith J, Hochachka WM, Stearns SC (1994)
Selection against inbred song sparrows during a natural popu-
lation bottleneck. *Nature* 372:356–357
- Kohn MH, York EC, Kamradt DA et al (1999) Estimating population
size by genotyping faeces. *Proc R Soc Lond B* 266:657–663
- Kraaijeveld-Smith FLJ, Beebee TJC, Griffiths RA, Moore RD,
Schley L (2005) Low gene flow but high genetic diversity in
the threatened Mallorcan midwife toad *Alytes muletensis*. *Mol*
Ecol 14:3307–3315
- Lacy RC (1997) The importance of genetic variation to the viability
of mammalian populations. *J Mammal* 78:320–335
- Lande R, Barrowclough G (1987) Effective population size, genetic
variation and their use in population management. In: Soulé ME
(ed) Viable populations for conservation. Cambridge University
Press, New York, pp 87–123
- Lynch M, Lande R (1998) The critical effective size for a genetically
secure population. *Anim Conserv* 1:70–72
- McKelvey KS, Schwartz MK (2004) Genetic errors associated with
population estimation using non-invasive molecular tagging:
problems and new solutions. *J Wildl Manage* 68:439–448
- Miller CR, Waits LP (2003) The history of effective population size
and genetic diversity in the Yellowstone grizzly (*Ursus arctos*):
implication for conservation. *Proc Natl Acad Sci USA* 7:4334–
4339
- Miller CR, Joyce P, Waits LP (2002) Assessing allelic drop-out and
genotype reliability using maximum likelihood. *Genetics*
160:357–366
- Moritz C (1994) Defining ‘Evolutionary significant units’ for
conservation. *Trends Ecol Evol* 9:373–375
- Naves J, Nores C (1997) Status of the brown bear in western
Cantabria, Spain. In: Servheen C, Herrero S, Peyton B (eds)
Bears: status survey, conservation action plan. International
Union for the Conservation of Nature, Natural Resources. Gland,
Switzerland, pp 104–111
- Naves J, Wiegand T, Fernandez A, Stephan T (1999) Riesgo de
extinción del oso pardo cantábrico La población occidental.
Fundación Oso de Asturias, Oviedo, Spain
- Naves J, Wiegand T, Revilla E, Delibes M (2003) Endangered species
constrained by natural and human factors: the case of brown
bears in northern Spain. *Conserv Biol* 17:1276–1286
- Nei M (1978) Estimation of average heterozygosity and genetic
distance from a small number of individuals. *Genetics* 89:
583–590
- Nsubuga AM, Robbins MM, Roeder AD et al (2004) Factors affecting
the amount of genomic DNA extracted from ape faeces and the
identification of an improved sample storage method. *Mol Ecol*
13:2089–2094
- Paetkau D, Strobeck C (1994) Microsatellite analysis of genetic
variation in black bear populations. *Mol Ecol* 3:489–495
- Paetkau D, Calvert W, Stirling I, Strobeck C (1995) Microsatellite
analysis of population structure in Canadian polar bears. *Mol*
Ecol 4:347–354
- Paetkau D, Waits LP, Clarkson P et al (1998) Variation in genetic
diversity across the range of North American brown bears.
Conserv Biol 12:418–429
- Piggot M, Bellemain E, Taberlet P, Taylor AC (2004) A multiplex
pre-amplification method that significantly improves microsat-
ellite amplification and error rates for faecal DNA in limiting
conditions. *Conserv Genet* 5:417–420
- Pritchard JK, Stephens P, Donnelly P (2000) Inference of population
structure using multilocus genotype data. *Genetics* 155:945–959
- Raymond M, Rousset F (1995) GENEPOP (version 1.2): population
genetics software for exact tests and ecumenicism. *J Hered*
86:248–249

735 Roeder AD, Archer FI, Poinar HN, Morin PA (2004) A novel method
 736 for collection and preservation of faeces for genetic studies. *Mol*
 737 *Ecol Notes* 4:761–764

738 Servheen C (1990) The status and conservation of the bears of the
 739 world. In: International conference of bear research and man-
 740 agement monography series, vol. 2, pp. 1–32

741 Sherwin WB, Moritz C (2000) Managing and monitoring genetic
 742 erosion. In: Young AJ, Clarke G (eds) *Demography genetics*
 743 *viability of fragmented populations*. Cambridge University
 744 Press, New York

745 Simberloff D (1988) The contribution of population and community
 746 biology to conservation science. *Annu Rev Ecol Syst* 19:473–
 747 511

748 Smith DA, Ralls K, Hurt A et al (2006) Assessing reliability of
 749 microsatellite genotypes from kit fox faecal samples using
 750 genetic and GIS analyses. *Mol Ecol* 15:387–406

751 Sokal RR, Rohlf FJ (1994) *Biometry: the principles and practice of*
 752 *statistics in biological research*. WH Freeman, New York

753 Soulé ME (ed) (1987) *Viable populations for conservation*. Cam-
 754 bridge University Press, Cambridge, UK

755 Taberlet P, Bouvet J (1994) Mitochondrial DNA polymorphism,
 756 phylogeography, and conservation genetics of the brown bear
 757 (*Ursus arctos*) in Europe. *Proc R Soc Lond B* 255:195–200

758 Taberlet P, Luikart G (1999) Non-invasive genetic sampling and
 759 individual identification. *Biol J Linn Soc* 68:41–55

760 Taberlet P, Griffin S, Goossens B et al (1996) Reliable genotyping of
 761 samples with very low DNA quantities using PCR. *Nucleic*
 762 *Acids Res* 26:3189–3194

763 Taberlet P, Camarra JJ, Griffin S et al (1997) Noninvasive genetic
 764 tracking of the endangered Pyrenean brown bear population. *Mol*
 765 *Ecol* 6:869–876

766 Taberlet P, Waits LP, Luikart G (1999) Noninvasive genetic
 767 sampling: look before you leap. *Trends Ecol Evol* 14:323–327

768 Taylor BL, Dizon AE (1999) First policy then science: why a
 769 management unit based solely on genetic criteria cannot work.
 770 *Mol Ecol* 8:S11–S16

771 Taylor AC, Sherwin WB, Wayne RK (1994) Genetic variation of
 772 microsatellite loci in a bottlenecked species: the northern hairy-
 773 nosed wombat *Lasiorhinus krefftii*. *Mol Ecol* 3:277–290

774 Valière N (2002) GIMLET, a computer program for analyzing genetic
 775 individual identification data. *Mol Ecol Notes* 2:377–379

776 Vigilant L (1999) An evaluation of techniques for the extraction and
 777 amplification of DNA from naturally shed hairs. *Biol Chem*
 778 380:1329–1331

779 Waits LP, Talbot S, Ward RH, Shields GF (1998) Mitochondrial
 780 DNA phylogeography of the North American brown bear and
 781 implications for conservation. *Conserv Biol* 12:408–417

782 Waits LP, Taberlet P, Swenson JE, Sandegren F (2000) Nuclear DNA
 783 microsatellite analysis of genetic diversity and gene flow in the
 784 Scandinavian brown bear (*Ursus arctos*). *Mol Ecol* 9:421–431

785 Waits LP, Luikart G, Taberlet P (2001) Estimating the probability of
 786 identity among genotypes in natural populations: cautions and
 787 guidelines. *Mol Ecol* 10:249–256

788 Weir BS (1996) *Genetic data analysis II*. Sinauer Associates Inc.,
 789 Massachusetts

790 Wiegand T, Naves J, Stephan T, Fernández A (1998) Assessing the
 791 risk of extinction for the brown bear (*Ursus arctos*) in the
 792 Cordillera Cantabrica, Spain. *Ecol Monogr* 68:539–571

793 Woods JG, Paetkau D, Lewis D et al (1999) Genetic tagging of free-
 794 ranging black and brown bears. *Wildl Soc B* 27:616–627

795 Zedrosser A, Dahle B, Swenson JE, Gerstl N (2001) Status and
 796 management of the brown bear in Europe. *Ursus* 12:9–20
 797

Author Proof

UNCORRECTED