

Search for the Decays $B_{(s)}^0 \rightarrow e^+ \mu^-$ and $B_{(s)}^0 \rightarrow e^+ e^-$ in CDF Run II

T. Aaltonen,²⁴ J. Adelman,¹⁴ T. Akimoto,⁵⁶ B. Álvarez González^{s, 12} S. Amerio^{y, 44} D. Amidei,³⁵ A. Anastassov,³⁹ A. Annovi,²⁰ J. Antos,¹⁵ G. Apollinari,¹⁸ A. Apresyan,⁴⁹ T. Arisawa,⁵⁸ A. Artikov,¹⁶ W. Ashmanskas,¹⁸ A. Attal,⁴ A. Aurisano,⁵⁴ F. Azfar,⁴³ P. Azzurri^{z, 47} W. Badgett,¹⁸ A. Barbaro-Galtieri,²⁹ V.E. Barnes,⁴⁹ B.A. Barnett,²⁶ V. Bartsch,³¹ G. Bauer,³³ P.-H. Beauchemin,³⁴ F. Bedeschi,⁴⁷ D. Beecher,³¹ S. Behari,²⁶ G. Bellettini^{z, 47} J. Bellinger,⁶⁰ D. Benjamin,¹⁷ A. Beretvas,¹⁸ J. Beringer,²⁹ A. Bhatti,⁵¹ M. Binkley,¹⁸ D. Bisello^{y, 44} I. Bizjak^{ee, 31} R.E. Blair,² C. Blocker,⁷ B. Blumenfeld,²⁶ A. Bocci,¹⁷ A. Bodek,⁵⁰ V. Boisvert,⁵⁰ G. Bolla,⁴⁹ D. Bortoletto,⁴⁹ J. Boudreau,⁴⁸ A. Boveia,¹¹ B. Brau^{a, 11} A. Bridgeman,²⁵ L. Brigliadori,⁴⁴ C. Bromberg,³⁶ E. Brubaker,¹⁴ J. Budagov,¹⁶ H.S. Budd,⁵⁰ S. Budd,²⁵ S. Burke,¹⁸ K. Burkett,¹⁸ G. Busetto^{y, 44} P. Bussey,²² A. Buzatu,³⁴ K. L. Byrum,² S. Cabrera^{u, 17} C. Calancha,³² M. Campanelli,³⁶ M. Campbell,³⁵ F. Canelli^{14, 18} A. Canepa,⁴⁶ B. Carls,²⁵ D. Carlsmith,⁶⁰ R. Carosi,⁴⁷ S. Carrillo^{n, 19} S. Carron,³⁴ B. Casal,¹² M. Casarsa,¹⁸ A. Castro^{x, 6} P. Catastini^{aa, 47} D. Cauz^{dd, 55} V. Cavaliere^{aa, 47} M. Cavalli-Sforza,⁴ A. Cerri,²⁹ L. Cerrito^{o, 31} S.H. Chang,²⁸ Y.C. Chen,¹ M. Chertok,⁸ G. Chiarelli,⁴⁷ G. Chlachidze,¹⁸ F. Chlebana,¹⁸ K. Cho,²⁸ D. Chokheli,¹⁶ J.P. Chou,²³ G. Choudalakis,³³ S.H. Chuang,⁵³ K. Chung,¹³ W.H. Chung,⁶⁰ Y.S. Chung,⁵⁰ T. Chwalek,²⁷ C.I. Ciobanu,⁴⁵ M.A. Ciocci^{aa, 47} A. Clark,²¹ D. Clark,⁷ G. Compostella,⁴⁴ M.E. Convery,¹⁸ J. Conway,⁸ M. Cordelli,²⁰ G. Cortiana^{y, 44} C.A. Cox,⁸ D.J. Cox,⁸ F. Crescioli^{z, 47} C. Cuenca Almenar^{u, 8} J. Cuevas^{s, 12} R. Culbertson,¹⁸ J.C. Cully,³⁵ D. Dagenhart,¹⁸ M. Datta,¹⁸ T. Davies,²² P. de Barbaro,⁵⁰ S. De Cecco,⁵² A. Deisher,²⁹ G. De Lorenzo,⁴ M. Dell'Orso^{z, 47} C. Deluca,⁴ L. Demortier,⁵¹ J. Deng,¹⁷ M. Deninno,⁶ P.F. Derwent,¹⁸ G.P. di Giovanni,⁴⁵ C. Dionisi^{cc, 52} B. Di Ruzza^{dd, 55} J.R. Dittmann,⁵ M. D'Onofrio,⁴ S. Donati^{z, 47} P. Dong,⁹ J. Donini,⁴⁴ T. Dorigo,⁴⁴ S. Dube,⁵³ J. Efron,⁴⁰ A. Elagin,⁵⁴ R. Erbacher,⁸ D. Errede,²⁵ S. Errede,²⁵ R. Eusebi,¹⁸ H.C. Fang,²⁹ S. Farrington,⁴³ W.T. Fedorko,¹⁴ R.G. Feild,⁶¹ M. Feindt,²⁷ J.P. Fernandez,³² C. Ferrazza^{bb, 47} R. Field,¹⁹ G. Flanagan,⁴⁹ R. Forrest,⁸ M.J. Frank,⁵ M. Franklin,²³ J.C. Freeman,¹⁸ I. Furic,¹⁹ M. Gallinaro,⁵² J. Galyardt,¹³ F. Garberson,¹¹ J.E. Garcia,²¹ A.F. Garfinkel,⁴⁹ K. Genser,¹⁸ H. Gerberich,²⁵ D. Gerdes,³⁵ A. Gessler,²⁷ S. Giagu^{cc, 52} V. Giakoumopoulou,³ P. Giannetti,⁴⁷ K. Gibson,⁴⁸ J.L. Gimmell,⁵⁰ C.M. Ginsburg,¹⁸ N. Giokaris,³ M. Giordani^{dd, 55} P. Giromini,²⁰ M. Giunta^{z, 47} G. Giurgiu,²⁶ V. Glagolev,¹⁶ D. Glenzinski,¹⁸ M. Gold,³⁸ N. Goldschmidt,¹⁹ A. Golossanov,¹⁸ G. Gomez,¹² G. Gomez-Ceballos,³³ M. Goncharov,³³ O. González,³² I. Gorelov,³⁸ A.T. Goshaw,¹⁷ K. Goulianos,⁵¹ A. Gresele^{y, 44} S. Grinstein,²³ C. Grosso-Pilcher,¹⁴ R.C. Group,¹⁸ U. Grundler,²⁵ J. Guimaraes da Costa,²³ Z. Gunay-Unalan,³⁶ C. Haber,²⁹ K. Hahn,³³ S.R. Hahn,¹⁸ E. Halkiadakis,⁵³ B.-Y. Han,⁵⁰ J.Y. Han,⁵⁰ F. Happacher,²⁰ K. Hara,⁵⁶ D. Hare,⁵³ M. Hare,⁵⁷ S. Harper,⁴³ R.F. Harr,⁵⁹ R.M. Harris,¹⁸ M. Hartz,⁴⁸ K. Hatakeyama,⁵¹ C. Hays,⁴³ M. Heck,²⁷ A. Heijboer,⁴⁶ J. Heinrich,⁴⁶ C. Henderson,³³ M. Herndon,⁶⁰ J. Heuser,²⁷ S. Hewamanage,⁵ D. Hidas,¹⁷ C.S. Hill^{c, 11} D. Hirschbuehl,²⁷ A. Hocker,¹⁸ S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,²⁹ B.T. Huffman,⁴³ R.E. Hughes,⁴⁰ U. Husemann,⁶¹ M. Hussein,³⁶ J. Huston,³⁶ J. Incandela,¹¹ G. Introzzi,⁴⁷ M. Iori^{cc, 52} A. Ivanov,⁸ E. James,¹⁸ D. Jang,¹³ B. Jayatilaka,¹⁷ E.-J. Jeon,²⁸ M.K. Jha,⁶ S. Jindariani,¹⁸ W. Johnson,⁸ M. Jones,⁴⁹ K.K. Joo,²⁸ S.Y. Jun,¹³ J.E. Jung,²⁸ T.R. Junk,¹⁸ T. Kamon,⁵⁴ D. Kar,¹⁹ P.E. Karchin,⁵⁹ Y. Kato^{l, 42} R. Kephart,¹⁸ J. Keung,⁴⁶ V. Khotilovich,⁵⁴ B. Kilminster,¹⁸ D.H. Kim,²⁸ H.S. Kim,²⁸ H.W. Kim,²⁸ J.E. Kim,²⁸ M.J. Kim,²⁰ S.B. Kim,²⁸ S.H. Kim,⁵⁶ Y.K. Kim,¹⁴ N. Kimura,⁵⁶ L. Kirsch,⁷ S. Klimenko,¹⁹ B. Knuteson,³³ B.R. Ko,¹⁷ K. Kondo,⁵⁸ D.J. Kong,²⁸ J. Konigsberg,¹⁹ A. Korytov,¹⁹ A.V. Kotwal,¹⁷ M. Kreps,²⁷ J. Kroll,⁴⁶ D. Krop,¹⁴ N. Krumnack,⁵ M. Kruse,¹⁷ V. Krutelyov,¹¹ T. Kubo,⁵⁶ T. Kuhr,²⁷ N.P. Kulkarni,⁵⁹ M. Kurata,⁵⁶ S. Kwang,¹⁴ A.T. Laasanen,⁴⁹ S. Lami,⁴⁷ S. Lammel,¹⁸ M. Lancaster,³¹ R.L. Lander,⁸ K. Lannon^{r, 40} A. Lath,⁵³ G. Latino^{aa, 47} I. Lazzizzera^{y, 44} T. LeCompte,² E. Lee,⁵⁴ H.S. Lee,¹⁴ S.W. Lee^{t, 54} S. Leone,⁴⁷ J.D. Lewis,¹⁸ C.-S. Lin,²⁹ J. Linacre,⁴³ M. Lindgren,¹⁸ E. Lipeles,⁴⁶ A. Lister,⁸ D.O. Litvintsev,¹⁸ C. Liu,⁴⁸ T. Liu,¹⁸ N.S. Lockyer,⁴⁶ A. Loginov,⁶¹ M. Loreti^{y, 44} L. Lovas,¹⁵ D. Lucchesi^{y, 44} C. Luci^{cc, 52} J. Lueck,²⁷ P. Lujan,²⁹ P. Lukens,¹⁸ G. Lungu,⁵¹ L. Lyons,⁴³ J. Lys,²⁹ R. Lysak,¹⁵ D. MacQueen,³⁴ R. Madrak,¹⁸ K. Maeshima,¹⁸ K. Makhoul,³³ T. Maki,²⁴ P. Maksimovic,²⁶ S. Malde,⁴³ S. Malik,³¹ G. Manca^{e, 30} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁹ C. Marino,²⁷ C.P. Marino,²⁵ A. Martin,⁶¹ V. Martin^{k, 22} M. Martínez,⁴ R. Martínez-Ballarín,³² T. Maruyama,⁵⁶ P. Mastrandrea,⁵² T. Masubuchi,⁵⁶ M. Mathis,²⁶ M.E. Mattson,⁵⁹ P. Mazzanti,⁶ K.S. McFarland,⁵⁰ P. McIntyre,⁵⁴ R. McNulty^{j, 30} A. Mehta,³⁰ P. Mehtala,²⁴ A. Menzione,⁴⁷ P. Merkel,⁴⁹ C. Mesropian,⁵¹ T. Miao,¹⁸ N. Miladinovic,⁷ R. Miller,³⁶ C. Mills,²³ M. Milnik,²⁷ A. Mitra,¹ G. Mitselmakher,¹⁹ H. Miyake,⁵⁶ N. Moggi,⁶ C.S. Moon,²⁸ R. Moore,¹⁸ M.J. Morello^{z, 47} J. Morlock,²⁷ P. Movilla Fernandez,¹⁸ J. Mülmenstädt,²⁹ A. Mukherjee,¹⁸ Th. Muller,²⁷ R. Mumford,²⁶ P. Murat,¹⁸ M. Mussini^{x, 6} J. Nachtman,¹⁸ Y. Nagai,⁵⁶ A. Nagano,⁵⁶ J. Naganoma,⁵⁶ K. Nakamura,⁵⁶ I. Nakano,⁴¹ A. Napier,⁵⁷ V. Necula,¹⁷

J. Nett,⁶⁰ C. Neu^{v,46} M.S. Neubauer,²⁵ S. Neubauer,²⁷ J. Nielsen^{g,29} L. Nodulman,² M. Norman,¹⁰ O. Norniella,²⁵ E. Nurse,³¹ L. Oakes,⁴³ S.H. Oh,¹⁷ Y.D. Oh,²⁸ I. Oksuzian,¹⁹ T. Okusawa,⁴² R. Orava,²⁴ K. Osterberg,²⁴ S. Pagan Griso^{y,44} E. Palencia,¹⁸ V. Papadimitriou,¹⁸ A. Papaikonomou,²⁷ A.A. Paramonov,¹⁴ B. Parks,⁴⁰ S. Pashapour,³⁴ J. Patrick,¹⁸ G. Pauletta^{dd,55} M. Paulini,¹³ C. Paus,³³ T. Peiffer,²⁷ D.E. Pellett,⁸ A. Penzo,⁵⁵ T.J. Phillips,¹⁷ G. Piacentino,⁴⁷ E. Pianori,⁴⁶ L. Pinera,¹⁹ K. Pitts,²⁵ C. Plager,⁹ L. Pondrom,⁶⁰ O. Poukhov^{*,16} N. Pounder,⁴³ F. Prakoshyn,¹⁶ A. Pronko,¹⁸ J. Proudfoot,² F. Ptohos^{i,18} E. Pueschel,¹³ G. Punzi^{z,47} J. Pursley,⁶⁰ J. Rademacker^{c,43} A. Rahaman,⁴⁸ V. Ramakrishnan,⁶⁰ N. Ranjan,⁴⁹ I. Redondo,³² P. Renton,⁴³ M. Renz,²⁷ M. Rescigno,⁵² S. Richter,²⁷ F. Rimondi^{x,6} L. Ristori,⁴⁷ A. Robson,²² T. Rodrigo,¹² T. Rodriguez,⁴⁶ E. Rogers,²⁵ S. Rolli,⁵⁷ R. Roser,¹⁸ M. Rossi,⁵⁵ R. Rossin,¹¹ P. Roy,³⁴ A. Ruiz,¹² J. Russ,¹³ V. Rusu,¹⁸ B. Rutherford,¹⁸ H. Saarikko,²⁴ A. Safonov,⁵⁴ W.K. Sakumoto,⁵⁰ O. Saltó,⁴ L. Santi^{dd,55} S. Sarkar^{cc,52} L. Sartori,⁴⁷ K. Sato,¹⁸ A. Savoy-Navarro,⁴⁵ P. Schlabach,¹⁸ A. Schmidt,²⁷ E.E. Schmidt,¹⁸ M.A. Schmidt,¹⁴ M.P. Schmidt^{*,61} M. Schmitt,³⁹ T. Schwarz,⁸ L. Scodellaro,¹² A. Scribano^{aa,47} F. Scuri,⁴⁷ A. Sedov,⁴⁹ S. Seidel,³⁸ Y. Seiya,⁴² A. Semenov,¹⁶ L. Sexton-Kennedy,¹⁸ F. Sforza,⁴⁷ A. Sfyrta,²⁵ S.Z. Shalhout,⁵⁹ T. Shears,³⁰ P.F. Shepard,⁴⁸ M. Shimojima^{q,56} S. Shiraishi,¹⁴ M. Shochet,¹⁴ Y. Shon,⁶⁰ I. Shreyber,³⁷ A. Sidoti,⁴⁷ P. Sinervo,³⁴ A. Sisakyan,¹⁶ A.J. Slaughter,¹⁸ J. Slaunwhite,⁴⁰ K. Sliwa,⁵⁷ J.R. Smith,⁸ F.D. Snider,¹⁸ R. Snihur,³⁴ A. Soha,⁸ S. Somalwar,⁵³ V. Sorin,³⁶ J. Spalding,¹⁸ T. Spreitzer,³⁴ P. Squillacioti^{aa,47} M. Stanitzki,⁶¹ R. St. Denis,²² B. Stelzer,³⁴ O. Stelzer-Chilton,³⁴ D. Stentz,³⁹ J. Strologas,³⁸ G.L. Strycker,³⁵ D. Stuart,¹¹ J.S. Suh,²⁸ A. Sukhanov,¹⁹ I. Suslov,¹⁶ T. Suzuki,⁵⁶ A. Taffard^{f,25} R. Takashima,⁴¹ Y. Takeuchi,⁵⁶ R. Tanaka,⁴¹ M. Tecchio,³⁵ P.K. Teng,¹ K. Terashi,⁵¹ J. Thom^{h,18} A.S. Thompson,²² G.A. Thompson,²⁵ E. Thomson,⁴⁶ P. Tipton,⁶¹ P. Ttito-Guzmán,³² S. Tkaczyk,¹⁸ D. Toback,⁵⁴ S. Tokar,¹⁵ K. Tollefson,³⁶ T. Tomura,⁵⁶ D. Tonelli,¹⁸ S. Torre,²⁰ D. Torretta,¹⁸ P. Totaro^{dd,55} S. Tourneur,⁴⁵ M. Trovato,⁴⁷ S.-Y. Tsai,¹ Y. Tu,⁴⁶ N. Turini^{aa,47} F. Ukegawa,⁵⁶ S. Vallecorsa,²¹ N. van Remortel^{b,24} A. Varganov,³⁵ E. Vataga^{bb,47} F. Vázquez^{n,19} G. Velez,¹⁸ C. Vellidis,³ M. Vidal,³² R. Vidal,¹⁸ I. Vila,¹² R. Vilar,¹² T. Vine,³¹ M. Vogel,³⁸ I. Volobouev^{t,29} G. Volpi^{z,47} P. Wagner,⁴⁶ R.G. Wagner,² R.L. Wagner,¹⁸ W. Wagner^{w,27} J. Wagner-Kuhr,²⁷ T. Wakisaka,⁴² R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³⁴ D. Waters,³¹ M. Weinberger,⁵⁴ J. Weinelt,²⁷ H. Wenzel,¹⁸ W.C. Wester III,¹⁸ B. Whitehouse,⁵⁷ D. Whiteson^{f,46} A.B. Wicklund,² E. Wicklund,¹⁸ S. Wilbur,¹⁴ G. Williams,³⁴ H.H. Williams,⁴⁶ P. Wilson,¹⁸ B.L. Winer,⁴⁰ P. Wittich^{h,18} S. Wolbers,¹⁸ C. Wolfe,¹⁴ T. Wright,³⁵ X. Wu,²¹ F. Würthwein,¹⁰ S. Xie,³³ A. Yagil,¹⁰ K. Yamamoto,⁴² J. Yamaoka,¹⁷ U.K. Yang^{p,14} Y.C. Yang,²⁸ W.M. Yao,²⁹ G.P. Yeh,¹⁸ J. Yoh,¹⁸ K. Yorita,⁵⁸ T. Yoshida^{m,42} G.B. Yu,⁵⁰ I. Yu,²⁸ S.S. Yu,¹⁸ J.C. Yun,¹⁸ L. Zanello^{cc,52} A. Zanetti,⁵⁵ X. Zhang,²⁵ Y. Zheng^{d,9} and S. Zucchelli^{x,6}

(CDF Collaboration[†])

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^xUniversity of Bologna, I-40127 Bologna, Italy*

⁷*Brandeis University, Waltham, Massachusetts 02254*

⁸*University of California, Davis, Davis, California 95616*

⁹*University of California, Los Angeles, Los Angeles, California 90024*

¹⁰*University of California, San Diego, La Jolla, California 92093*

¹¹*University of California, Santa Barbara, Santa Barbara, California 93106*

¹²*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹³*Carnegie Mellon University, Pittsburgh, PA 15213*

¹⁴*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

¹⁵*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹⁶*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁷*Duke University, Durham, North Carolina 27708*

¹⁸*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

¹⁹*University of Florida, Gainesville, Florida 32611*

²⁰*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

²¹*University of Geneva, CH-1211 Geneva 4, Switzerland*

²²*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²³*Harvard University, Cambridge, Massachusetts 02138*

²⁴*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

²⁵*University of Illinois, Urbana, Illinois 61801*

²⁶*The Johns Hopkins University, Baltimore, Maryland 21218*

²⁷*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*

- ²⁸Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon, 305-806, Korea; Chonnam National University, Gwangju, 500-757, Korea
- ²⁹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720
- ³⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³¹University College London, London WC1E 6BT, United Kingdom
- ³²Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
- ³³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
- ³⁴Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³⁵University of Michigan, Ann Arbor, Michigan 48109
- ³⁶Michigan State University, East Lansing, Michigan 48824
- ³⁷Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ³⁸University of New Mexico, Albuquerque, New Mexico 87131
- ³⁹Northwestern University, Evanston, Illinois 60208
- ⁴⁰The Ohio State University, Columbus, Ohio 43210
- ⁴¹Okayama University, Okayama 700-8530, Japan
- ⁴²Osaka City University, Osaka 588, Japan
- ⁴³University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴⁴Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^uUniversity of Padova, I-35131 Padova, Italy
- ⁴⁵LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104
- ⁴⁷Istituto Nazionale di Fisica Nucleare Pisa, ^zUniversity of Pisa, ^{aa}University of Siena and ^{bb}Scuola Normale Superiore, I-56127 Pisa, Italy
- ⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260
- ⁴⁹Purdue University, West Lafayette, Indiana 47907
- ⁵⁰University of Rochester, Rochester, New York 14627
- ⁵¹The Rockefeller University, New York, New York 10021
- ⁵²Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, ^{cc}Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵³Rutgers University, Piscataway, New Jersey 08855
- ⁵⁴Texas A&M University, College Station, Texas 77843
- ⁵⁵Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, ^{dd}University of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁶University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁵⁷Tufts University, Medford, Massachusetts 02155
- ⁵⁸Waseda University, Tokyo 169, Japan
- ⁵⁹Wayne State University, Detroit, Michigan 48201
- ⁶⁰University of Wisconsin, Madison, Wisconsin 53706
- ⁶¹Yale University, New Haven, Connecticut 06520

We report results from a search for the lepton flavor violating decays $B_{(s)}^0 \rightarrow e^+\mu^-$, and the flavor-changing neutral-current decays $B_{(s)}^0 \rightarrow e^+e^-$. The analysis uses data corresponding to 2 fb^{-1} of integrated luminosity of $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$ collected with the upgraded Collider Detector (CDF II) at the Fermilab Tevatron. The observed number of $B_{(s)}^0$ candidates is consistent with background expectations. The resulting Bayesian upper limits on the branching ratios at 90% credibility level are $\mathcal{B}(B_s^0 \rightarrow e^+\mu^-) < 2.0 \times 10^{-7}$, $\mathcal{B}(B^0 \rightarrow e^+\mu^-) < 6.4 \times 10^{-8}$, $\mathcal{B}(B_s^0 \rightarrow e^+e^-) < 2.8 \times 10^{-7}$ and $\mathcal{B}(B^0 \rightarrow e^+e^-) < 8.3 \times 10^{-8}$. From the limits on $\mathcal{B}(B_{(s)}^0 \rightarrow e^+\mu^-)$, the following lower bounds on the Pati-Salam leptoquark masses are also derived: $M_{LQ}(B_s^0 \rightarrow e^+\mu^-) > 47.8\text{ TeV}/c^2$, and $M_{LQ}(B^0 \rightarrow e^+\mu^-) > 59.3\text{ TeV}/c^2$, at 90% credibility level.

PACS numbers: 13.20.He 13.30.Ce 12.15.Mm 12.60.-i

*Deceased

†With visitors from ^aUniversity of Massachusetts Amherst, Amherst, Massachusetts 01003, ^bUniversiteit Antwerpen, B-2610 Antwerp, Belgium, ^cUniversity of Bristol, Bristol BS8 1TL,

United Kingdom, ^dChinese Academy of Sciences, Beijing 100864, China, ^eIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^fUniversity of California Irvine, Irvine, CA 92697, ^gUniversity of California Santa Cruz,

Rare particle decays that are either forbidden within the standard model of particle physics (SM), or are expected to have very small branching ratios provide excellent signatures with which to look for new physics and allow to probe subatomic processes that are beyond the reach of direct searches. The decays $B_{(s)}^0 \rightarrow e^+\mu^-$ [1] are forbidden within the SM, in which lepton number and lepton flavor are conserved. However the observation of neutrino oscillations indicates that lepton flavor is not conserved. To date, no lepton flavor violating (LFV) decays in the charged sector such as $B_{(s)}^0 \rightarrow e^+\mu^-$ have been observed. These decays are allowed in models where the SM has been extended by heavy singlet Dirac neutrinos [2]. The LFV decays are also allowed in some physics scenarios beyond the SM, such as the Pati-Salam model [3] and supersymmetry (SUSY) models [4]. The grand-unification theory by J. Pati and A. Salam predicts a new interaction to mediate transitions between leptons and quarks via exchange of spin-1 gauge bosons, which are called Pati-Salam leptoquarks (LQ), that carry both color and lepton quantum numbers [3]. The lepton and quark components of the leptoquarks are not necessarily from the same generation [5, 6], and the decays $B_s^0 \rightarrow e^+\mu^-$ and $B^0 \rightarrow e^+\mu^-$ can be mediated by different types of leptoquarks. Processes involving flavor-changing neutral currents (FCNCs) can occur in the SM only through higher-order Feynman diagrams where new physics contributions can provide a significant enhancement. Compared to $B_{(s)}^0 \rightarrow \mu^+\mu^-$ [7], the FCNC decays of $B_{(s)}^0 \rightarrow e^+e^-$ are further suppressed by the square of the ratio of the electron and muon masses $(m_e/m_\mu)^2$. The SM expectations for branching ratios of $B_{(s)}^0 \rightarrow e^+e^-$ are of the order of 10^{-15} [8].

In this Letter we report on a search for the LFV decays $B_{(s)}^0 \rightarrow e^+\mu^-$ and the FCNC decays $B_{(s)}^0 \rightarrow e^+e^-$, using a data sample corresponding to 2 fb^{-1} of integrated luminosity collected in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$. With no evidence for either the LFV or FCNC decays, we set upper limits on their branching ratios using the common reference decay $B^0 \rightarrow K^+\pi^-$, which has a precisely-known branching ratio. This is the first time a search for

$B_s^0 \rightarrow e^+e^-$ has been performed.

A detailed description of the CDF II detector can be found in Ref. [9]. Here we give a brief description of the detector elements most relevant to this analysis. Charged particle tracking is provided by a silicon microstrip detector together with the surrounding open-cell wire drift chamber (COT), both immersed in a 1.4 T axial magnetic field. The tracking system provides precise vertex and momentum measurement for charged particles in the pseudorapidity range $|\eta| < 1.0$ [10]. Surrounding the tracking system are electromagnetic (CEM) and hadronic sampling calorimeters, arranged in a projective geometry. Drift chambers and scintillation counters are located behind the calorimeters to detect muons within $|\eta| < 0.6$ (CMU) and $0.6 < |\eta| < 1.0$ (CMX).

We use a data sample enriched in two-body B -decays selected by a three-level trigger system using the extremely fast tracker [11] at level-1, and the silicon vertex trigger [12] at level-2. The trigger requires two oppositely-charged tracks, each with a transverse momentum $p_T > 2\text{ GeV}/c$, and an impact parameter [13] $0.1 < d_0 < 1\text{ mm}$. It also requires the scalar sum of the transverse momenta of the two tracks to be greater than $5.5\text{ GeV}/c$, the difference in the azimuthal angles of the tracks $20^\circ < \Delta\varphi < 135^\circ$, and a transverse decay length [14] $L_{xy} > 200\text{ }\mu\text{m}$. At the level-3 trigger stage, and in the offline analysis, the trigger selections are enforced with a more accurate determination of the same quantities. In the off-line analysis, additionally we require: the B -meson isolation $I > 0.675$ [15], the pointing angle $\Delta\phi < 6.3^\circ$ [16], and a tighter selection of $L_{xy} > 375\text{ }\mu\text{m}$. These three thresholds were optimized in an unbiased way to obtain the best sensitivity for the searches using the procedure described in Ref. [17].

Electron and muon identification is applied in the selection of $B_{(s)}^0 \rightarrow e^+\mu^-$ and $B_{(s)}^0 \rightarrow e^+e^-$ decay modes. The electron identification [18] requires that both the specific ionization (dE/dx) measured in the COT, and the transverse and longitudinal shower shape as measured in the CEM, be consistent with the hypothesis that the particle is an electron. The performance of electron identification is optimized using pure electron samples reconstructed from $\gamma \rightarrow e^+e^-$ conversions and hadron and muon samples from $D^0 \rightarrow K^-\pi^+$, $\Lambda \rightarrow p\pi^-$, and $J/\psi \rightarrow \mu^+\mu^-$ decays. We find the identification efficiency to be around 70% for electrons. The muon identification starts from tracks in the COT that are extrapolated into the muon detectors and are required to match hits in the muon systems. The muon selection is fully efficient for muons with $p_T > 2\text{ GeV}/c$ in CMU or CMX.

The mass resolution σ_m of fully-reconstructed B -meson decays to two charged particles is about $28\text{ MeV}/c^2$. Energy loss due to bremsstrahlung by electrons generates a tail on the low side of the mass distribution. This tail is more prominent for the $B_{(s)}^0 \rightarrow e^+e^-$ channels, where two electrons are involved. We define search windows of $(5.262\text{--}5.477)\text{ GeV}/c^2$ for $B_s^0 \rightarrow e^+\mu^-$

Santa Cruz, CA 95064, ^hCornell University, Ithaca, NY 14853, ⁱUniversity of Cyprus, Nicosia CY-1678, Cyprus, ^jUniversity College Dublin, Dublin 4, Ireland, ^kUniversity of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, ^lUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017 ^mKinki University, Higashi-Osaka City, Japan 577-8502 ⁿUniversidad Iberoamericana, Mexico D.F., Mexico, ^oQueen Mary, University of London, London, E1 4NS, England, ^pUniversity of Manchester, Manchester M13 9PL, England, ^qNagasaki Institute of Applied Science, Nagasaki, Japan, ^rUniversity of Notre Dame, Notre Dame, IN 46556, ^sUniversity de Oviedo, E-33007 Oviedo, Spain, ^tTexas Tech University, Lubbock, TX 79609, ^uIFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain, ^vUniversity of Virginia, Charlottesville, VA 22904, ^wBergische Universität Wuppertal, 42097 Wuppertal, Germany, ^{ee}On leave from J. Stefan Institute, Ljubljana, Slovenia,

and (5.171–5.387) GeV/c² for $B^0 \rightarrow e^+\mu^-$. These correspond to a window around the nominal values of the B_s^0 and B^0 masses [19] of approximately $\pm 3\sigma_m$. To recover some of the acceptance loss due to electron bremsstrahlung for the $B_{(s)}^0 \rightarrow e^+e^-$ channels, we choose wider and asymmetric search windows ranging from 6 σ_m below to 3 σ_m above the nominal values of the B_s^0 and B^0 masses. The search windows are (5.154–5.477) GeV/c² for the B_s^0 and (5.064–5.387) GeV/c² for the B^0 . The sideband regions (4.800–5.028) GeV/c² and (5.549–5.800) GeV/c² are used to estimate the combinatorial backgrounds.

The background contributions considered include combinations of random track pairs and partial B decays that accidentally meet the selection requirement (combinatorial), and hadronic two-body B decays in which both final particles are misidentified as leptons. The combinatorial background is evaluated by extrapolating the normalized number of events found in the sidebands to the signal region. The double-lepton misidentification rate is determined by applying electron and muon misidentification probabilities to the number of two-body decays found in the search window.

Figure 1 shows the invariant mass distribution for $e^+\mu^-$ candidates. We observe one event in the B_s^0 mass window, and two events in the B^0 mass window, consistent with the estimated total background of 0.8 ± 0.6 events in the B_s^0 search window, and 0.9 ± 0.6 in the B^0 window. The combinatorial background in both channels is estimated to be 0.7 ± 0.6 events. The number of events where two tracks are misidentified as electron and muon is estimated to be 0.09 ± 0.02 for the B_s^0 case and 0.22 ± 0.04 for the B^0 case.

Figure 2 shows the invariant mass distributions for e^+e^- candidate pairs where both tracks were identified as electrons. We observe one event in the B_s^0 mass window, and two events in the B^0 mass window. We estimate the total background contributions to be 2.7 ± 1.8 events in both the B_s^0 and B^0 mass windows. The dominant contribution comes from combinatorial background: 2.7 ± 1.8 compared to the contribution where both tracks are misidentified as electrons: 0.038 ± 0.008 for both B_s^0 or B^0 .

We use the reference decay $B^0 \rightarrow K^+\pi^-$ to set a limit on $\mathcal{B}(B_s^0 \rightarrow e^+\ell^-)$ (where ℓ is either e or μ), using the following expression:

$$\mathcal{B}(B_s^0 \rightarrow e^+\ell^-) = \frac{N(B_s^0 \rightarrow e^+\ell^-) \cdot \mathcal{B}(B^0 \rightarrow K^+\pi^-) \cdot f_d/f_s}{\epsilon_{B_s^0 \rightarrow e^+\ell^-}^{rel} \cdot N(B^0 \rightarrow K^+\pi^-)}.$$

The expression for the B^0 channels is identical, except that the ratio of b -quark fragmentation probabilities: f_d/f_s is not present. In the expression, $N(B_s^0 \rightarrow e^+\ell^-)$ is the calculated upper limit on the number of $B_s^0 \rightarrow e^+\ell^-$ events, $N(B^0 \rightarrow K^+\pi^-)$ is the observed number of events from the reference channel $B^0 \rightarrow K^+\pi^-$, $\mathcal{B}(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$ [19] is the branch-

ing ratio for the $B^0 \rightarrow K^+\pi^-$ decay, and $\epsilon_{B_s^0 \rightarrow e^+\ell^-}^{rel}$ is the detector acceptance and event selection efficiency for reconstructing $B_s^0 \rightarrow e^+\ell^-$ decays relative to that for $B^0 \rightarrow K^+\pi^-$. The value of f_d/f_s is 3.86 ± 0.59 , where the (anti-)correlation between the uncertainties has been accounted for [20]. To calculate the detector acceptance, we use simulated events with a detailed simulation of the CDF II detector and event selection. We obtain $\epsilon_{B_s^0 \rightarrow e^+\mu^-}^{rel} = 0.207 \pm 0.016$, $\epsilon_{B^0 \rightarrow e^+\mu^-}^{rel} = 0.210 \pm 0.012$, $\epsilon_{B_s^0 \rightarrow e^+e^-}^{rel} = 0.129 \pm 0.011$, and $\epsilon_{B^0 \rightarrow e^+e^-}^{rel} = 0.128 \pm 0.011$. The uncertainties listed above are the combined statistical and systematic uncertainties. The later include uncertainties from detector fiducial coverage, electron and muon identification efficiencies, detector material determination, $B_{(s)}^0$ p_T spectrum, and $B_{(s)}^0$ lifetimes. The reference channel $B^0 \rightarrow K^+\pi^-$ has been reconstructed using the same selection criteria except lepton identification. We find 6387 ± 214 $B^0 \rightarrow K^+\pi^-$ events, using a fitting procedure similar to that described in Ref. [21].

The upper limit on the branching ratio in each search window is obtained using the Bayesian approach [19], assuming a flat prior, and incorporating Gaussian uncertainties into the limit. The total systematic uncertainties, listed in Table I, are used as input for the limit calculation. Table II lists the upper limits we obtain on the branching ratios at 90% (95%) credibility level (C.L.).

Within the Pati-Salam leptoquark model, the following relationship between the $\mathcal{B}(B_{(s)}^0 \rightarrow e^+\mu^-)$ and the leptoquark mass (M_{LQ}) can be derived [5]:

$$\mathcal{B}(B_{(s)}^0 \rightarrow e^+\mu^-) = \pi \alpha_s^2(M_{LQ}) \frac{1}{M_{LQ}^4} F_{B_{(s)}^0}^2 m_{B_{(s)}^0}^3 R^2 \cdot \frac{\tau_{B_{(s)}^0}}{\hbar},$$

where $R = \frac{m_{B_{(s)}^0}}{m_b} \left(\frac{\alpha_s(M_{LQ})}{\alpha_s(m_t)} \right)^{-\frac{4}{3}} \left(\frac{\alpha_s(m_t)}{\alpha_s(m_b)} \right)^{-\frac{12}{23}}$. The values and uncertainties of the quantities used in the calculation of M_{LQ} are the following [19]: the top-quark mass m_t (171.2 ± 2.1 GeV/c²), the bottom quark mass m_b (4.20 ± 0.17 GeV/c²), the charm quark mass m_c (1.27 ± 0.11 GeV/c²), the B^0 -meson mass m_{B^0} (5.27953 ± 0.00033 GeV/c²), the B_s^0 -meson mass $m_{B_s^0}$ (5.3663 ± 0.0006 GeV/c²), the B^0 -meson lifetime τ_{B^0} (1.530 ± 0.009 ps), the B_s^0 -meson lifetime $\tau_{B_s^0}$ (1.470 ± 0.027 ps), the coupling strength F_{B^0} (0.178 ± 0.014 GeV), and $F_{B_s^0}$ (0.200 ± 0.014 GeV) [22]. For the strong coupling constant we use $\alpha_s(M_{Z^0}) = 0.115$, which is evolved to M_{LQ} using the Marciano approximation [23] assuming no colored particles exist with masses between m_t and M_{LQ} . Using the limits on the branching ratios listed in Table II, we calculate limits on the masses of the corresponding Pati-Salam leptoquarks of $M_{LQ}(B_s^0 \rightarrow e^+\mu^-) > 47.8$ (44.9) TeV/c² and $M_{LQ}(B^0 \rightarrow e^+\mu^-) > 59.3$ (56.3) TeV/c² at 90 (95)% C.L. Figure 3 shows the limit and the relation between the leptoquark mass and the branching ratio for the B_s^0 meson.

TABLE I: Values used to calculate the limits on $\mathcal{B}(B_{(s)}^0 \rightarrow e^+\mu^-)$ and $\mathcal{B}(B_{(s)}^0 \rightarrow e^+e^-)$ and their uncertainties.

Source	Values	$\mathcal{B}(B_s^0 \rightarrow e^+\mu^-)$	$\mathcal{B}(B^0 \rightarrow e^+\mu^-)$	$\mathcal{B}(B_s^0 \rightarrow e^+e^-)$	$\mathcal{B}(B^0 \rightarrow e^+e^-)$
$N(B^0 \rightarrow K^+\pi^-)$	6387 ± 214	3.4%	3.4%	3.4%	3.4%
$\mathcal{B}(B^0 \rightarrow K^+\pi^-)$	$(19.4 \pm 0.6) \times 10^{-6}$	3.1%	3.1%	3.1%	3.1%
$f_{B^0}/f_{B_s^0}$	3.86 ± 0.59	15.3%	-	15.3%	-
$\epsilon_{B^0 \rightarrow e^+\mu^-}^{rel}$	0.207 ± 0.016	7.6%	-	-	-
$\epsilon_{B^0 \rightarrow e^+\mu^-}^{rel}$	0.210 ± 0.012	-	5.9%	-	-
$\epsilon_{B^0 \rightarrow e^+e^-}^{rel}$	0.129 ± 0.011	-	-	8.9%	-
$\epsilon_{B^0 \rightarrow e^+e^-}^{rel}$	0.128 ± 0.011	-	-	-	8.9%
Total		17.7%	7.5%	18.3%	10.0%





 FIG. 1: Invariant mass distribution of $e^+\mu^-$ pairs for events where one track passed the electron identification and the other track the muon identification. The B_s^0 (B^0) search window is indicated by the solid (dotted) line. The sideband regions are indicated by the dashed lines.

 FIG. 2: Invariant mass distributions of e^+e^- pairs for events where both tracks passed the electron identification. The B_s^0 (B^0) search window is indicated by the solid (dotted) line. The sideband regions are indicated by dashed lines.

TABLE II: Branching ratio limits at 90(95) % C.L.

$\mathcal{B}(B_s^0 \rightarrow e^+\mu^-) < 2.0 \text{ (2.6)} \times 10^{-7}$
$\mathcal{B}(B^0 \rightarrow e^+\mu^-) < 6.4 \text{ (7.9)} \times 10^{-8}$
$\mathcal{B}(B_s^0 \rightarrow e^+e^-) < 2.8 \times 10^{-7}$
$\mathcal{B}(B^0 \rightarrow e^+e^-) < 8.3 \times 10^{-8}$


 FIG. 3: Leptoquark mass limit corresponding to the 90 (95) % C.L. on $\mathcal{B}(B_s^0 \rightarrow e^+\mu^-)$. The error band is obtained by varying the values entering the theoretical calculation within their uncertainties. The uncertainties stemming from approximating α_s are not included.

In summary, we report on a search for the lepton flavor violating decays $B_{(s)}^0 \rightarrow e^+\mu^-$ and the flavor changing neutral current decays $B_{(s)}^0 \rightarrow e^+e^-$ using data corresponding to 2 fb^{-1} of integrated luminosity collected in $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$. This is the first search for $B_s^0 \rightarrow e^+e^-$ decays. We observe no evidence for these decays and set limits that are the most stringent to date. These results represent a significant improvement compared to the previous measurement [24] by CDF and the best results from B -Factories [25, 26, 27].

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of

Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

-
- [1] Throughout this Letter inclusion of charge conjugate reactions is implied.
- [2] A. Ilakovac, Phys. Rev. D **62**, 036010 (2000).
- [3] J.C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974).
- [4] R. A. Diaz, R. Martinez, C. E. Sandoval, Eur. Phys. J. C **41**, 305 (2005).
- [5] G. Valencia and S. Willenbrock, Phys. Rev. D **50**, 6843 (1994).
- [6] M. Blanke *et al.* J. High Energy Phys. 05 (2007) 103.
- [7] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **100**, 101802 (2008).
- [8] M. Misiak and J. Urban, Phys. Lett. B **451**, 161 (1999); G. Buchalla and A. J. Burn, Nucl. Phys. **B548**, 309 (1999).
- [9] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005); and references therein.
- [10] The polar angle (θ) in cylindrical coordinates is measured with respect to the proton beam direction, which defines the z -axis. Pseudorapidity (η) is defined as $\eta = -\ln(\tan \frac{\theta}{2})$.
- [11] E.J. Thomson *et al.*, IEEE Trans. Nucl. Sci. **49**, 1063 (2002).
- [12] W. Ashmanskas *et al.*, Nucl. Instrum. Methods A **518**, 532 (2004).
- [13] The impact parameter d_0 is the distance of closest approach of the track to the beam line.
- [14] F. Abe *et al.* (CDF Collaboration), Phys. Rev. D **57**, 5382 (1998).
- [15] Due to the hard b -quark fragmentation, B -mesons carry most of the momentum of the b -quark. The isolation is defined as $I = p_T(B)/(\sum_i p_T^i + p_T(B))$, where $p_T(B)$ is the transverse momentum of the B candidate, and the sum runs over all other tracks within a cone of radius 1 in $\eta - \phi$ space around the B flight direction.
- [16] For track pairs coming from the two-body decay of a B , the vector pointing from the primary vertex to the B decay vertex in the transverse plane $\vec{l}_{xy}$ should point in the same direction as the transverse momentum vector $\vec{p}_T(B)$ of the B candidate. $\Delta\phi$ is defined as the angle between $\vec{l}_{xy}$ and $\vec{p}_T(B)$.
- [17] G. Punzi, arXiv:physics/0308063v2.
- [18] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **97**, 012002 (2006).
- [19] C. Amsler *et al.*, Physics Letters B **667**, 1 (2008)
- [20] Heavy Flavor Averaging Group, arXiv:hep-ex/0704.3575v1.
- [21] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **97**, 211802 (2006).
- [22] J. Bordes *et al.* J. High Energy Phys. 12 (2004) 064.
- [23] W. J. Marciano, Phys. Rev. D **29**, 580 (1984).
- [24] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **81**, 5742 (1998).
- [25] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **99**, 251803 (2007).
- [26] M. C. Chang *et al.* (Belle Collaboration), Phys. Rev. D **68**, 111101 (2003).
- [27] T. Bergfeld *et al.* (CLEO Collaboration), Phys. Rev. D **62**, 091102 (2000).