

Myxococcus xanthus induces actinorhodin overproduction and aerial mycelium formation by *Streptomyces coelicolor*

Juana Pérez,¹ José Muñoz-Dorado,¹
Alfredo F. Braña,² Lawrence J. Shimkets,³
Laura Sevillano⁴ and Ramón I. Santamaría^{4*}

¹Departamento de Microbiología, Facultad de Ciencias, Universidad de Granada, E-18071 Granada, Spain.

²Area de Microbiología, Departamento de Biología Funcional, Facultad de Medicina, IUBA, Universidad de Oviedo, 33006 Oviedo, Asturias, Spain.

³Department of Microbiology, University of Georgia, Athens, GA, USA.

⁴Instituto de Microbiología Bioquímica/Departamento de Microbiología y Genética. Consejo Superior de Investigaciones Científicas (CSIC)/Universidad de Salamanca, Edificio Departamental. Campus Miguel de Unamuno. 37007-Salamanca, Spain.

Summary

Interaction of the predatory myxobacterium *Myxococcus xanthus* with the non-motile, antibiotic producer *Streptomyces coelicolor* was examined using a variety of experimental approaches. *Myxococcus xanthus* cells prey on *S. coelicolor*, forming streams of ordered cells that lyse the *S. coelicolor* hyphae in the contact area between the two colonies. The interaction increases actinorhodin production by *S. coelicolor* up to 20-fold and triggers aerial mycelium production. Other bacteria are also able to induce these processes in *S. coelicolor* though to a lesser extent. These studies offer new clues about the expression of genes that remain silent or are expressed at low level in axenic cultures and open the possibility of overproducing compounds of biotechnological interest by using potent inducers synthesized by other bacteria.

Introduction

Bacterial populations in natural habitats are complex communities containing many species that exhibit competition and/or collaboration in order to survive with limiting nutri-

tional resources. The study of these interactions has attracted much interest (Shank and Kolter, 2009; Straight and Kolter, 2009; Vos and Velicer, 2009). Laboratory co-cultures sometimes trigger the expression of genes that remain silent in pure cultures (Yamanaka *et al.*, 2005). The number of silent or poorly expressed genes under laboratory conditions may be much higher than originally thought (Schneiker *et al.*, 2007). Perhaps in natural communities unknown signals trigger gene expression, sometimes in other organisms (Bassler and Losick, 2006).

In this work we have examined the confrontation between two typical soil inhabitants, immobile *Streptomyces coelicolor* and the mobile predator *Myxococcus xanthus*. Both bacteria have genomes in excess of 8 Mb and are endowed with the capacity to produce many secondary metabolites. *Streptomyces coelicolor* contains 23 gene clusters related to secondary metabolite production (Bentley *et al.*, 2002). However, only four antibiotics have been detected under laboratory conditions, actinorhodin (ACT), undecylprodigiosin (RED) and calcium dependent antibiotic (CDA), synthesized by proteins encoded by the chromosome, and methylenomicin, whose biosynthetic enzymes are encoded by plasmid SCP1. ACT and RED are pigmented and their production is easily visualized. The genome sequence of *M. xanthus* DK1622 has revealed the presence of at least 18 clusters of polyketide/non-ribosomal peptide genes, most of which are not expressed under laboratory conditions (Wenzel and Muller, 2009). In fact, no antibiotics were identified in *M. xanthus* cultures until 2005 when the use of high-performance liquid chromatograph mass spectrometry (HPLC-MS) technology provided a sensitive method to identify five antibiotic families (Wenzel and Muller, 2009).

There are several groups of bacterial predators. *Myxococcus xanthus* has been extensively studied (Berleman and Kirby, 2009; Velicer and Mendes-Soares, 2009a) and moves on solid surfaces by two surface translocation mechanisms, the adventurous (A motility) and social (S motility) motility systems (Mauriello and Zusman, 2007). This myxobacterium preys on a wide variety of microorganisms by secreting lytic enzymes and toxic molecules (Velicer and Mendes-Soares, 2009a). It attacks in groups like a wolf pack by surrounding the prey (Velicer and

Received 8 July, 2010; accepted 31 July, 2010. *For correspondence. E-mail santa@usal.es; Tel. (+34) 923 294732; Fax (+34) 923 224876.

© 2010 The Authors

Journal compilation © 2010 Society for Applied Microbiology and Blackwell Publishing Ltd

Mendes-Soares, 2009a). *Myxococcus* is not a specialized predator, and can feed on a single species including *Escherichia coli*, *Corynebacterium glutamicum*, *Micrococcus luteus* and *Saccharomyces cerevisiae* (Hillesland *et al.*, 2007; Berleman and Kirby, 2009). Other predators are specialists and only feed a single species (Velicer and Mendes-Soares, 2009a).

During *M. xanthus* S motility, cells glide in groups. This social behaviour has been more extensively studied during fruiting body development following nutrition depletion. Development culminates when the rod-shaped vegetative cells differentiate into metabolically quiescent, spherical spores that germinate when nutrients are supplied (Dworkin, 1996). Fruiting body development and predation of other organisms induce rippling behaviour during which the cells organize themselves in parallel ridges that move coordinately (Berleman *et al.*, 2006). In addition, the social and multicellular behaviour of this bacterium requires intercellular communication (Kroos, 2007; Velicer and Vos, 2009b).

Streptomyces also has a complex developmental cycle that begins with the germination of a spore to form multigenomic substrate mycelia. Some mycelia erect aerial mycelia that generate unigenomic spores by transverse division of the tips (Chater, 1993; Flardh and Buttner, 2009). All this summarized process needs the action of a wide number of genes and signals among which the surfactant SapB and eight chaplins play an important role in the development of aerial mycelium (Capstick *et al.*, 2007).

In nature, actinomycetes are very abundant and they contribute to the fertility of soil degrading organic material and interacting with other organisms that live free or form part of the rhizosphere (Mazzola, 2007; Tamilarasi *et al.*, 2008; Chater *et al.*, 2010). These interactions may trigger the induction of otherwise silent secondary metabolite pathways and they are starting to be described (Straight *et al.*, 2007; Kurosawa *et al.*, 2008; Schroeckh *et al.*, 2009).

In this article we show that *M. xanthus* induces antibiotic production and differentiation by *S. coelicolor*. These results reinforce the idea that examination of interactions between microorganisms can increase the production of secondary metabolites and/or lead to the discovery of new metabolic compounds.

Results

Predation and competition between *Myxococcus* and *Streptomyces*

The interaction between *Myxococcus* and *Streptomyces* was examined in co-culture on an agar surface. A lawn of *M. xanthus* DK1622 cells was inoculated on a CTT agar plate (3×10^7 cells per plate). One line of *S. coelicolor*




Fig. 1. Predatory activity of *M. xanthus* on *Streptomyces*.

A. A CTT agar plate inoculated with a lawn of *M. xanthus* cells was streaked with a line of *S. coelicolor* (Sc) cells and another of *S. lividans* (Sl) cells. Photograph was taken after 120 h of incubation.

B. A CTT plate streaked with a line of *S. coelicolor* (Sc) cells and another of *S. lividans* (Sl) cells.

C. *Streptomyces coelicolor* cells (Sc) were exposed to three *M. xanthus* strains, the wt (DK1622), a mutant leaky in S motility (DZF1) and the non-motile *mgl* mutant (DK6204). Pictures, of the surface of the colonies, were taken at 72 h and should be compared with the Sc/Sc control.

All pictures were taken at the same magnification. Bar represents 2 mm.

M145 and another of *Streptomyces lividans* 1326 were streaked across the *M. xanthus* cells. The plates were incubated at 30°C for 5 days. Production of the blue antibiotic ACT was observed around the *S. coelicolor* M145 cells (Fig. 1A). However, ACT production by *S. coelicolor* M145 was very low in a control plate on which *S. coelicolor* M145 and *S. lividans* 1326 were inoculated alone (Fig. 1B).

Interactions between *M. xanthus* and *S. coelicolor* were examined in more detail by inoculating drops of each microorganism next to each other. Two drops of *Streptomyces* at the same distance were used as a control. Three *M. xanthus* strains were examined, the fully motile wild-type (wt) strain DK1622, the one with reduced S motility DZF1 strain and the non-motile mutant DK6204. Only the two motile strains moved towards *Streptomyces* (Fig. 1C). The *mgl* mutant (DK6204) was unable to migrate towards the *Streptomyces* but it induces coloured antibiotic production and aerial mycelium formation suggesting produc-

tion of a bioactive compound or lytic enzyme. More detailed visualization of predation can be observed in the movie attached as supplementary information.

M. xanthus induces formation of abnormal *S. coelicolor* hyphae

As shown in the movie, *M. xanthus* DK1622 cells moved towards *S. coelicolor* spores lysing them even from some distance. However, *Streptomyces* recovers coincident with spore germination suggesting that growing *Streptomyces* cells are partially resistant (see below). As the *M. xanthus* cells enter the *S. coelicolor* colony intense lysis of *Streptomyces* hyphae occurs. The use of the LIVE/DEAD BacLight kit demonstrates that control cultures of *S. coelicolor* without *Myxococcus* contained mainly living cells (Fig. 2A). However, the co-cultures were predominantly red indicating a high proportion of dead mycelia (Fig. 2A). Scanning electron microscopy showed that the cells at the distal edges of the *Streptomyces* and *Myxococcus* drops that did not have contact with each other looked healthy (Fig. 2B). *Myxococcus* cells in the interaction zone also

appeared healthy having the normal bacillar shape whereas *Streptomyces* hyphae exhibited aberrant morphologies (Fig. 2B). The cell density for both bacterial species in this zone was lower than at the distal edges due to competition and predation.

M. xanthus induces antibiotic production and differentiation in *S. coelicolor*

In *S. coelicolor* colonies growing next to *M. xanthus* more ACT is produced in the interaction zone after 48–72 h of incubation than in distal regions of the spots or with the *S. coelicolor* only controls (Fig. 3). ACT production is upregulated by each of the three different *M. xanthus* strains used in this experiment suggesting that motility is not necessary (Fig. 1C). In addition, aerial mycelia containing grey spores are formed near the interaction zone (Figs 1C and 3). Aerial mycelia were induced by 48 h with each of the three *M. xanthus* strains suggesting that production is not dependent on movement or direct contact between organisms, as illustrated by non-motile mutant DK6204




Fig. 2. Effect of *M. xanthus* predation on *S. coelicolor* cells.

A. Microscopic observation of *S. coelicolor* (Sc) or the interface of the co-culture of *S. coelicolor* and *M. xanthus* DK1622 (Sc/Mx) visualized by phase-contrast or fluorescence microscopy after staining with SYTO 9 and propidium iodide. Bar represents 5 μ m. B. Scanning electron micrographs of *S. coelicolor* and *M. xanthus* DK1622 cells on CTT agar plates. The colour picture shows the macroscopic appearance of the adjacent drops of *M. xanthus* (left) and *S. coelicolor* (right) cells that were spotted next to each other. The white arrows indicate the regions observed by scanning electron microscopy. Blue arrows point to aberrant hyphae and yellow arrows to *M. xanthus* cells. Not all *M. xanthus* cells were marked. Bar represents 1 μ m.




Fig. 3. Induction of ACT in *S. coelicolor*. Sc/Sc and Sc/Mx indicates that either two drops of *S. coelicolor*, or one drop of *S. coelicolor* and another of *M. xanthus* DK1622, respectively, were spotted next to each other. Pictures were taken with a digital camera at the times indicated in each panel. Bar represents 2 mm. Pictures were taken through the bottom of the Petri plate.

(Fig. 1C). If anything, the non-motile mutant seems to stimulate more sporulation than wt *M. xanthus* cells (Fig. 1C).

Production of ACT and aerial mycelia is also stimulated by co-culture with other microorganisms (Table 1). Induction of ACT was observed in co-cultures with *Bacillus megaterium*, *B. subtilis*, *B. thuringensis* and *Serratia*, although to much lower extent than with *M. xanthus*. Some of these bacteria were also able to induce the aerial mycelium formation (Table 1). From organisms such as

Klebsiella pneumoniae, which induces aerial mycelia but not ACT and *B. megaterium*, which induces ACT but not aerial mycelia it would appear that the two processes are not strictly coupled.

ACT overproduction is induced in liquid co-culture of S. coelicolor and M. xanthus

Stimulation of antibiotic production was also studied in CTT liquid cultures. ACT was clearly overproduced by *S. coelicolor* in co-culture with *Myxococcus* strains DK1622 or DZF1 (Fig. 4A). Colorimetric quantification of ACT indicated that the presence of either *M. xanthus* strain increased *Streptomyces* ACT production profusely. However, very low and similar levels of CDA were detected under the conditions assayed in the control or the co-cultures (data not shown).

Ultra high pressure liquid chromatography (UHPLC) allowed quantification of the extracted compounds produced in the control cultures, *S. coelicolor* (Fig. 4B) or *M. xanthus* DK1622 (Fig. 4F), and in the co-culture with both strains (Fig. 4D). Several compounds eluting in the region between 4.5 and 7 min in the *Streptomyces* control culture (Fig. 4B) were overproduced in the co-cultures (Fig. 4D), and absent in the *M. xanthus* DK1622 control (Fig. 4F). These peaks shared the same absorption spectrum, which corresponded to that of the ACT family. When optimized to 500 nm, a wavelength suited to detect ACT, the types of ACT molecules were similar between the *S. coelicolor* control and the co-culture, but the amounts were roughly 20-fold higher in the co-culture (Fig. 4C and E; notice the difference in the scale). These compounds

Table 1. Induction of ACT and aerial mycelia by *S. coelicolor* in co-culture with different bacteria species.

Bacteria	ACT ^a	AM ^b
<i>Myxococcus xanthus</i>	++	++
<i>Bacillus laterosporus</i>	-	-
<i>Bacillus licheniformis</i>	-	-
<i>Bacillus megaterium</i>	+	-
<i>Bacillus subtilis</i>	+	++
<i>Bacillus thuringensis</i>	+	++
<i>Micrococcus</i> sp.	-	-
<i>Mycobacterium phlei</i>	-	-
<i>Mycobacterium smegmatis</i>	-	-
<i>Staphylococcus aureus</i>	-	-
<i>Escherichia coli</i>	-	-
<i>Klebsiella pneumoniae</i>	-	++
<i>Proteus</i> sp.	-	-
<i>Salmonella</i> sp.	-	-
<i>Serratia</i> sp.	+	++


a. ACT indicates actinorhodin production.

b. AM indicates the development of aerial mycelia.

++ and + indicate actinorhodin production or aerial mycelia development after 48 and 72 h, respectively, and - indicates that actinorhodin or aerial mycelia were observed at the same time as the control which contains only *S. coelicolor* cells.



Fig. 4. Induction of ACT in *S. coelicolor* by *M. xanthus* in liquid CTT cultures. Flasks containing cultures with only *S. coelicolor* wt, or *S. coelicolor* plus *M. xanthus* DK1622 or DZF1 are shown in (A). The chromatograms of extracts of *S. coelicolor*, *M. xanthus* DK1622 or a co-culture of both strains are shown in (B), (F) and (D) respectively. The chromatograms are maxplots (i.e. chromatograms at absorbance maximum for each analyte) obtained from spectrophotometric detection in the range from 200 to 500 nm. In (C) (*S. coelicolor* wt), (E) (co-culture) and (G) (*M. xanthus* DK1622), data were processed to obtain chromatograms at 500 nm, a wavelength suited to observe the production of ACT. Note the differences in scales between graphs.



were not detected in the *M. xanthus* control culture (Fig. 4G). HPLC-MS analysis of the same peaks gave $m/z[H^+]$ values ranging from 631 to 666, as expected for ACT family members (data not shown). RED was not detected in the control culture or in the co-culture. These experiments confirm that *M. xanthus* increases production of ACT by *S. coelicolor*.

S. coelicolor ACT biosynthesis mutants repel *M. xanthus* less effectively

To determine whether ACT production provides an advantage to *S. coelicolor* strains against *M. xanthus* predation, mutants impaired in the production of RED (M510), ACT (M511) or both antibiotics (M512) were used. ACT was produced by strains M145 (wild type) and M510 (Fig. 5).

RED was detected in M145 and M511 (Fig. 5). Migration of *M. xanthus* DK1622 cells towards all the *Streptomyces* strains was observed. However, this migration was more evident with the *Streptomyces* strains that did not produce ACT (M511 and M512 strains). *Myxococcus xanthus* seemed to more aggressively attack strains lacking ACT (see the blue arrows in Fig. 5 taken after 200 h co-culture). This observation predicted that strain M512 would be more sensitive to *M. xanthus* attack. However, the general appearance of the *Streptomyces* colonies suggested that the three mutants were as resistant to *M. xanthus* predation as the wt strain.

To obtain a clearer understanding predation was quantified by dilution plating. *Myxococcus xanthus* DK1622 was co-cultured in liquid CTT medium with the *S. coelicolor* strains M145 or M512. After incubation for 3 days,




Fig. 5. Analysis of the interaction between different mutants of *S. coelicolor* and *M. xanthus* DK1622 on CTT agar plates. Blue arrows point to *M. xanthus* cells moving towards *S. coelicolor*. Bars represent 2 mm.

cells were diluted and inoculated onto R2YE plates (*Myxococcus* does not grow in this medium). The number of *S. coelicolor* colonies that survive co-culture with *M. xanthus* was similar for both strains, approximately one-fifth of the colonies obtained when the *Streptomyces* strains were grown in the absence of *M. xanthus*. This result demonstrates that the double antibiotic mutant has a similar level of resistance to *Myxococcus* predation.

Discussion

Genome sequencing efforts have revealed that, under laboratory conditions, microorganisms have a wide number of genes that remains silent. Recently co-culture of two microorganisms has been considered as a strategy to partially mimic natural communities that exchange chemical signals. This new approach has permitted the discovery of new capabilities that remained silent in axenic cultures. For example, previous work on *Streptomyces* interactions has demonstrated alteration of its developmental program due to availability of new carbon sources during co-culture of *S. lividans* with yeasts (Santamaria *et al.*, 2002). *Bacillus subtilis* production of a surfactant inhibits chaplin and SapB production, which are required for aerial mycelium formation and sporulation in *S. coelicolor* (Straight *et al.*, 2006). Description of the interaction of *Streptomyces olivaceoviridis* with *Aspergil-*

lus proliferans via a protein targeting chitin on the fungus cell wall has been done (Siemieniowicz and Schrempf, 2007). Direct physical interaction between *Streptomyces hygroscopicus* and the fungus *Aspergillus nidulans* is necessary for induction of polyketide synthesis. In this interaction the bacterium also triggers the production of lecanoric acid by the fungus, a metabolite that inhibits ATP synthesis and may be used in self-defence by the fungus (Schroeckh *et al.*, 2009).

Myxobacteria and actinomycetes are normal inhabitants of soil where they act as scavengers to recycle cellular debris using hydrolytic enzymes. Both types of organisms produce molecules with antibiotic activity that may act in defence or as communication signals. *Myxococcus xanthus*, the model myxobacterium, behaves as an active predator, able to consume other microorganisms and even worms. To do so, myxobacterial cells detect potential preys and surround them to facilitate predation. Non-motile *Streptomyces*, the most abundant actinomycete in soil, produces dense colonies of mycelia in which the position of cells in the colony determine the pattern of gene expression. Although *Streptomyces* is not a motile predator, it lyses other organisms by secreting antibiotics and hydrolytic enzymes. In the soil, both *Streptomyces* and *Myxococcus* coexist and there is evidence for horizontal gene transfer between these bacteria in the case of an endoglucanase (CelA) gene transferred from

Streptomyces to *Myxococcus* ancestors (Quillet *et al.*, 1995). Horizontal transfer between Gram-positive and Gram-negative bacteria is widely represented in other genes and could originate with predation and lead to the incorporation of prey DNA in the genome of the predator.

Interspecies signalling has the potential to induce silent metabolic pathways or to obtain new hybrid compounds. Genome sequencing projects have revealed the capacity of the organisms to produce small molecules that are not produced under laboratory conditions. Some conclusions may be extracted from the study reported here, where *Myxococcus* and *Streptomyces* have been co-cultured. *Myxococcus xanthus* stimulates *S. coelicolor* production of the blue polyketide antibiotic ACT but not the tripyrrol antibiotic RED or the cyclic lipopeptide CDA. Theoretically, ACT does not have antibiotic activity on Gram-negative bacteria like *Myxococcus* because it is not able to enter in the cell bind DNA. However, *S. coelicolor* strains that produce ACT are not surrounded by *M. xanthus* in the same manner as those that do not synthesize this antibiotic. It remains possible that *S. coelicolor* uses ACT as a repellent signal for *Myxococcus*. In addition, antibiotic production in *S. coelicolor* by a signal produced by other bacteria might have important biotechnological applications to improve the yield of clinically relevant antibiotics.

Co-culture also stimulates aerial mycelium formation by *S. coelicolor* suggesting a chemical induction pathway. It could be argued that nutrient depletion by *M. xanthus* may lead to more rapid development of *S. coelicolor*. However, co-cultures of *M. xanthus* and *S. lividans* do not induce aerial mycelia in this bacteria (data not shown). In addition, aerial mycelia are not observed when *S. coelicolor* colonies are juxtaposed or confronted with other bacterial strains. These results suggest that a signal produced by *Myxococcus* is recognized by receptor encoded by *S. coelicolor* but not *S. lividans*. It will be of interest to determine whether the *M. xanthus* molecule that induces ACT production in *S. coelicolor* is the same as the one that stimulates differentiation.

Experimental procedures

Bacterial strains and media

Streptomyces coelicolor M145 and *S. lividans* 66 (Kieser *et al.*, 2000) were used as prey along with mutant derivatives of *S. coelicolor* M145: M510 ($\Delta redD$), M511 ($\Delta actII-ORF4$) and M512 ($\Delta redD-\Delta actII-ORF4$) (Floriano and Bibb, 1996) that do not produce the antibiotics RED, ACT or both respectively. The wt *M. xanthus* DK1622 (Kaiser, 1979), the *pilQ1* mutant DZF1, leaky in S motility (Morrison and Zusman, 1979), and the non-motile *mgI* mutant DK6204 (Hartzell and Kaiser, 1991) were used as predators. CTT solid (1.5% Bacto agar) and liquid media were used to grow *M. xanthus* (Hodgkin and Kaiser, 1977). R2YE was used for *Streptomy-*

ces cultures (Kieser *et al.*, 2000). Several other bacteria have been used to examine *S. coelicolor* antibiotic production and aerial mycelium formation. All of them were grown in Luria-Bertani medium (Sambrook and Russell, 2001). These bacteria were obtained from the 'Colección del Departamento de Microbiología' (Universidad de Granada, Spain) (*Bacillus laterosporus*, *B. licheniformis*, *B. megaterium*, *B. subtilis*, *B. thuringensis*, *Mycobacterium smegmatis*, *K. pneumoniae*, *Salmonella* sp. and *Serratia* sp.), and from the 'Colección Española de Cultivos Tipo' (CECT) [*Micrococcus* sp. (CECT 241), *Mycobacterium phlei* (CECT 3009), *Staphylococcus aureus* (CECT 240), *E. coli* (CECT 101) and *Proteus* sp. (CECT 484)].

Predation experiments

Myxococcus xanthus strains were grown in CTT for 24 h, washed twice with sterile TM buffer (10 mM Tris-HCl, pH 7.6, 1 mM MgSO₄) and concentrated to a final cell density of 4.5×10^9 cells ml⁻¹. Drops of 5 or 10 μ l were deposited on the surface of CTT agar plates and allowed to dry. Next, drops of 5 or 10 μ l of *Streptomyces* spores (2×10^8 spores ml⁻¹) were spotted close to the *Myxococcus* spot to leave a separation of no more than 1 mm. Plates were incubated at 30°C and images were taken directly with a digital camera or under a Zeiss Stemi SV11 or Wild-Heerbrugg dissecting microscope. Each experiment was repeated at least four times. The same approach was used when *S. coelicolor* was plated next to other bacterial and yeast species.

Liquid co-cultures and antibiotic quantification

Co-cultures of *S. coelicolor* with *M. xanthus* were carried out in liquid CTT. A total of 10^7 *S. coelicolor* spores were inoculated into 100 ml baffled flasks containing 10 ml of CTT and incubated at 28°C for 24 h. In parallel a culture of *M. xanthus* DK1622 was incubated under identical conditions. Different amounts of *M. xanthus* DK1622 (from 3×10^7 to 3×10^4 cells ml⁻¹) were added to the *S. coelicolor* cultures, except to the control where only *Streptomyces* was grown. Incubation was continued for 3–5 days. Production of the coloured antibiotics was quantified by colorimetric assays (Kieser *et al.*, 2000) and production of CDA was determined by bioassay using *B. subtilis* as the sensitive organism.

Chromatographic analysis

Culture supernatants from 10 ml of liquid cultures were extracted twice with an equal volume of ethyl acetate containing 1% formic acid. The solvent was evaporated and the residue redissolved in 100 μ l of dimethyl sulfoxide : methanol (50:50). These samples were fractionated by reversed phase in an Acquity UPLC with a BEH C18 column (1.7 μ m, 2.1×100 mm, Waters) using acetonitrile and 0.1% trifluoroacetic acid in water. Samples were eluted with 10% acetonitrile for 1 min, followed by a linear gradient from 10% to 100% over 15 min at a flow rate of 0.5 ml min⁻¹ and a column temperature of 30°C. For HPLC-MS analysis, an Alliance chromatographic system coupled to a ZQ4000 mass spectrometer and a Symmetry C18 (2.1×150 mm, Waters) was

used. Solvents were the same as above and elution was performed with an initial isocratic hold with 10% acetonitrile for 4 min followed by a linear gradient from 10% to 88% acetonitrile over 26 min, at 0.25 ml min⁻¹. MS analysis were performed by electrospray ionization in the positive mode, with a capillary voltage of 3 kV and a cone voltage of 20 V. Detection and spectral characterization of peaks was performed in both cases by photodiode array detection and Empower software (Waters).

Fluorescence microscopy

Cellular viability was detected by using the LIVE/DEAD Bac Light Bacterial Viability kit L-13152 (Molecular Probes), which includes the dyes SYTO 9 (Green) and propidium iodide (red). SYTO 9 stains living cells green while propidium iodide stains damaged cells red (Haugland, 2002). The kit was used as indicated by the manufacturer to stain control cultures of *Streptomyces* or co-cultures of both organisms. Samples were observed in a Leica DMRXA microscope equipped for bright-field and epifluorescence and photographed with an Orca-ER C4742-80 camera (Hamamatsu, Bridgewater, NJ).

Scanning electron microscopy

Approximately 72 h co-cultures of *S. coelicolor* and *M. xanthus* on CTT agar plates were fixed with 3% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.4 for 24 h at 4°C. Then, samples were washed three times (5 min each) with the same buffer. Dehydration was accomplished by a graded series of ethanol. Samples were then critical-point dried and sputter coated with carbon. Photographs were taken in a LEO 1530 scanning electron microscope.

Videomicroscopy

Cell spots of a co-culture of *S. coelicolor* and *M. xanthus* were filmed with a Wild Heerbrugg M7 S dissecting microscope at room temperature. Photographs were taken every 5 min as jpg files with a Spot Insight 2 camera using SPOT software v4.5 (Diagnostic Instruments). The movies were compiled from the images using Quicktime Pro (Apple) at six frames per second. The field of view is 3.8 mm across. The movie was compressed using the H.264 video codec in QuickTime Pro 7.

Acknowledgements

This work has been supported by grants from Ministerio de Ciencia e Innovación (EUI2008-03631 to R.I.S.; BFU2009-07565/BMC, 70% funded by FEDER, and the programme CONSOLIDER-INGENIO 2010, CSD2009-00006, both to J.M.-D.), Spain. This material is based on work supported in part by the National Science Foundation under Grant 0742976 to L.J.S. Thanks are due to Dr. M. Bibb for *S. coelicolor* M510, M511 and M512 strains and to M.J. Jiménez Rufo for her excellent technical work.

References

Bassler, B.L., and Losick, R. (2006) Bacterially speaking. *Cell* **125**: 237–246.

- Bentley, S.D., Chater, K.F., Cerdano-Tarraga, A.M., Challis, G.L., Thomson, N.R., James, K.D., *et al.* (2002) Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* **417**: 141–147.
- Berleman, J.E., and Kirby, J.R. (2009) Deciphering the hunting strategy of a bacterial wolfpack. *FEMS Microbiol Rev* **33**: 942–957.
- Berleman, J.E., Chumley, T., Cheung, P., and Kirby, J.R. (2006) Rippling is a predatory behavior in *Myxococcus xanthus*. *J Bacteriol* **188**: 5888–5895.
- Capstick, D.S., Willey, J.M., Buttner, M.J., and Elliot, M.A. (2007) SapB and the chaplins: connections between morphogenetic proteins in *Streptomyces coelicolor*. *Mol Microbiol* **64**: 602–613.
- Chater, K.F. (1993) Genetics of differentiation in *Streptomyces*. *Annu Rev Microbiol* **47**: 685–713.
- Chater, K.F., Biro, S., Lee, K.J., Palmer, T., and Schrepf, H. (2010) The complex extracellular biology of *Streptomyces*. *FEMS Microbiol Rev* **34**: 171–198.
- Dworkin, M. (1996) Recent advances in the social and developmental biology of the myxobacteria. *Microbiol Rev* **60**: 70–102.
- Flardh, K., and Buttner, M.J. (2009) *Streptomyces* morphogenetics: dissecting differentiation in a filamentous bacterium. *Nat Rev Microbiol* **7**: 36–49.
- Floriano, B., and Bibb, M. (1996) *afsR* is a pleiotropic but conditionally required regulatory gene for antibiotic production in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **21**: 385–396.
- Hartzell, P., and Kaiser, D. (1991) Upstream gene of the *mgl* operon controls the level of MglA protein in *Myxococcus xanthus*. *J Bacteriol* **173**: 7625–7635.
- Haugland, R.P. (2002) *Handbook of Fluorescent Probes and Research Products*, 9th ed. Probes, G.J.M. (ed.). Eugene, OR, USA: Molecular Probes.
- Hillesland, K.L., Lenski, R.E., and Velicer, G.J. (2007) Ecological variables affecting predatory success in *Myxococcus xanthus*. *Microb Ecol* **53**: 571–578.
- Hodgkin, J., and Kaiser, D. (1977) Cell-to-cell stimulation of movement in nonmotile mutants of *Myxococcus*. *Proc Natl Acad Sci USA* **74**: 2938–2942.
- Kaiser, D. (1979) Social gliding is correlated with the presence of pili in *Myxococcus xanthus*. *Proc Natl Acad Sci USA* **76**: 5952–5956.
- Kieser, T., Bibb, M.J., Buttner, M.J., Chater, K.F., and Hopwood, D.A. (2000) *Practical Streptomyces genetics*. Norwich, UK: The John Innes Foundation.
- Kroos, L. (2007) The *Bacillus* and *Myxococcus* developmental networks and their transcriptional regulators. *Annu Rev Genet* **41**: 13–39.
- Kurosawa, K., Ghiviriga, I., Sambandan, T.G., Lessard, P.A., Barbara, J.E., Rha, C., and Sinskey, A.J. (2008) Rhodostreptomycins, antibiotics biosynthesized following horizontal gene transfer from *Streptomyces padanus* to *Rhodococcus fascians*. *J Am Chem Soc* **130**: 1126–1127.
- Mauriello, E.M., and Zusman, D.R. (2007) Polarity of motility systems in *Myxococcus xanthus*. *Curr Opin Microbiol* **10**: 624–629.
- Mazzola, M. (2007) Manipulation of rhizosphere bacterial communities to induce suppressive soils. *J Nematol* **39**: 213–220.

- Morrison, C.E., and Zusman, D.R. (1979) *Myxococcus xanthus* mutants with temperature-sensitive, stage-specific defects: evidence for independent pathways in development. *J Bacteriol* **140**: 1036–1042.
- Quillet, L., Barray, S., Labedan, B., Petit, F., and Guespin-Michel, J. (1995) The gene encoding the beta-1,4-endoglucanase (CelA) from *Myxococcus xanthus*: evidence for independent acquisition by horizontal transfer of binding and catalytic domains from actinomycetes. *Gene* **158**: 23–29.
- Sambrook, J., and Russell, D. (2001) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor, NY, USA: Cold Spring Harbor Laboratory.
- Santamaria, R.I., Leal, F., Diaz, M., and Fernandez-Abalos, J.M. (2002) Morphological and physiological changes in *Streptomyces lividans* induced by different yeasts. *Arch Microbiol* **177**: 259–266.
- Schneiker, S., Perlova, O., Kaiser, O., Gerth, K., Alici, A., Altmeyer, M.O., et al. (2007) Complete genome sequence of the myxobacterium *Sorangium cellulosum*. *Nat Biotechnol* **25**: 1281–1289.
- Schroeckh, V., Scherlach, K., Nutzmann, H.W., Shelest, E., Schmidt-Heck, W., Schuemann, J., et al. (2009) Intimate bacterial–fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proc Natl Acad Sci USA* **106**: 14558–14563.
- Shank, E.A., and Kolter, R. (2009) New developments in microbial interspecies signaling. *Curr Opin Microbiol* **12**: 205–214.
- Siemieniowicz, K.W., and Schrempf, H. (2007) Concerted responses between the chitin-binding protein secreting *Streptomyces olivaceoviridis* and *Aspergillus proliferans*. *Microbiology* **153**: 593–600.
- Straight, P.D., and Kolter, R. (2009) Interspecies chemical communication in bacterial development. *Annu Rev Microbiol* **63**: 99–118.
- Straight, P.D., Willey, J.M., and Kolter, R. (2006) Interactions between *Streptomyces coelicolor* and *Bacillus subtilis*: role of surfactants in raising aerial structures. *J Bacteriol* **188**: 4918–4925.
- Straight, P.D., Fischbach, M.A., Walsh, C.T., Rudner, D.Z., and Kolter, R. (2007) A singular enzymatic megacomplex from *Bacillus subtilis*. *Proc Natl Acad Sci USA* **104**: 305–310.
- Tamilarasi, S., Nanthakumar, K., Karthikeyan, K., and Lakshmanaperumalsamy, P. (2008) Diversity of root associated microorganisms of selected medicinal plants and influence of rhizomicroorganisms on the antimicrobial property of *Coriandrum sativum*. *J Environ Biol* **29**: 127–134.
- Velicer, G.J., and Mendes-Soares, H. (2009a) Bacterial predators. *Curr Biol* **19**: R55–R56.
- Velicer, G.J., and Vos, M. (2009b) Sociobiology of the myxobacteria. *Annu Rev Microbiol* **63**: 599–623.
- Vos, M., and Velicer, G.J. (2009) Social conflict in centimeter- and global-scale populations of the bacterium *Myxococcus xanthus*. *Curr Biol* **19**: 1763–1767.
- Wenzel, S.C., and Muller, R. (2009) Myxobacteria ‘microbial factories’ for the production of bioactive secondary metabolites. *Mol Biosyst* **5**: 567–574.
- Yamanaka, K., Oikawa, H., Ogawa, H.O., Hosono, K., Shinmachi, F., Takano, H., et al. (2005) Desferrioxamine E produced by *Streptomyces griseus* stimulates growth and development of *Streptomyces tanashiensis*. *Microbiology* **151**: 2899–2905.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Movie S1. Interaction between *M. xanthus* (left) and *S. coelicolor* (right) on solid CTT medium. The pictures were taken every 5 minutes and compiled with Quicktime Pro (Apple).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.